

# Effects of Free Fatty Acid Elevation on Postabsorptive Endogenous Glucose Production and Gluconeogenesis in Humans

Michael Roden, Harald Stingl, Visvanathan Chandramouli, William C. Schumann, Astrid Hofer, Bernard R. Landau, Peter Nowotny, Werner Waldhäusl, and Gerald I. Shulman

**Effects of free fatty acids (FFAs) on endogenous glucose production (EGP) and gluconeogenesis (GNG) were examined in healthy subjects ( $n = 6$ ) during stepwise increased Intralipid/heparin infusion (plasma FFAs  $0.8 \pm 0.1$ ,  $1.8 \pm 0.2$ , and  $2.8 \pm 0.3$  mmol/l) and during glycerol infusion (plasma FFAs  $\sim 0.5$  mmol/l). Rates of EGP were determined with D-[6,6- $^2\text{H}_2$ ]glucose and contributions of GNG from  $^2\text{H}$  enrichments in carbons 2 and 5 of blood glucose after  $^2\text{H}_2\text{O}$  ingestion. Plasma glucose concentrations decreased by  $\sim 10\%$  ( $P < 0.01$ ), whereas plasma insulin increased by  $\sim 47\%$  ( $P = 0.02$ ) after 9 h of lipid infusion. EGP declined from  $9.3 \pm 0.5$  (lipid) and  $9.0 \pm 0.8$   $\mu\text{mol} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$  (glycerol) to  $8.4 \pm 0.5$  and  $8.2 \pm 0.7$   $\mu\text{mol} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ , respectively ( $P < 0.01$ ). Contribution of GNG similarly rose ( $P < 0.01$ ) from  $46 \pm 4$  and  $52 \pm 3\%$  to  $65 \pm 8$  and  $78 \pm 7\%$ . To exclude interaction of FFAs with insulin secretion, the study was repeated at fasting plasma insulin ( $\sim 35$  pmol/l) and glucagon ( $\sim 90$  ng/ml) concentrations using somatostatin-insulin-glucagon clamps. Plasma glucose increased by  $\sim 50\%$  ( $P < 0.005$ ) during lipid but decreased by  $\sim 12\%$  during glycerol infusion ( $P < 0.005$ ). EGP remained unchanged over the 9-h period ( $9.9 \pm 1.2$  vs.  $9.0 \pm 1.1$   $\mu\text{mol} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ ). GNG accounted for  $62 \pm 5$  (lipid) and  $60 \pm 6\%$  (glycerol) of EGP at time 0 and rose to  $74 \pm 3\%$  during lipid infusion only ( $P < 0.05$  vs. glycerol:  $64 \pm 4\%$ ). In conclusion, high plasma FFA concentrations increase the percent contribution of GNG to EGP and may contribute to increased rates of GNG in patients with type 2 diabetes. *Diabetes* 49:701–707, 2000**

From the Division of Endocrinology and Metabolism (M.R., H.S., A.H., P.N., W.W.), Department of Internal Medicine III, University of Vienna Medical School, Vienna, Austria; the Departments of Medicine (V.C., W.C.S., B.R.L.) and Biochemistry (B.R.L.), Case Western Reserve University, Cleveland, Ohio; and the Department of Internal Medicine (G.I.S.) and Howard Hughes Medical Institute (G.I.S.), Yale University School of Medicine, New Haven, Connecticut.

Address correspondence and reprint requests to Michael Roden, MD, Division of Endocrinology and Metabolism, Department of Internal Medicine III, University of Vienna Medical School, General Hospital of Vienna, Währinger Gürtel 18-20, A-1090 Vienna, Austria. E-mail: michael.roden@akh-wien.ac.at.

Received for publication 1 September 1999 and accepted in revised form 21 January 2000.

CV, coefficient of variation; EGP, endogenous glucose production; FFA, free fatty acid; GNG, gluconeogenesis; HMT, hexamethylenetetramine.

**E**levation of plasma free fatty acid (FFA) concentrations is often associated with obesity (1) and type 2 diabetes (2). The close correlation between whole-body glucose uptake and fasting plasma FFA concentrations in lean normoglycemic offspring of type 2 diabetic parents (3) indicates that FFAs might play a pivotal role in the early events leading to insulin resistance (4,5).

Plasma FFA elevation induced by lipid/heparin infusions during hyperinsulinemic clamps has repeatedly been shown to decrease insulin-dependent whole-body glucose disposal (5–8). Reports on a correlation between plasma FFAs and hepatic insulin sensitivity are more controversial. Fasting plasma FFAs correlate with the magnitude of hyperglycemia and endogenous glucose production (EGP) (9), which has been attributed to increased lipid oxidation in type 2 diabetes (10). Under hyperinsulinemic conditions, lipid/heparin infusion either increased (6,11) or had no effect on (12–14) EGP. At postabsorptive plasma insulin concentrations, plasma FFA elevation caused marked increases in EGP during somatostatin-insulin clamps (12,15), but not after an overnight fast (15,16). Similarly, inhibition of lipolysis by nicotinic acid or its derivative, acipimox, decreased basal EGP in some (17,18) but not other (19,20) studies. These apparent discrepancies could result from FFA-induced insulin secretion counterbalancing the stimulatory effect of FFAs on EGP (15) or from hepatic autoregulation preventing an increase in EGP under conditions that might favor hepatic gluconeogenesis (GNG) (16). Of note, increased GNG was documented in type 2 diabetes from a variety of precursors (21,22), whereas contradictory effects of FFAs on the contribution of GNG to EGP have been reported during lipid/heparin infusion or acipimox studies (16,18,20,23).

In most studies, glycerol was not infused during control experiments to match the lipid-induced rise in plasma glycerol concentrations. Thus, stimulation of hepatic GNG by FFAs could also be exclusively due to increased availability of glycerol (12), possibly the major gluconeogenic substrate under these conditions (24). A wide range of estimates of the contribution of GNG to EGP has been reported in humans; however, the methods used to make those estimates have several limitations (25–27). Recently, a method was introduced to overcome those limitations (28,29). Quantitative estimates of the contributions of GNG to EGP are made from  $^2\text{H}$  enrichments in the hydrogens bound to carbons 2 and 5 of glucose on deuterated water ( $^2\text{H}_2\text{O}$ ) ingestion.

The present study was therefore designed to compare concentration-dependent effects of FFAs with those of glycerol on 1) the time-course of plasma glucose, insulin, and C-peptide concentrations; 2) the rates of EGP; and 3) the contribution of endogenous GNG to EGP. The study combined oral administration of  $^2\text{H}_2\text{O}$  with bolus/continuous infusion of dideuterated glucose ( $\text{D-[6,6-}^2\text{H}_2\text{]}_2\text{glucose}$ ) under conditions of prolonged fasting.

## RESEARCH DESIGN AND METHODS

**Subjects.** For the basal study, 6 healthy volunteers (3 men and 3 women, ages 25–37 years, BMI  $22.8 \pm 1.2 \text{ kg/m}^2$ ) without family history of diabetes, dyslipidemia, or bleeding disorders fasted overnight for 8 h before the beginning of the experiments. For the pancreatic clamp study, 6 healthy subjects (5 men and 1 woman, ages 26–38 years, BMI  $22.7 \pm 0.7 \text{ kg/m}^2$ ) were studied after identical overnight fasts; 3 of them had participated in the first study. Female subjects were studied in the follicular phase of the menstrual cycle. The studies were spaced by intervals of 6–13 weeks. No changes in diet, weight, or lifestyle were recorded from the time of recruitment until completion of both studies. The protocols were reviewed and approved by the local human ethics board, and informed consent was obtained from all subjects.

**Study protocols.** For the basal study, experiments were begun at 5:30 A.M. (time -390 min) with insertion of Teflon catheters in antecubital veins of the right and left arm for blood sampling and infusions, respectively. At -375 min, to assess endogenous GNG, subjects started to drink a total of 5 g  $^2\text{H}_2\text{O}$  (99.9%  $^2\text{H}$ , Cambridge Isotope Laboratories, Andover, MA) per kilogram body water divided into 4 equal doses spaced at intervals of 45 min (30). Body water was assumed to be 50% of body weight in women and 60% in men. Thereafter, subjects had free access to water containing 0.5% of  $^2\text{H}_2\text{O}$  to maintain isotopic equilibrium in body water. To allow calculation of EGP, a bolus ( $[6.606 \mu\text{mol/kg} \times \text{body weight in kg} \times \text{fasting blood glucose in } \mu\text{mol/l}]/500 \mu\text{mol/l}$ )-continuous ( $0.067 \mu\text{mol/kg} \times \text{body weight in kg/min}$ ) infusion of  $\text{D-[6,6-}^2\text{H}_2\text{]}_2\text{glucose}$  (99%, Cambridge Isotope Laboratories) was commenced at -210 min (31). Plasma FFA concentrations were raised stepwise every 3 h by infusion of a triglyceride emulsion (0.5, 1.0, and 1.5 ml/min) (Intralipid 20%; Kabi Pharmacia, Uppsala, Sweden) combined with heparin (bolus 200 U; continuous  $0.2 \text{ U} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ ). During control experiments, glycerol was infused from 120 to 540 min ( $0.7 \text{ mg} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$  in 0.9% saline) to control for glycerol effects per se.

For the pancreatic clamp study, the protocol was identical to that in the basal study except 1) somatostatin ( $0.1 \mu\text{g} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ ) was infused from -10 min until 540 min to inhibit hormone secretion, 2) insulin ( $0.07 \text{ mU} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ ) and glucagon ( $0.65 \text{ ng} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ ) were infused to maintain postabsorptive plasma concentrations from 0 to 540 min, and 3) the glycerol infusion was increased stepwise to mimic the increase in plasma glycerol during lipid infusion (135–240 min:  $0.23$ ; 240–360 min:  $0.47$ ; 360–540:  $1.5 \text{ mg} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ ). During both studies, blood samples were taken at timed intervals, immediately

chilled, and centrifuged, and the supernatants were stored at  $-80^\circ\text{C}$  until determination of hormones and metabolites.

**Analytical procedures.** Plasma glucose concentrations were measured using glucose oxidase (Glucose Analyzer II; Beckman, Fullerton, CA). Plasma concentrations of FFAs (intra- and interassay coefficients of variation [CVs] 4.3 and 5.7%), glycerol (CVs 2.0 and 3.4%), and lactate (CVs 3.3%) were determined by enzymatic methods (FFAs: Wako, Neuss, Germany; glycerol: Boehringer-Mannheim, Mannheim, Germany; lactate: Sigma, St. Louis, MO). Plasma insulin (CVs  $\leq 8\%$ ), C-peptide (CVs  $\leq 9\%$ ), glucagon (CVs  $\leq 8\%$ ), and growth hormone (CVs  $\leq 0.7\%$ ) were measured by radioimmunoassay (RIA) (insulin: Pharmacia-Upjohn, Uppsala, Sweden; C-peptide: CIS, Gif-sur-Yvette, France; glucagon: Serono Diagnostics, Freiburg, Germany; growth hormone: Sorin Biomedica, Saluggia, Italy). Plasma cortisol (CVs  $\leq 6\%$ ) was determined after extraction and charcoal-dextran separation by RIA (32).

Enrichments of  $^2\text{H}$  in hydrogens bound to carbon 2 (C2), C5, and C6 of blood glucose were determined using gas chromatography-mass spectrometry by isolating the hydrogens in formaldehyde derivatized to hexamethylenetetramine (HMT) as previously detailed (28–30). Briefly, blood was diluted with an equal volume of water, deproteinized with  $\text{ZnSO}_4/\text{Ba}(\text{OH})_2$ , and deionized, and the glucose was isolated by high-performance liquid chromatography. To estimate fractional GNG, aliquots of glucose were converted to arabitol-5-phosphate and ribitol-5-phosphate and to xylose, which were oxidized with periodate to yield formaldehyde containing C2 or C5 of glucose, respectively, with their hydrogens. For assessing EGP, the hydrogens bound to C6 of glucose were isolated in formaldehyde by direct oxidation with periodate of an aliquot of glucose. Each formaldehyde was condensed with ammonia to form HMT. The HMTs were assayed for  $^2\text{H}$  enrichments by mass spectrometry (28,29), mass 141 for the enrichment of the hydrogen at carbons C2 and C5 and mass 142 for those at C6. Enrichments in plasma water were determined by Metabolic Solutions (Merrimack, NH) using an isotope ratio mass spectrometer.

**Calculations and data analysis.** Rates of EGP were calculated from the tracer infusion rate of  $\text{D-[6,6-}^2\text{H}_2\text{]}_2\text{glucose}$  and its enrichment divided by the mean percent enrichment of plasma  $\text{D-[6,6-}^2\text{H}_2\text{]}_2\text{glucose}$  for the 3 collections minus the infusion rate. The percent contribution of GNG to EGP was set to 100 times the ratio of  $^2\text{H}$  enrichment bound at C5 to that at C2 of blood glucose. The absolute contribution of GNG to EGP was calculated by multiplying percent GNG by EGP. All data are presented as means  $\pm$  SE. Statistical comparisons between lipid and glycerol experiments were performed using the paired Student's *t* test.

## RESULTS

**Basal study.** Lipid infusion at rates of 0.5, 1.0, and 1.5 ml/min resulted in a stepwise increase in plasma FFA concentrations, which did not change from baseline values during glycerol (control) infusion (Table 1). Plasma glycerol concentrations continuously increased, being higher ( $P < 0.01$ ) during

TABLE 1  
Basal study: plasma concentrations of FFAs, glycerol, lactate, growth hormone, and cortisol during lipid infusion and glycerol infusion (control)

|                               | 0 min           | 150–180 min                | 330–360 min                 | 510–540 min                 |
|-------------------------------|-----------------|----------------------------|-----------------------------|-----------------------------|
| FFAs (mmol/l)                 |                 |                            |                             |                             |
| Lipid                         | $0.48 \pm 0.07$ | $0.81 \pm 0.09^{*\dagger}$ | $1.82 \pm 0.18^{*\ddagger}$ | $2.76 \pm 0.29^{*\ddagger}$ |
| Control                       | $0.35 \pm 0.09$ | $0.52 \pm 0.08$            | $0.50 \pm 0.10$             | $0.59 \pm 0.08$             |
| Glycerol (mmol/l)             |                 |                            |                             |                             |
| Lipid                         | $0.10 \pm 0.03$ | $0.18 \pm 0.02^{*\dagger}$ | $0.28 \pm 0.03^{\S\dagger}$ | $0.46 \pm 0.06^{\ddagger}$  |
| Control                       | $0.14 \pm 0.02$ | $0.36 \pm 0.03^{\ddagger}$ | $0.47 \pm 0.04^{\ddagger}$  | $0.48 \pm 0.04^{\ddagger}$  |
| Lactate (mmol/l)              |                 |                            |                             |                             |
| Lipid                         | $0.52 \pm 0.05$ | $0.42 \pm 0.07$            | $0.42 \pm 0.06$             | $0.42 \pm 0.02$             |
| Control                       | $0.40 \pm 0.06$ | $0.50 \pm 0.07$            | $0.43 \pm 0.05$             | $0.46 \pm 0.06$             |
| Growth hormone (ng/ml)        |                 |                            |                             |                             |
| Lipid                         | $4.5 \pm 2.1$   | $6.3 \pm 2.9$              | $1.5 \pm 0.9$               | $1.8 \pm 1.1$               |
| Control                       | $2.3 \pm 0.8$   | $5.2 \pm 2.6$              | $4.2 \pm 1.7$               | $4.8 \pm 1.8$               |
| Cortisol ( $\mu\text{g/dl}$ ) |                 |                            |                             |                             |
| Lipid                         | $9.3 \pm 1.6$   | $8.0 \pm 1.2$              | $5.8 \pm 1.1^{\ddagger}$    | $2.3 \pm 0.6^{\ddagger}$    |
| Control                       | $8.0 \pm 1.5$   | $7.2 \pm 1.6$              | $5.7 \pm 1.3$               | $2.8 \pm 0.5^{\ddagger}$    |

Data are means  $\pm$  SE of 6 paired experiments. \* $P < 0.0005$ ;  $\dagger P < 0.05$ ;  $\ddagger P < 0.005$  vs. basal;  $\S P < 0.005$  vs. control.




FIG. 1. Basal study: plasma concentrations of glucose, insulin, C-peptide, and glucagon during lipid infusion (●) and glycerol infusion (control; ○). Data are given as means  $\pm$  SE of 6 paired experiments. \* $P < 0.01$  vs. control; § $P < 0.01$  vs. 0 min.

glycerol infusion until 360 min, but were not different at 510–540 min. Plasma cortisol decreased ( $P < 0.05$ ) in both experiments, whereas plasma lactate and growth hormone did not change.

Plasma glucose concentrations remained constant during the first 3 h of both parts of the basal study (Fig. 1). Then plasma glucose slightly declined, by  $\sim 7\%$ , during the final period of lipid infusion ( $P < 0.00001$  vs. time 0), but not during glycerol infusion. Plasma insulin concentrations continuously increased during lipid infusion and were  $\sim 48\%$  higher than during glycerol infusion at 510–540 min ( $P < 0.05$ ). In parallel, plasma C-peptide concentrations increased by  $\sim 50\%$  during lipid infusion ( $P < 0.05$  vs. baseline) but decreased by  $\sim 30\%$  during glycerol infusion ( $P < 0.025$  vs. baseline). At the




FIG. 2. Pancreatic clamp study: plasma concentrations of glucose, insulin, C-peptide, and glucagon during lipid infusion (●) and glycerol infusion (control; ○) in the presence of somatostatin-insulin-glucagon infusions. Data are given as means  $\pm$  SE of 6 paired experiments. \* $P < 0.05$  vs. control; § $P < 0.01$  vs. 0 min.

end of the study, plasma C-peptide concentrations were  $\sim 2.4$ -fold higher during lipid than during glycerol infusion ( $P = 0.0015$ ). Plasma glucagon concentrations were similar before and during both experiments (Fig. 1).

Rates of EGP were similar (time 0:  $9.3 \pm 0.5$  vs.  $9.0 \pm 0.8$   $\mu\text{mol} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ ) and declined to  $8.4 \pm 0.5$  and  $8.2 \pm 0.7$   $\mu\text{mol} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$  ( $P < 0.01$  and  $P < 0.005$ ) during lipid and glycerol experiments, respectively (Fig. 3). GNG contributed  $46 \pm 4$  (lipid) and  $52 \pm 3\%$  (control) to EGP at time 0 and  $55 \pm 7$  (lipid) and  $61 \pm 9\%$  (control) at 180 min. Then GNG continuously rose ( $P < 0.01$  vs. time 0) to  $66 \pm 8$  and  $73 \pm 7\%$  at 360–540 min, without a difference between lipid and glycerol experiments. **Pancreatic clamp study.** Plasma FFA concentrations increased  $\sim 5.7$ -fold ( $P < 0.01$ ) during lipid infusion and

TABLE 2  
Pancreatic clamp study: plasma concentrations of FFAs, glycerol, lactate, growth hormone, and cortisol during lipid infusion and glycerol infusion (control) in the presence of somatostatin-insulin-glucagon infusion

|                        | 0 min       | 180 min       | 360 min       | 540 min       |
|------------------------|-------------|---------------|---------------|---------------|
| FFAs (mmol/l)          |             |               |               |               |
| Lipid                  | 0.37 ± 0.09 | 1.04 ± 0.16*† | 1.64 ± 0.26‡  | 2.05 ± 0.30*§ |
| Control                | 0.50 ± 0.08 | 0.29 ± 0.08†  | 0.26 ± 0.07†  | 0.28 ± 0.07†  |
| Glycerol (mmol/l)      |             |               |               |               |
| Lipid                  | 0.14 ± 0.01 | 0.17 ± 0.01*† | 0.26 ± 0.04*† | 0.36 ± 0.04*† |
| Control                | 0.18 ± 0.02 | 0.24 ± 0.01†  | 0.35 ± 0.03§  | 0.47 ± 0.06§  |
| Lactate (mmol/l)       |             |               |               |               |
| Lipid                  | 0.75 ± 0.15 | 0.53 ± 0.09   | 0.54 ± 0.07   | 0.50 ± 0.07   |
| Control                | 0.54 ± 0.13 | 0.47 ± 0.07   | 0.55 ± 0.07   | 0.62 ± 0.10   |
| Growth hormone (ng/ml) |             |               |               |               |
| Lipid                  | 0.7 ± 0.1   | <0.5†         | <0.5†         | <0.5†         |
| Control                | 0.6 ± 0.0   | <0.5†         | <0.5†         | <0.5†         |
| Cortisol (µg/dl)       |             |               |               |               |
| Lipid                  | 9.9 ± 1.9   | 10.2 ± 1.9    | 4.9 ± 1.1†    | 4.6 ± 1.2†    |
| Control                | 9.8 ± 1.6   | 11.8 ± 1.3    | 7.9 ± 1.0     | 5.9 ± 0.8     |

Data are means ± SE of 6 paired experiments. \* $P < 0.05$ ; † $P < 0.05$ ; ‡ $P < 0.005$  vs. control; § $P < 0.005$  vs. basal.

remained constant during glycerol infusion (Table 2). Plasma glycerol concentrations rose during both protocols ( $P < 0.05$ ) and were slightly but significantly higher at 360 and 540 min of the glycerol protocol ( $P < 0.05$ ). Plasma growth hormone concentrations decreased during both experiments ( $P < 0.05$ ). Plasma concentrations of cortisol and lactate were not different between the 2 protocols.

Plasma glucose declined by ~19% ( $P < 0.05$ ) and ~29% ( $P < 0.005$ ) during the first 90 min of lipid and glycerol infusion, respectively (Fig. 2). In contrast to the basal study, plasma glucose concentrations increased thereafter by ~50% to  $7.9 \pm 0.6$  mmol/l (540 min;  $P < 0.005$  vs. baseline and vs. control) during lipid infusion, being ~1.8-fold higher than during glycerol infusion ( $4.4 \pm 0.7$  mmol/l). Plasma insulin was maintained at fasting peripheral concentrations throughout the experiments, so portal insulin concentrations were likely lower than during the basal study. Plasma C-peptide concentrations were similar at baseline and rapidly declined below the detection limit of 0.3 ng/ml ( $P < 0.0005$ ) during somatostatin exposure in both protocols. Plasma glucagon concentrations were maintained at their fasting concentrations (Fig. 2).

Absolute rates of GNG and EGP at the end of the study are presented in Fig. 3. Rates of EGP were similar to those of the basal study at time 0 (lipid  $9.6 \pm 1.3$ ; glycerol  $9.0 \pm 1.1$   $\mu\text{mol} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ ) and were unchanged at the end of the experiments (lipid  $9.9 \pm 1.2$ ; glycerol  $9.0 \pm 1.1$   $\mu\text{mol} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ ). Absolute mean enrichments at C6 ranged from 0.67 to 0.79%, and percent enrichments at C6 of glucose at -10, -5, and 0 min were essentially constant, as were those at 510, 520, and 540 min (Table 3). At time 0, GNG accounted for  $62 \pm 5$  and  $60 \pm 6\%$  of EGP during lipid and glycerol infusion, respectively. Until 180 min, GNG remained unchanged in both protocols (lipid  $59 \pm 3$ ; control  $63 \pm 4\%$ ; NS). During high-dose lipid infusion, GNG rose to  $74 \pm 3\%$  ( $P < 0.005$  vs. time 0) and was higher than during glycerol infusion (360–540 min:  $64 \pm 4\%$ ;  $P < 0.05$  vs. lipid). Using the ratio of  $^2\text{H}$  enrichments at C5 and C2, the enrichment at C2 is assumed to approximate that in body water (25,26). The enrichment at C2 was compared with that in plasma water at 540 min. The C2/water ratio was  $1.00 \pm 0.05$  in the glycerol and  $1.12 \pm 0.05$  in the lipid experiments ( $P \leq 0.05$ ).

## DISCUSSION

These studies demonstrate that short-term plasma FFA elevation induces insulin secretion in humans that is sufficient to prevent an increase in plasma glucose concentration. If plasma insulin is maintained at fasting peripheral concentrations by somatostatin-insulin infusion, FFA exposure results in a marked rise of plasma glucose concentrations and augments the contribution of gluconeogenesis to endogenous glucose production.

Plasma FFA elevation per se was likely responsible for the stimulation of insulin secretion, because 1) plasma glucose concentrations were not higher than during control experiments and 2) reduction of hepatic insulin clearance is unlikely to occur (5). The results are partly in line with a previous study reporting a small increase in plasma C-peptide without changes in plasma insulin (15). Insufficient duration or lower degree of plasma FFA elevation may explain the difference in results. Stimulation by FFAs of glucose-stimulated insulin secretion was demonstrated in humans (33,34) and rats (35), whereas reduction of plasma FFAs by acipimox




FIG. 3. EGP and the contribution of GNG after 540 min of lipid and glycerol infusion (control) without (basal study) or with (pancreatic clamp) simultaneous somatostatin-insulin-glucagon infusions. Data are given as means ± SE of 6 paired experiments. \* $P < 0.001$  vs. control-pancreatic clamp and  $P < 0.00001$  vs. lipid-basal study.

TABLE 3

Enrichments of plasma D-[6,6-<sup>2</sup>H<sub>2</sub>]glucose collected at -10 and -5 min as a percentage of enrichments at 0 min and collected at 510 and 525 min as a percentage of enrichments at 540 min

|                        | -10/0 min   | -5/0 min   | 510/540 min | 525/540 min |
|------------------------|-------------|------------|-------------|-------------|
| Basal study            |             |            |             |             |
| Control*               | 102.0 ± 2.0 | 98.1 ± 1.8 | 101.8 ± 1.7 | 104.1 ± 1.3 |
| Lipid†                 | 97.4 ± 1.9  | 99.8 ± 2.2 | 102.3 ± 1.7 | 100.0 ± 1.7 |
| Pancreatic clamp study |             |            |             |             |
| Control‡               | 99.9 ± 2.5  | 98.8 ± 1.4 | 99.2 ± 2.4  | 103.7 ± 1.5 |
| Lipid§                 | 98.6 ± 1.3  | 97.3 ± 1.2 | 96.3 ± 0.6  | 98.8 ± 1.3  |

Data are means ± SE of 6 experiments per group. Absolute enrichments: \*0.747 ± 0.061 at 0 min, 0.789 ± 0.060 at 540 min; †0.697 ± 0.042 at 0 min, 0.773 ± 0.062 at 540 min; ‡0.706 ± 0.044 at 0 min, 0.697 ± 0.048 at 540 min; §0.673 ± 0.045 at 0 min, 0.651 ± 0.040 at 540 min.

(20,23) or nicotinic acid (18) did not consistently decrease plasma insulin concentrations.

The FFA-dependent increase of insulin secretion may explain the small but significant decline of plasma glucose concentrations during the basal study. Insulin levels can be expected to be ~2.5-fold higher in the portal than the peripheral vein (36), which could attenuate EGP (37) by a decrease in GNG or, more likely, in hepatic glycogenolysis (38,39). However, EGP and GNG were not lower during lipid than during glycerol infusion, which could be partly due to gradual overreplacement of glycerol during the first 6 h of control experiments. Although glycerol contributes only ~3–10% to GNG in the postabsorptive state (40), increased availability of glycerol might have augmented percent GNG (41).

Under conditions of basal peripheral insulinemia and glucagonemia, plasma FFA elevation induced a rise of ~3.5 mmol/l in plasma glucose concentrations. This effect may result from 1 or more of the following: 1) cessation of the insulin-dependent inhibition of EGP, 2) enhanced EGP due to stimulated GNG or glycogenolysis by FFA or glycerol, or 3) reduced whole-body glucose clearance due to peripheral (skeletal muscle) insulin resistance (7,8).

The role of FFAs in the regulation of EGP might depend on the prevailing portal vein insulin concentrations. In the presence of hyperinsulinemia, lipid/heparin infusion increases EGP (5) or does not affect EGP in healthy (12,13) and diabetic (14) humans. During somatostatin-insulin clamps at postabsorptive plasma insulin concentrations, plasma FFA elevation caused a marked increase in EGP (12,15) but not under similar conditions after an overnight fast (15,16). In the present study, conditions of portal vein hypoinsulinemia and lower portal insulin/glucagon ratios might have favored glycogenolysis (38,39) and potentially GNG. Since plasma insulin and glucagon concentrations were similar in both protocols, this possibility cannot completely explain the increase of GNG during lipid infusion.

Plasma FFA elevation was shown to increase GNG in humans (16,42) by pyruvate carboxylase activation (43) and increased availability of NADH and ATP (44). Because glycerol was not infused during most previous control studies to match the lipid-induced rise in plasma glycerol, stimulation by FFAs of hepatic GNG could result exclusively from increased availability of glycerol (12,24). In the present study, EGP was not higher despite an ~12% increase in the contribution of GNG to EGP during lipid infusion. This is in agreement with a study reporting that overnight inhibition of lipolysis by acipimox decreases GNG from lactate without affecting glycemia and

EGP in type 2 diabetic subjects (20). Similarly, preventing the fall of plasma FFAs by fat-heparin infusion during an overnight fast increased Cori cycle activity but not EGP in non-diabetic humans (16). Also, the authors of that study could not exclude a gluconeogenic effect of glycerol, whose concentration was markedly increased (16). In any case, they proposed that the increase in GNG is normally matched by a decrease of glycogenolysis so that EGP will remain unchanged. Recently, evidence for such hepatic autoregulation by FFAs was supported by a study reporting that a rebound of plasma FFA concentrations of up to 1.8 mmol/l after administration of nicotinic acid augmented the contribution of GNG to EGP (18). However, effects of nicotinic acid per se on hepatic metabolism could have contributed to these results. In the present study, the increased contribution of GNG regardless of unchanged EGP can be also due to the concomitant rise in glucose concentration to ~8 mmol/l, which will suppress glucose production, or to intracellular accumulation of glucose-6-phosphate, which will inhibit glycogen phosphorylase activity and stimulate glycogen synthase in the absence of counterregulatory hormones (44).

The contribution of GNG to EGP was in good agreement with previous studies (28,29,45) and was higher only at the end of lipid infusion compared with baseline or control conditions. Percent GNG from the ratio of <sup>2</sup>H enrichments in C5/C2 of blood glucose might result in overestimation of the contribution of GNG to EGP to the extent that glucose-6-phosphate before its conversion to glucose undergoes triose phosphate cycling and underestimation of it to the extent that glycogen cycling occurs. Glycogen cycling is undetected in the fasted state (46), and triose phosphate cycling as the pentose cycle probably contributes no more than 2% to any overestimation of fractional GNG in the fasted state (30,45,47). Nevertheless, changes in their contributions induced by FFAs or glycerol cannot be ruled out. Estimates using the C5/C2 enrichment ratio assume complete isotopic equilibration of glucose-6-phosphate with fructose-6-phosphate. If so, at steady state, the enrichment in C2 equals that in body water (18,45,46). The reason in the lipid study for the C2/body water ratio of ~1.12 is unclear. However, using the C5/water rather than C5/C2 ratio would only increase the percent GNG in the lipid compared with the glycerol study.

Finally, the increase of plasma glucose concentrations could also result indirectly from a reduction in whole-body glucose clearance (5–7,48) which is due to direct inhibition of skeletal muscle glucose transport activity with subsequent reduction in rates of glucose oxidation and glycogen syn-

thesis (7,8,48). Under pancreatic clamp conditions of the present study, the inability to counteract any decline of whole-body glucose clearance by increased secretion of insulin to inhibit EGP and stimulate glucose uptake may then lead to a net increase of plasma glucose concentrations.

In conclusion, these findings are consistent with the hypothesis that FFAs are involved in the regulation of hepatic glucose metabolism (49). However, only at high plasma FFA concentrations did the contribution of gluconeogenesis to glucose production increase, whereas endogenous glucose production remained unchanged in healthy humans.

#### ACKNOWLEDGMENTS

This study was supported by grants from the Austrian Science Foundation to M.R. (P13213-MOB), from the Austrian National Bank to W.W. and M.R. (ÖNB 8196), and from the National Institutes of Health to B.R.L. (ROI-DK-14507) and G.I.S. (ROI-DK-49230 and P30-DK-45735). H.S. was supported by the Austrian Academy of Sciences (427/1997). G.I.S. is an investigator for the Howard Hughes Medical Institute.

We thank the Metabolic Unit (O.H. Lentner) and the laboratory staff of the Division of Clinical Endocrinology and Metabolism, Department of Internal Medicine III, General Hospital of Vienna, for their skillful work.

#### REFERENCES

- Felber J-P, Ferrannini E, Golay A, Mejean H, Thiebaut D, Curchod B, Maeder E, Jequier E, DeFronzo RA: Role of lipid oxidation in pathogenesis of insulin resistance of obesity and type II diabetes. *Diabetes* 36:1341-1350, 1987
- Reaven GM, Hollenbeck C, Jeng CY, Wu MS, Chen YD: Measurement of plasma glucose free fatty acid, lactate, and insulin for 24 h in patients with NIDDM. *Diabetes* 37:1020-1024, 1988
- Perseghin G, Ghosh S, Gerow K, Shulman GI: Metabolic defects in lean non-diabetic offspring of NIDDM parents: a cross-sectional study. *Diabetes* 46:1001-1009, 1997
- McGarry JD: What if Minkowski had been ageusic? An alternative angle on diabetes. *Science* 258:766-770, 1992
- Boden G: Role of fatty acids in the pathogenesis of insulin resistance and NIDDM. *Diabetes* 46:3-10, 1997
- Kelley DE, Mokan M, Simoneau JA, Mandarino LJ: Interaction between glucose and free fatty acid metabolism in human skeletal muscle. *J Clin Invest* 92:91-98, 1993
- Roden M, Price TB, Perseghin G, Falk-Petersen K, Rothman DL, Cline GW, Shulman GI: Mechanism of free fatty acid-induced insulin resistance in humans. *J Clin Invest* 97:2859-2865, 1996
- Roden M, Krssak M, Stingl H, Gruber S, Hofer A, Fürnsinn C, Moser E, Waldhäusl W: Rapid impairment of skeletal muscle glucose transport/phosphorylation by free fatty acids in humans. *Diabetes* 48:358-364, 1999
- Bogardus C, Lillioja S, Howard BV, Reaven G, Mott D: Relationships between insulin secretion, insulin action, and fasting plasma glucose concentration in nondiabetic and noninsulin-dependent diabetic subjects. *J Clin Invest* 74:1238-1246, 1984
- Groop LC, Saloranta C, Shank M, Bonadonna RC, Ferrannini E, DeFronzo RA: The role of free fatty acid metabolism in the pathogenesis of insulin resistance in obesity and noninsulin-dependent diabetes mellitus. *J Clin Endocrinol Metab* 72:96-107, 1991
- Saloranta C, Koivisto V, Widen E, Falholt K, DeFronzo RA, Harkonen M, Groop L: Contribution of muscle and liver to glucose-fatty acid cycle in humans. *Am J Physiol* 264:E599-E605, 1993
- Ferrannini E, Barrett EJ, Bevilacqua S, DeFronzo RA: Effect of fatty acids on glucose production and utilization in man. *J Clin Invest* 72:1737-1747, 1983
- Bonadonna RC, Zych K, Boni C, Ferrannini E, DeFronzo RA: Time dependence of the interaction between lipid and glucose in humans. *Am J Physiol* 257:E49-E56, 1989
- Yki-Järvinen H, Puhakainen I, Saloranta C, Groop L, Taskinen MR: Demonstration of a novel feedback mechanism between FFA oxidation from intracellular and intravascular sources. *Am J Physiol* 260:E680-E689, 1991
- Boden G, Jadhav F: Effects of lipid on basal carbohydrate metabolism in normal men. *Diabetes* 40:686-692, 1991
- Clore JN, Glickman PS, Nestler JE, Blackard WG: In vivo evidence for hepatic autoregulation during FFA-stimulated gluconeogenesis in normal humans. *Am J Physiol* 261:E425-E429, 1991
- Fulcher GR, Walker M, Catalano C, Agius L, Alberti KG: Metabolic effects of suppression of nonesterified fatty acid levels with acipimox in obese NIDDM subjects. *Diabetes* 41:1400-1408, 1992
- Chen X, Iqbal N, Boden G: The effect of free fatty acids on gluconeogenesis and glycogenolysis in normal subjects. *J Clin Invest* 103:365-372, 1999
- Vaag A, Sködt P, Gall M-A, Richter EA, Beck-Nielsen H: Effect of the antilipolytic nicotinic acid analogue acipimox on whole-body and skeletal muscle glucose metabolism in patients with non-insulin-dependent diabetes mellitus. *J Clin Invest* 88:1282-1290, 1991
- Puhakainen I, Yki-Järvinen H: Inhibition of lipolysis decreases lipid oxidation and gluconeogenesis from lactate but not fasting hyperglycemia or total hepatic glucose production in NIDDM. *Diabetes* 42:1694-1699, 1993
- Magnusson I, Rothman DL, Katz LD, Shulman RG, Shulman GI: Increased rate of gluconeogenesis in type II diabetes mellitus: a <sup>13</sup>C nuclear magnetic resonance study. *J Clin Invest* 90:1323-1327, 1992
- Gerich JE, Nurjhan N: Gluconeogenesis in type 2 diabetes. *Adv Exp Med Biol* 334:253-258, 1993
- Fery F, Plat L, Melot C, Balasse EO: Role of fat-derived substrates in the regulation of gluconeogenesis during fasting. *Am J Physiol* 270:E822-E830, 1996
- Waldhäusl W, Bratusch-Marrain P, Gasic S, Korn A, Nowotny P: Insulin production rate, hepatic insulin retention and splanchnic carbohydrate metabolism after oral glucose ingestion in hyperinsulinaemic type 2 (non-insulin-dependent) diabetes mellitus. *Diabetologia* 23:6-15, 1982
- Landau BR: Estimating gluconeogenic rates in NIDDM. *Adv Exp Med Biol* 334:205-220, 1992
- Schumann WC, Magnusson I, Chandramouli V, Kumaran K, Wahren J, Landau BR: Metabolism of [<sup>2-14</sup>C]acetate and its use in assessing hepatic Krebs cycle activity and gluconeogenesis. *J Biol Chem* 266:6985-6990, 1991
- Landau BR, Fernandez CA, Previs SF, Ekberg K, Chandramouli V, Wahren J, Kalhan SC, Brunengraber H: A limitation in the use of mass isotopomer distributions to measure gluconeogenesis in fasting humans. *Am J Physiol* 269:E18-E26, 1995
- Landau BR, Wahren J, Chandramouli V, Schumann WC, Ekberg K, Kalhan SC: Use of <sup>2</sup>H<sub>2</sub>O for estimating rates of gluconeogenesis: application to the fasted state. *J Clin Invest* 95:172-178, 1995
- Landau BR, Wahren J, Chandramouli V, Schumann WC, Ekberg K, Kalhan SC: Contributions of gluconeogenesis to glucose production in the fasted state. *J Clin Invest* 98:378-385, 1996
- Chandramouli V, Ekberg K, Schumann WC, Kalhan SC, Wahren J, Landau BR: Quantifying gluconeogenesis during fasting. *Am J Physiol* 36:E1209-E1215, 1997
- Hother-Nielsen O, Beck-Nielsen H: On the determination of basal glucose production rate in patients with type 2 (non-insulin-dependent) diabetes mellitus using primed-continuous 3-<sup>3</sup>H-glucose infusion. *Diabetologia* 33:603-610, 1990
- Waldhäusl W, Herkner K, Nowotny P, Bratusch-Marrain P: Combined 17- and 18-hydroxylase deficiency associated with complete male pseudohermaphroditism and hypoadosteronism. *J Clin Endocrinol Metab* 46:236-246, 1978
- Boden G, Chen X, Rosner J, Barton M: Effects of a 48-h fat infusion on insulin secretion and glucose utilization. *Diabetes* 44:1239-1242, 1995
- Pelkonen R, Miettinen TA, Taskinen M-R, Nikkilä EA: Effect of acute elevation of plasma glycerol, triglyceride and FFA levels on glucose utilization and plasma insulin. *Diabetes* 17:76-82, 1968
- Stein DT, Esser V, Stevenson BE, Lane KE, Whiteside JH, Daniels MB, Chen S, McGarry JD: Essentiality of circulating fatty acids for glucose-stimulated insulin secretion in the fasted rat. *J Clin Invest* 97:2728-2735, 1996
- Blackard WG, Nelson NC: Portal and peripheral vein immunoreactive insulin concentrations before and after glucose infusion. *Diabetes* 19:302-306, 1970
- Groop LC, Bonadonna RC, DelPrato S, Ratheiser K, Zych K, Ferrannini E, DeFronzo RA: Glucose and free fatty acid metabolism in non-insulin-dependent diabetes mellitus: evidence for multiple sites of insulin resistance. *J Clin Invest* 84:205-213, 1989
- Sindelar DK, Igawa K, Chu CA, Balcom JH, Neal DW, Cherrington AD: Effect of a selective rise in hepatic artery insulin on hepatic glucose production in the conscious dog. *Am J Physiol* 276:E806-E813, 1999
- Roden M, Perseghin G, Falk Petersen K, Hwang J-H, Cline GW, Gerow K, Rothman DL, Shulman GI: The roles of insulin and glucagon in the regulation of hepatic glycogen synthesis and turnover in humans. *J Clin Invest* 97:642-648, 1996
- Nurjhan N, Campbell PJ, Kennedy FP, Miles JM, Gerich JE: Insulin dose-response characteristics for suppression of glycerol release and conversion to glucose in humans. *Diabetes* 35:1326-1331, 1986
- Previs S, Cline GW, Shulman GI: A critical evaluation of mass isotopomer dis-

- tribution analysis of gluconeogenesis in vivo. *Am J Physiol* 277:E154–E160, 1999
42. Wahren J, Hagenfeldt L, Felig P: Splanchnic and leg exchange of glucose, amino acids, and free fatty acids during exercise in diabetes mellitus. *J Clin Invest* 55:1303–1314, 1975
  43. Bahl JJ, Matsuda M, DeFronzo RA, Bressler R: In vitro and in vivo suppression of gluconeogenesis by inhibition of pyruvate carboxylase. *Biochem Pharmacol* 53:67–74, 1997
  44. Youn H, Bergman RN: Enhancement of hepatic glycogen by gluconeogenic precursors: substrate flux or metabolic control? *Am J Physiol* 258:E899–E906, 1990
  45. Chandramouli V, Ekberg K, Schumann WC, Wahren J, Landau BR: Origins of the hydrogen bound to carbon 1 of glucose in fasting: significance in gluconeogenesis quantitation. *Am J Physiol* 277:E717–E723, 1999
  46. Petersen KF, Laurent D, Rothman DL, Cline GW, Shulman GI: Mechanism by which glucose and insulin inhibit net hepatic glycogenolysis in humans. *J Clin Invest* 101:1203–1209, 1998
  47. Magnusson I, Chandramouli V, Schumann WC, Kumaran K, Wahren J, Landau BR: Pentose pathway in human liver. *Proc Natl Acad Sci USA* 85:4682–4685, 1988
  48. Dresner A, Laurent D, Marcucci M, Griffin ME, Dufour S, Cline GW, Slezak LA, Andersen DK, Hundal RS, Rothman DL, Petersen KF, Shulman GI: Effects of free fatty acids on glucose transport and IRS-1-associated phosphatidylinositol 3-kinase activity. *J Clin Invest* 103:253–259, 1999
  49. Rebrin K, Steil GM, Getty L, Bergman RN: Free fatty acids as a link in the regulation of hepatic glucose output by peripheral insulin. *Diabetes* 44:1038–1045, 1995