

Differential Effects of Tumor Necrosis Factor- α on Protein Kinase C Isoforms α and δ Mediate Inhibition of Insulin Receptor Signaling

Tovit Rosenzweig,¹ Liora Braiman,¹ Asia Bak,¹ Addy Alt,¹ Toshio Kuroki,² and Sanford R. Sampson,¹

Tumor necrosis factor- α (TNF- α) is a multifunctional cytokine that interferes with insulin signaling, but the molecular mechanisms of this effect are unclear. Because certain protein kinase C (PKC) isoforms are activated by insulin, we examined the role of PKC in TNF- α inhibition of insulin signaling in primary cultures of mouse skeletal muscle. TNF- α , given 5 min before insulin, inhibited insulin-induced tyrosine phosphorylation of insulin receptor (IR), IR substrate (IRS)-1, insulin-induced association of IRS-1 with the p85 subunit of phosphatidylinositol 3-kinase (PI3-K), and insulin-induced glucose uptake. Insulin and TNF- α each caused tyrosine phosphorylation and activation of PKCs δ and α , but when TNF- α preceded insulin, the effects were less than that produced by each substance alone. Insulin induced PKC δ specifically to coprecipitate with IR, an effect blocked by TNF- α . Both PKC α and δ are constitutively associated with IRS-1. Whereas insulin decreased coprecipitation of IRS-1 with PKC α , it increased coprecipitation of IRS-1 with PKC δ . TNF- α blocked the effects of insulin on association of both PKCs with IRS-1. To further investigate the involvement of PKCs in inhibitory actions of TNF- α on insulin signaling, we overexpressed specific PKC isoforms in mature myotubes. PKC α overexpression inhibited basal and insulin-induced IR autophosphorylation, whereas PKC δ overexpression increased IR autophosphorylation and abrogated the inhibitory effect of TNF- α on IR autophosphorylation and signaling to PI3-K. Blockade of PKC α antagonized the inhibitory effects of TNF- α on both insulin-induced IR tyrosine phosphorylation and IR signaling to PI3-K. We suggest that the effects of TNF- α on IR tyrosine phosphorylation are mediated via alteration of insulin-induced activation and association of PKC δ and α with upstream signaling molecules. *Diabetes* 51:1921–1930, 2002

Tumor necrosis factor- α (TNF- α) (1) is a multifunctional cytokine, produced mainly by activated macrophages and involved in immune and proinflammatory responses. It has been suggested that TNF- α may also play a role in obesity-linked insulin resistance. Thus, adipose tissues of obese animals and human patients (1) produce high levels of TNF- α mRNA accompanied by overexpression of the cytokine. TNF- α is also expressed in human muscle and is found at higher levels in muscle from insulin-resistant and diabetic patients than in normal control subjects (2,3). Moreover, depletion of TNF- α activity, using a chimeric antibody or as occurs in transgenic animals lacking either TNF- α or TNF- α receptor, improves sensitivity to insulin in obese animals (4,5). Although the mechanism of action of TNF- α in these insulin-related effects is not fully understood, it is known that a major component involves TNF- α inhibition of upstream insulin signaling. In this regard, a number of studies have shown that TNF- α both reduces insulin-stimulated tyrosine phosphorylation of insulin receptor (IR) and IR substrate (IRS)-1 and inhibits tyrosine phosphorylation and activity of phosphatidylinositol 3-kinase (PI3-K) (6,7). Inhibition of IR signaling by TNF- α in skeletal muscle is controversial, and although some studies have confirmed this effect (8), others could not demonstrate this pathway (9,10).

Several studies have shown that the protein kinase C (PKC) family of serine threonine kinases may be implicated in various aspects of TNF- α signaling. The PKC family is composed of at least 11 isoforms, which are categorized into three groups (conventional PKCs α , β I, β II, and γ ; novel PKCs δ , ϵ , η , and θ ; and atypical PKCs ζ and λ/ι) according to their structure and mechanisms of activation (11,12). Phosphorylation on tyrosine residues is associated with activation of PKC δ , PKC β II, and PKC ζ (13–15). It is currently believed that tyrosine phosphorylation may be important in the determination of substrate specificity of the enzyme (16).

Activation and inhibition of PKC by TNF- α in a variety of cell types have been reported (17,18). With regard to the involvement of PKC isoforms in TNF- α inhibition of insulin signaling, it was shown in adipocytes that the cytokine inhibits insulin-induced translocation of PKC β and ζ to the membrane (19). In addition, overexpression of PKC ϵ increased the inhibitory effect of TNF- α on insulin signaling in HEK293 cells (20).

From the ¹Faculty of Life Sciences, Gonda-Goldschmied Center, Bar-Ilan University, Ramat-Gan, Israel; and the ²Institute of Molecular Oncology, Showa University, Hatanodai, Shinagawa-ku, Tokyo, Japan.

Address correspondence and reprint requests to S. R. Sampson, Department of Life Sciences, Bar-Ilan University, Ramat-Gan 52900, Israel. E-mail: sampsos@mail.biu.ac.il.

Received for publication 17 May 2001 and accepted in revised form 11 March 2002.

2DG, 2-deoxy-D-glucose; IR, insulin receptor; IRS, IR substrate; PI3-K, phosphatidylinositol 3-kinase; PKC, protein kinase C; TNF- α , tumor necrosis factor- α .

Recent studies implicate certain PKC isoenzymes in the insulin-signaling cascade. Insulin activates PKCs α , β II, and ζ in several cell types, including cell lines of skeletal muscle (21–24). We recently reported that insulin stimulates glucose uptake and induces tyrosine phosphorylation, translocation, and activation of PKCs β II, δ , and ζ in primary cultures of rat skeletal myotubes (15). Activation of PKCs β II and ζ apparently occurs via a PI3-K-dependent pathway, whereas PKC δ appears to be activated upstream (15,25), perhaps by the IR itself or by the IRS proteins. The various PKC isoforms are both activated by insulin and can also modulate insulin signaling by regulating the kinase activity of IR. Indeed, PKC δ was found to upregulate IR tyrosine phosphorylation in rat skeletal muscle cells (26), whereas PKCs α , δ , and θ were found to inhibit this phosphorylation in HEK293 cells (27). Because various PKC isoforms have been found to interact with both IR and IRS, it is still not clear whether specific PKC isoforms regulate IR directly or through IRS-1.

TNF- α induces serine phosphorylation of IRS-1 and reduces its ability to interact with the juxtamembrane region of IR (28). Whereas IRS-1 from cells treated with TNF- α inhibits insulin-induced IR tyrosine phosphorylation, it is not certain whether this effect occurs directly by interaction of IRS-1 with IR or indirectly through another element in the pathway (29,30). PKC isoforms, which are activated by insulin, are possible candidates for phosphorylating the receptor on serine residues and affecting subsequent signaling.

The preparation of primary skeletal muscle cultures obtained from neonatal mice pups is a useful model for the study of regulation of insulin signaling by TNF- α . These cells, plated initially as individual myoblasts, align and fuse into multinucleated muscle fibers by day 3–4 in vitro. The mature fibers display resting membrane and action potentials that are nearly identical to those seen in vivo. In addition, the physiological expression of a number of membrane proteins in this preparation, in contrast to muscle cell lines such as L6, closely resemble that obtained in vivo (31–33). In this study, we investigated the possibility that TNF- α may differentially affect the interaction of specific PKC isoforms with upstream elements in the IR signaling cascade to modulate IR signaling. The results show that insulin induces PKC δ to associate with IR and causes PKC α , which is constitutively associated with IRS-1, to disassociate from this docking protein. TNF- α interferes with these insulin-induced effects, resulting in a decrease in insulin-induced tyrosine phosphorylation of IR. Overexpression of PKC δ on one hand and inhibition of PKC α on the other reverse the influence of TNF- α on IR autophosphorylation. These results strongly suggest that TNF- α alters IR signaling in skeletal muscle through its effects on PKC α and δ .

RESEARCH DESIGN AND METHODS

Materials. Tissue culture media and serum were purchased from Biological Industries (Beit HaEmek, Israel). TNF- α was purchased from Sigma (St. Louis, MO). Antibodies to various proteins were obtained from the following sources: monoclonal antibodies to IR β were purchased from Transduction Laboratories (Lexington, KY). Monoclonal antiphosphotyrosine was obtained from Upstate Biotechnology (Lake Placid, NY). Antibodies to IRS-1, IRS-2, and specific PKC isoforms were purchased from Santa Cruz Biotechnology (Santa Cruz, CA; polyclonal) and from Transduction Laboratories (monoclonal). Horseradish peroxidase and anti-rabbit and anti-mouse IgG were obtained

from BioRad (Israel). Leupeptin, aprotinin, PMSF (phenylmethylsulfonyl fluoride), orthovanadate, and pepstatin were purchased from Sigmas. G06976 was purchased from Calbiochem (San Diego, CA).

Methods

Preparation of mouse muscle cell cultures. Skeletal muscle cultures were freshly prepared for each experiment from the limb muscles obtained from 1- to 2-day neonatal mice as described (15,25,34). Fusion of plated myoblasts into mature myotubes occurs 3–4 days after plating. On day 5 in culture, myotubes were transferred to low-glucose (4.5 mmol/l) serum-free DMEM containing 1% BSA for 24 h before study.

Immunoprecipitation. Culture dishes (90 mm; Nunc) containing the muscle cultures were washed with Ca²⁺/Mg²⁺-free PBS and then mechanically detached in RIPA (radioactive immunoprecipitation assay) buffer (50 mmol/l Tris HCl, 150 mmol/l NaCl, 1 mmol/l EDTA, 10 mmol/l NaF, 1% Triton X-100, 0.1% SDS, and 1% Na deoxycholate; pH 7.4) containing a cocktail of antiproteases (20 μ g/ml leupeptin, 10 μ g/ml aprotinin, 0.1 mmol/l PMSF, and 1 mmol/l dithiothreitol) and antiphosphatases (200 mmol/l orthovanadate and 2 μ g/ml pepstatin). After scraping, the preparation was centrifuged at 20,000g for 20 min at 4°C. The supernatant was used for immunoprecipitation as described (15,25).

Western blot analysis. Crude and fractionated lysates of control and treated cultures were subjected to SDS-PAGE and electrophoretic transfer to Immobilon-P (Millipore) membranes. The membranes were subjected to standard blocking procedures and incubated with monoclonal and polyclonal antibodies (25).

Glucose uptake. Glucose transport was evaluated by measuring 2-deoxy-D-glucose (2DG) uptake as described (15,25). After appropriate treatment, cells were washed three times with 0.5 ml PBS, and the final wash was replaced immediately with 0.5 ml PBS containing 2 mmol/l glucose and tracer amounts (1 μ Ci/ml) of [³H]-2DG. Cells were then incubated for 15 min at 37°C, washed four times with 0.5 ml cold PBS, and then detached from the wells by addition of 300 μ l Triton X-100 (1%) and incubation for 30 min. The contents of each well were transferred to counting vials, and 3.5 ml scintillation fluid was added to each vial. Samples were counted in the [³H] window of a Tricarb scintillation counter. Nonspecific uptake was determined in the presence of excess (100 μ mol/l) D-glucose. Experiments were carried in triplicate. Net specific uptake was then calculated as the difference between the total and nonspecific values.

PKC recombinant adenoviruses and viral infection of cultures. The recombinant adenovirus vectors were constructed as described (35). The dominant-negative mutant of mouse PKC δ was generated by substitution of the lysine residue at the ATP binding site with alanine (36). The mutant δ cDNA was cut from the SRD expression vector with *Eco*RI and ligated into the pAxCALw cosmid cassette to construct the Ax vector. Dominant-negative activity was demonstrated by abrogation of autophosphorylation (36).

After differentiation of cultured rat myoblasts into myotubes, the culture medium was aspirated and cultures were infected with medium containing PKC recombinant adenoviruses as described (25).

PKC activity. Specific PKC activity was determined in freshly prepared immunoprecipitates from mature muscle cultures after appropriate treatments as described (15,25). These lysates were prepared in RIPA buffer without NaF. Activity was measured using the SignaTECT PKC Assay System (Promega, Madison, WI). The kit contains phosphatidylserine and diacylglycerol and uses Neurogranin as substrate.

RESULTS

Characterization of TNF α effects on upstream IR signaling in cultured skeletal muscle. TNF- α has been shown to inhibit insulin-induced phosphorylation of the receptor tyrosine kinase in several cell types (6,30,37). Figure 1 shows the effects of TNF- α on insulin-induced tyrosine phosphorylation in cultured mouse skeletal muscle. Under control conditions, immunoprecipitated IR displayed an increase in tyrosine phosphorylation within 1 min of insulin stimulation. This remained elevated for at least 5 min. IR phosphorylation then decreased toward prestimulation levels by 15 min. As the maximum level of IR tyrosine phosphorylation occurred within 1–5 min of insulin stimulation, this time period of insulin stimulation was used in all subsequent experiments. Effects of TNF- α on insulin-induced tyrosine phosphorylation of IR are shown in Fig. 1B. Pretreatment of cells with TNF- α (1




FIG. 1. TNF- α inhibits insulin-induced tyrosine phosphorylation of IR in cultured mouse skeletal myotubes. Studies were performed on 6-day-old cultured myotubes, which were transferred to serum-free low-glucose Eagle's medium, 24 h before experiments were conducted. **A:** Time course of insulin-induced tyrosine phosphorylation of IR in control cells. Protein extracts from untreated cultures or insulin-stimulated cultures treated for various time periods (1, 5, 15, or 30 min) were immunoprecipitated with specific antiphosphotyrosine antibodies. Immunoprecipitates were run on SDS-PAGE, transferred to Immobilon-P membranes, and immunoblotted with specific anti-IR antibodies. **B:** Effects of pretreatment with TNF- α on insulin-induced tyrosine phosphorylation of IR. Protein extracts were prepared from untreated cultures or cultures stimulated with insulin for 5 min with and without prior treatment with TNF- α (1 nmol/l) for 5, 15, or 30 min. Extracts were immunoprecipitated with specific antiphosphotyrosine antibodies. Immunoprecipitates were run on SDS-PAGE, transferred to Immobilon-P membranes, and immunoblotted with specific anti-IR antibodies. The Western blots shown are representative of three blots from three separate cultures. **C:** TNF- α inhibits insulin-induced tyrosine phosphorylation of IRS-1 in cultured mouse skeletal myotubes. Cells were either untreated (C) or treated with insulin for 5 min (I), with 1 nmol/l TNF- α for 5 min (T), or with TNF- α for 5 min before addition of insulin for 5 min (T+I). Extracts were immunoprecipitated with specific antiphosphotyrosine antibodies. Immunoprecipitates were run on SDS-PAGE, transferred to Immobilon-P membranes, and immunoblotted with specific anti-IRS-1 antibody. The Western blots shown are representative of three blots from three separate cultures. **D:** TNF- α inhibits insulin-induced glucose uptake. Myotubes were transferred to serum-free low-glucose medium for 24 h. Cells were either untreated or treated with insulin for 20 min, with 1 nmol/l TNF- α for 5 or 30 min, or with TNF- α before addition of insulin. The results are described as percent of control. Glucose transport was evaluated by measuring [3 H]-2DG for 15 min as described in RESEARCH DESIGN AND METHODS. Data are presented as percent of basal (18 ± 1.3 pmol \cdot mg protein $^{-1}$ \cdot min $^{-1}$), which was set at 100%. Each bar represents the means \pm SE of triplicate values in each of three experiments performed on separate cultures.

nmol/l) for 5 min completely inhibited insulin-induced phosphorylation of IR. Longer pretreatment (15–30 min) with TNF α not only blocked the insulin-induced phosphorylation of IR but also reduced the level of phosphorylation to below pretreatment levels.

The first element downstream of IR is the docking family of IRS proteins. In cultured mouse skeletal myotubes, the primary IRS protein involved in insulin signaling is IRS-1, which is also rapidly phosphorylated on tyrosine. As shown in Fig. 1C, immunoprecipitated IRS-1 is tyrosine phosphorylated within 5 min after insulin stimulation. Pretreatment of cells with TNF- α (1 nmol/l for 5 min) markedly reduced phosphorylation of IRS-1 by insulin. In addition, TNF- α inhibited insulin-induced association of IRS-1 with the p85 subunit of PI3-K (Fig. 9, left panel). Effects of TNF- α on the uptake of 2-deoxy-D-glucose (2-DG) were also studied (Fig. 1D). Treatment of myotubes with TNF- α for 5 min did not alter basal 2-DG uptake but significantly reduced the stimulatory effect of insulin. Treatment with TNF- α for 30 min caused a slight but significant increase in basal 2-DG uptake, which was not further increased by insulin. Thus, these findings in cul-

tured mouse skeletal myotubes confirm that TNF- α inhibits early upstream insulin signaling as well as glucose transport.

TNF- α affects insulin-induced association between specific PKC isoforms and upstream elements. Several studies indicate that TNF- α inhibition of insulin signaling is mediated by other, as yet unknown, elements. Among the possible candidates is the PKC family of serine-threonine kinases, because TNF- α is known to affect certain members of this family, and some PKC isoforms can modulate IR autophosphorylation. In preliminary studies, we found that cultured mouse skeletal muscle cells express PKC isoforms α , β 2, δ , ϵ , θ , and ζ , in agreement with other studies on mammalian skeletal muscle in vivo and in culture (15,38–40). In a recent study on rat skeletal muscle in primary culture, we showed that insulin induces a rapid and specific physical association between IR and PKC δ and that this physical linkage is essential for the continuation of the IR signaling cascade (26). We therefore reasoned that interference of PKC interactions with upstream elements might be involved in TNF- α effects on IR signaling. Accordingly, in this series of experiments, we




FIG. 2. TNF- α inhibits insulin-induced coimmunoprecipitation of PKC δ with IR (A and B) and IRS-1 (C and D). Cells were either untreated (C) or treated with insulin for 5 min (I), with 1 nmol/l TNF- α for 5 min (T), or with TNF- α for 5 min before addition of insulin for 5 min (T+I). Cell lysates were immunoprecipitated with anti-IR or anti-IRS-1 antibodies and then subjected to SDS-PAGE and immunoblotted with specific anti-PKC δ antibodies. A and C: Western blots of PKC δ induced to coimmunoprecipitate with IR and IRS-1, respectively. B and D: Optical density measurements of the Western blots. The values represent the means \pm SE of three measurements in three separate cultures.

sought to determine whether TNF- α affects any physical interactions induced by insulin between various PKC isoforms and IR, IRS-1, and IRS-2. In these studies, we immunoprecipitated IR, IRS-1, and IRS-2 from control cells, insulin-stimulated cells, and insulin-stimulated cells after pretreatment with TNF- α . After SDS-PAGE and transfer, the immunoprecipitated proteins were probed with specific anti-PKC antibodies. Of the PKC isoforms examined (PKCs α , β II, δ , ϵ , and ζ), only PKCs α and δ were found to coimmunoprecipitate with upstream elements. PKC δ was constitutively associated with both IR and IRS-1, and this association was increased within 5 min by insulin stimulation. PKC α was found to be constitutively associated with IRS-1 only, and this association was decreased within 5 min by insulin stimulation (Figs. 3 and 4). None of the other PKC isoforms examined coassociated with IR, IRS-1, or IRS-2. Moreover, IRS-2 did not coimmunoprecipitate with either PKC δ or α under basal or insulin-stimulated conditions (not shown). Therefore, all subsequent studies were done on IR and IRS-1 interactions with PKCs δ and α .

Treatment of muscle cells with TNF- α for 5 min before insulin stimulation had opposite effects on PKC association with IR and IRS-1. Thus, on one hand, as shown in Fig. 2, TNF- α increased the association between IR and IRS-1 with PKC δ and reduced the effect of insulin to increase coassociation of these elements with PKC δ . On the other hand, as illustrated in Fig. 3, this cytokine, while also inducing an increase in PKC α -IRS-1 association, prevented the ability of insulin to cause IRS-1 to dissociate from PKC α .

TNF- α activates specific PKC isoforms. Previous results from our laboratory have shown that insulin stimulation of cultured rat skeletal muscle cells results in activation of specific PKC isoforms (15). We therefore considered the possibility that the inhibitory effect of TNF- α on IR tyrosine phosphorylation might involve modulation of the activity of PKCs δ and α . Accordingly, we examined the effects of insulin and TNF- α on the activity of these PKC isoforms. Figure 4 shows the effects of insulin and TNF- α , separately and in combination, on the activity of specific PKC isoforms. In these studies, PKC isoforms were immunoprecipitated with specific antibodies from control cells and from cells treated with insulin, or with insulin after TNF- α , and the PKC activity of the immunoprecipitated protein was measured by an activity assay, as described in RESEARCH DESIGN AND METHODS. As can be seen, both insulin and TNF- α increased the activity of each of the PKC isoforms, the effect of the latter being at least as effective as that of the former. However, the activity of each of the enzymes from cells treated with TNF- α before insulin was less than that in cells treated with each substance alone.

Activation of PKC isoforms in skeletal muscle by insulin is associated with an increase in the tyrosine phosphorylation state (15). Accordingly, we examined effects of insulin and TNF- α on tyrosine phosphorylation of PKC α and δ . In one series of experiments, we immunoprecipitated specific PKC isoforms and performed Western blotting with antiphosphotyrosine antibodies. The results are exemplified in Fig. 5, which shows that within 5 min after insulin stimulation, there was a strong increase in tyrosine




FIG. 3. TNF- α prevents insulin-induced decrease in PKC α /IRS-1 association. Cells were either untreated (C) or treated with insulin for 5 min (I), with 1 nmol/l TNF- α for 5 min (T), or with TNF- α for 5 min before addition of insulin for 5 min (T+I). Cell lysates were immunoprecipitated with anti-IR or anti-IRS-1 antibody, subjected to SDS-PAGE, and then immunoblotted with anti-PKC α antibodies. **A:** lack of either constitutive or insulin-induced association between PKC α and IR. **B:** Effects of TNF- α on insulin-induced dissociation between PKC α and IRS-1. The Western blots shown are representative of results obtained in three separate experiments on different cultures. **C:** Graphs of optical density measurements of the Western blots. The values represent the means \pm SE of three measurements in three separate cultures.

phosphorylation of PKCs α and δ . Similar to insulin, TNF- α also induced phosphorylation of the α and δ isoforms within 5 min. As both insulin and TNF- α induced tyrosine phosphorylation of PKC isoforms α and δ , we expected that addition of TNF- α followed by insulin might produce an additive effect on these isoforms. However, when myotubes were treated with TNF- α before insulin, the level of tyrosine phosphorylation of each of the PKC isoforms induced by insulin stimulation was lower than that induced by insulin alone. In another series of studies, we immunoprecipitated with antiphosphotyrosine anti-




FIG. 4. TNF- α reduces insulin-induced activity of specific PKC isoforms in cultured mouse skeletal myotubes. Cells were either untreated (C) or treated with insulin for 5 min (I), with 1 nmol/l TNF- α for 5 min (T), or with TNF- α for 5 min before addition of insulin for 5 min (T+I). Cell lysates were immunoprecipitated with specific antibodies against PKC α (A) or PKC δ (B). Activity was determined with a SignaTECT PKC Assay System. The values represent the means \pm SE of three measurements in three separate cultures.

bodies and immunoblotted with specific anti-PKC antibodies to each of the isoforms. The results were essentially the same as those obtained with immunoprecipitation of the PKC isoforms.

Overexpression of specific PKC isoforms modifies effects of TNF- α on IR signaling. The upstream locations of PKC isoforms α and δ in the insulin signaling cascade suggest that these isoforms may play a direct role in the regulation of IR function. Indeed, our results so far have shown that TNF- α not only affects the interaction of the α and δ isoforms with IR and IRS-1 but also modifies the effects of insulin on these interactions. Thus, it is possible that TNF- α modulation of the activity of these PKC isoforms may affect IR autophosphorylation. We therefore examined more directly whether the ability of TNF- α to reduce insulin-induced activity of PKCs α and δ might play a role in the effects of this cytokine on IR autophosphorylation. To this end, we overexpressed these isoforms using an adenovirus expression system. When




FIG. 5. TNF- α reduces insulin-induced tyrosine phosphorylation of specific PKC isoforms in cultured mouse skeletal myotubes. Cells were either untreated (C) or treated with insulin for 5 min (I), with 1 nmol/l TNF- α for 5 min (T), or with TNF- α for 5 min before addition of insulin for 5 min (T+I). Extracts were immunoprecipitated with either specific antiphosphotyrosine or anti-PKC antibodies. Immunoprecipitates were run on SDS-PAGE and transferred to Immobilon-P membranes. Anti-PKC immunoprecipitates were immunoblotted with antiphosphotyrosine antibodies, and antiphosphotyrosine immunoprecipitates were immunoblotted with specific antibodies against PKC- α (A) or PKC δ (B). The Western blots shown are representative of three blots from three separate cultures. The lower row in each blot shows the protein level of each individual PKC isoform.

infected with recombinant adenovirus constructs containing cDNA for wild-type PKC δ , kinase-inactive dominant-negative PKC δ , or wild-type PKC α , myotubes displayed elevated protein expression of the transfected isoform compared with the expression of the endogenous isoforms (Fig. 6). Overexpression of each isoform, with the exception of dominant-negative PKC δ , also resulted in elevated kinase activity of the specific PKC isoform transfected (data not shown).

Next, we studied the effect of this specific overexpression of PKC α or - δ on insulin-induced IR tyrosine phosphorylation. As shown in Fig. 7A, overexpression of PKC δ resulted in an increase in tyrosine phosphorylation of IR in control and TNF- α treated cells (lanes 1, 2, 7, and 8). This effect was not additive to that of insulin, since insulin stimulation in cells overexpressing PKC δ failed to further increase the phosphorylation state (lanes 2-4). Furthermore, overexpression of PKC δ prevented the inhibitory effect of TNF- α on insulin-induced IR tyrosine phosphorylation (lanes 5 and 6). To confirm that PKC δ overexpression increases IR tyrosine phosphorylation, we infected cells with kinase-inactive dominant-negative PKC δ . The infected cells displayed high protein levels of the inactive PKC δ (Fig. 6). Overexpression of dominant-negative PKC δ abrogated both basal and insulin-induced IR tyrosine phosphorylation (Fig. 7B, lanes 1-4). In addition, the




FIG. 6. Overexpression of specific PKC isoforms. Five-day-old myotube cultures were infected with adenovirus constructs of specific PKC isoenzymes for 12 h. Whole-cell lysates of nonoverexpressing and specific PKC isoform-overexpressing cells were subjected to SDS-PAGE, transferred to Immobilon-P membranes, and immunoblotted with specific anti-PKC antibodies. Western blots of wild-type PKC δ (A; O.E. δ), dominant-negative PKC δ (B; D.N. δ), and wild-type PKC α (C; O.E. α) in noninfected (C) and adenovirus infected cells (right-side blots). The blots show strong overexpression of the PKC isoforms in adenovirus-infected compared with wild-type noninfected cells.




FIG. 7. Effects of overexpression of WTPKC δ and dominant-negative PKC δ on insulin-induced tyrosine phosphorylation of IR. Control noninfected and infected cells overexpressing wild-type PKC δ (**A**) and dominant-negative PKC δ (**B**) were treated for 5 min separately with insulin or TNF- α or 5 min with TNF- α followed by 5 min with insulin. Cell lysates were immunoprecipitated with antiphosphotyrosine antibodies. Immunoprecipitates were run on SDS-PAGE, transferred to Immobilon-P membranes, and immunoblotted with specific anti-IR antibodies. The Western blot shown is representative of results obtained in four separate experiments on four separate cultures.

effect of TNF- α on both basal and insulin-induced tyrosine phosphorylation in cells overexpressing dominant-negative PKC δ was not significantly greater than either TNF- α or dominant-negative PKC δ alone (Fig. 7B, lanes 5–8).

In contrast to the effects of PKC δ overexpression, overexpression of PKC α inhibited both basal and insulin-induced IR tyrosine phosphorylation, as shown in Fig. 8A (lanes 1–4). Interestingly, insulin did appear to increase tyrosine phosphorylation of IR in PKC α -overexpressing cells that had been treated initially with TNF- α (lanes 6 and 8), but the level was considerably less than that in control noninfected myotubes (lane 1). To determine whether PKC α is necessary for the inhibitory function of TNF- α on IR autophosphorylation, we used a selective inhibitor of the α and β PKC isoforms, GÖ6976 (Fig. 8B).




FIG. 8. Effects of overexpression and blockade of PKC α on TNF- α inhibition of insulin-induced tyrosine phosphorylation of IR. Cell lysates were prepared from cells treated for 5 min with insulin or TNF- α or 5 min with TNF- α followed by 5 min with insulin. The lysates were immunoprecipitated with antiphosphotyrosine antibodies, and immunoprecipitates were run on SDS-PAGE, transferred to Immobilon-P membranes, and immunoblotted with specific anti-IR antibodies. **A:** Overexpression of PKC α inhibits insulin-induced autophosphorylation of IR. Lysates were prepared from control noninfected and PKC α -overexpressing cells. **B:** Blockade of PKC α prevents TNF- α inhibition of IR autophosphorylation. Lysates were prepared from control cells and cells pretreated for 20 min with GÖ6976 (10 nmol/l). The Western blots are representative of results obtained in three separate experiments on three separate cultures.

(Because PKC β I is not detected in primary cultures of mouse skeletal muscle, we used this inhibitor as a selective inhibitor against PKC α in this system.) In contrast to effects of PKC α overexpression, inhibition of this isoform increased basal tyrosine phosphorylation of IR (lanes 1 and 2). In addition, inhibition of PKC α abrogated the ability of TNF- α to inhibit IR autophosphorylation (lanes 5 and 6). The results indicate that one component of the effects of TNF- α on IR signaling involves the activity of




FIG. 9. Effects of overexpression of PKC δ and blockade of PKC α on insulin-induced association between IRS-1 and PI3-K. Control noninfected cells (left panel), infected cells overexpressing WTPKC δ (middle panel), and cells treated with GÖ6976 (right panel) were treated for 5 min separately with insulin or TNF- α or 5 min with TNF- α followed by 5 min with insulin. Cell lysates were immunoprecipitated with anti-IRS-1 antibodies. Immunoprecipitates were run on SDS-PAGE, transferred to Immobilon-P membranes, and immunoblotted with specific anti-p85 PI3-K antibodies. The Western blot shown is representative of results obtained in three separate experiments on three separate cultures.

PKC α and its association with IRS-1. In parallel, TNF- α reduced responses to insulin by inhibiting insulin-induced activity of PKC δ and its association with IR and IRS-1.

The results therefore show that overexpression of PKC δ and inhibition of PKC α were able to reverse the deleterious effects of TNF- α on IR phosphorylation. To determine whether these treatments of the PKCs δ and α might also influence progression of the IR signal, we examined the association between IRS-1 and PI3-K which is one of the downstream steps in the IR signaling pathway. As shown in Fig. 9 (left panel), insulin induced PI3-K to associate with IRS-1, and TNF- α reduced this effect. Overexpression of PKC δ increased basal association between these two proteins, even in the absence of insulin, and insulin did not further increase this effect. Moreover, the inhibitory effect of TNF- α on IRS-1/PI3-K association was completely abrogated (Fig. 9, middle panel). Effects of inhibition of PKC α by GÖ6976 were essentially the same as those obtained with overexpression of PKC δ (Fig. 9, right panel). Thus, either PKC δ overexpression or PKC α blockade effectively antagonizes the inhibitory effects of TNF- α on signaling from IR to PI3-K.

DISCUSSION

In this study, we have shown that TNF- α has two distinct mechanisms to oppose insulin action on upstream elements in the IR signaling pathway. In each case, the effect is related to interaction with specific PKC isoforms. These interactions are summarized in Fig. 10. Thus, on one hand, insulin induced a physical interaction of PKC δ with IR and IRS-1 (Fig. 10A), and TNF- α prevented this interaction (Fig. 10B). On the other hand, insulin disrupted the constitutive association that occurs in this preparation between IRS-1 and PKC α (Fig. 10A), and TNF- α caused this association to increase (Fig. 10B). The effects of

insulin and TNF- α on both PKC α and - δ were accompanied by tyrosine phosphorylation and activation. Although it has been shown that stimulation of PKCs α and δ can inhibit IR autophosphorylation, and although various effects of TNF- α on PKC have been reported (17), a link between these effects and TNF- α inhibition of IR tyrosine phosphorylation has not been established.

This is the first report to demonstrate that PKC α may actually be constitutively associated with IRS-1 and that insulin causes these elements to physically dissociate. Others have shown, however, that activation of PKC α not only inhibits IR signaling (41,42) but also that PKC α may require IRS-1 for inhibition of IR tyrosine kinase activity (27). Activation of PKC α and its increased physical association with IRS-1 in response to TNF- α , as reported here, are thus consistent with the involvement of IRS-1 in PKC α inhibition of IR signaling. Moreover, the ability of insulin




FIG. 10. Schematic diagram of insulin (A) and TNF- α (B) effects on PKC interactions in insulin signaling. A. Insulin induces tyrosine phosphorylation of PKCs α and δ , as well as association of PKC δ with IR and IRS-1, and causes PKC α to dissociate from IRS-1. B. Pretreatment with TNF- α reduces insulin-induced tyrosine phosphorylation of PKCs α and δ . In addition, TNF- α interferes with insulin-induced interactions between the PKC isoforms and IR and IRS-1; PKC δ association with IR and IRS-1 is inhibited, whereas association between PKC α and IRS-1 is strengthened.

to cause dissociation between IRS-1 and PKC α may, by reducing serine phosphorylation, allow the insulin signal to continue. The results demonstrating that TNF- α activation of PKC α is involved in inhibition of IR signaling are also consistent with an earlier study conducted on a line of CHO cells overexpressing PKC α . Activation of PKC α in these cells by the phorbol ester, TPA (tetradecanoylphorbol acetate), also inhibited tyrosine phosphorylation of IRS-1 in response to insulin stimulation (42). The results obtained in studies on overexpression and blockade of PKC α indicate that this isoform is essential for the inhibitory effect of TNF α on insulin signaling. Thus, overexpression of PKC α alone was sufficient to block both basal and insulin-induced IR tyrosine phosphorylation, and treatment with a PKC α inhibitor elevated basal IR autophosphorylation and blocked the inhibitory effect of TNF α . Moreover, the fact that PKC α associates with IRS-1 and not with IR further indicates that IRS-1 is required for the inhibitory effect of PKC α and that it plays a role in reduction of insulin signaling, as previously suggested (27,29).

As we have shown, PKC δ is induced by insulin to associate with both the IR and its first downstream element, IRS-1. This association, upstream in insulin signal transduction, is consistent with the suggested role for PKC δ in regulating the function of IR itself (25,26). Our results, showing that an important component of the ability of TNF- α to interfere with IR signaling involves disturbance of the insulin-induced physical association between PKC δ and IR/IRS-1, further strengthen this idea. Additional support for an upstream location of PKC δ in insulin signaling is provided by results obtained in rat skeletal muscle showing that specific inhibitors of PI3K fail to inhibit insulin-induced activation of PKC δ (15).

We found that overexpression of PKC α reduced both basal and insulin-induced tyrosine phosphorylation of IR, whereas inhibition of PKC α increased IR tyrosine phosphorylation in the basal state. These results are in agreement with a previous study (27) in which it was shown that PKC isoforms α , δ , and θ , when coexpressed with HIR (human insulin receptor) and IRS-1 in HEK293 cells, inhibited HIR tyrosine phosphorylation, probably through serine/threonine phosphorylation of IRS-1. PKC δ , on the other hand, appears to have a dual effect on IR phosphorylation. In addition to serine phosphorylation, which was observed in cells overexpressing wild-type PKC δ even without insulin stimulation (data not shown; see also 26), overexpression of PKC δ resulted in IR tyrosine phosphorylation. This was not reported by Kellerer et al. (27). Because PKCs are serine/threonine kinases, the effect of PKC δ to increase IR tyrosine phosphorylation must be mediated by some other factor or factors. One possibility is that PKC δ , by its physical interaction with IR, may have a permissive effect on the autophosphorylation mechanism. Blockade of IR autophosphorylation by overexpression of DMPKC δ is consistent with this possibility. Alternatively, or additionally, PKC δ may affect other elements that mediate this phosphorylation. Another possibility may be related to the reported effects of PKC δ on IR routing (26,43). Recent results from our laboratory suggest that PKC δ increases the routing of IR by modulating the activity of specific proteins involved in the internalization.

Internalized IR is relatively highly phosphorylated on tyrosine residues (44).

The results regarding the opposing effects of both insulin and TNF- α on PKCs α and δ suggest that the two isoforms may play opposite roles in insulin signaling. Opposing effects of these isoforms have been reported in other systems and in signaling cascades other than the insulin signaling pathway, such as induction or inhibition of apoptosis, regulation of proliferation and differentiation, and mediation of cell transformation (45–47). Our additional finding that blockade of PKC α abrogated TNF- α inhibition of IR signaling further demonstrates that both PKC isoforms are required for its effect.

The results of this study indicate that TNF- α inhibition of insulin-induced IR and IRS-1 phosphorylation, IRS-1/PI3K association, and glucose transport occurs, in part, via modulation of the activity of PKC isoforms α and δ and their association with upstream elements in the insulin signaling cascade. We, therefore, propose that TNF- α activates PKC α and δ and influences their association with IR and IRS-1 in a manner that interferes with the ability of insulin to regulate these isoforms. We further suggest that tyrosine phosphorylation, which may occur at distinct and separate sites in response to insulin than to TNF- α , is a key element in this phenomenon. This would be consistent with studies that indicate the importance of tyrosine phosphorylation in directing the substrate specificity of PKC, in particular PKC δ (16). The mechanisms and specific sites of tyrosine phosphorylation of PKCs α and δ in response to insulin and TNF- α are currently being investigated. The apparently mutually antagonistic effects of insulin and TNF- α on major elements upstream in the insulin signaling cascade indicate that there may be a delicate balance among different substrates for the satisfactory propagation of the insulin signal. Moreover, the effects of this cytokine on specific PKC isoform interactions with these upstream elements support a role for TNF- α in insulin resistance of skeletal muscle.

ACKNOWLEDGMENTS

This study was supported in part by the Sorrell Foundation, the Ben and Effie Raber Research Fund, the Harvett-Aviv Neuroscience Research Fund, and grants from the Israel Science Foundation (founded by the Israel Academy of Sciences and Humanities) and the Chief Scientists Office of the Israel Ministry of Health. S.R.S. is the incumbent of the Louis Fisher Chair in Cellular Pathology.

This work represents an essential portion of a thesis submitted by T.R. in partial fulfillment of the PhD degree in the Faculty of Life Sciences at Bar-Ilan University.

REFERENCES

- Hotamisligil GS, Spiegelman BM: Tumor necrosis factor α : a key component of the obesity-diabetes link. *Diabetes* 43:1271–1278, 1994
- Hotamisligil GS, Arner P, Caro JF, Atkinson RL, Spiegelman BM: Increased adipose tissue expression of tumor necrosis factor- α in human obesity and insulin resistance. *J Clin Invest* 95:2409–2415, 1995
- Kern PA, Saghizadeh M, Ong JM, Bosch RJ, Deem R, Simsolo RB: The expression of tumor necrosis factor in human adipose tissue: regulation by obesity, weight loss, and relationship to lipoprotein lipase. *J Clin Invest* 95:2111–2119, 1995
- Uysal KT, Scheja L, Wiesbrock SM, Bonner-Weir S, Hotamisligil GS: Improved glucose and lipid metabolism in genetically obese mice lacking aP2. *Endocrinology* 141:3388–3396, 2000

5. Uysal KT, Wiesbrock SM, Marino MW, Hotamisligil GS: Protection from obesity-induced insulin resistance in mice lacking TNF- α function. *Nature* 389:610–614, 1997
6. Kroder G, Bossenmaier B, Kellerer M, Capp E, Stoyanov B, Muhlhofer A, Berti L, Horikoshi H, Ullrich A, Haring H: Tumor necrosis factor- α - and hyperglycemia-induced insulin resistance: evidence for different mechanisms and different effects on insulin signaling. *J Clin Invest* 97:1471–1477, 1996
7. Kanety H, Feinstein R, Papa MZ, Hemi R, Karasik A: Tumor necrosis factor α -induced phosphorylation of insulin receptor substrate-1 (IRS-1): possible mechanism for suppression of insulin-stimulated tyrosine phosphorylation of IRS-1. *J Biol Chem* 270:23780–23784, 1995
8. Del Aguila LF, Claffey KP, Kirwan JP: TNF- α impairs insulin signaling and insulin stimulation of glucose uptake in C2C12 muscle cells. *Am J Physiol* 276:E849–E855, 1999
9. Nolte LA, Hansen PA, Chen MM, Schluter JM, Gulve EA, Holloszy JO: Short-term exposure to tumor necrosis factor- α does not affect insulin-stimulated glucose uptake in skeletal muscle. *Diabetes* 47:721–726, 1998
10. Patiag D, Gray S, Idris I, Donnelly R: Effects of tumour necrosis factor- α and inhibition of protein kinase C on glucose uptake in L6 myoblasts. *Clin Sci (Lond)* 99:303–307, 2000
11. Azzi A, Boscoboinik D, Hensey C: The protein kinase C family. *Eur J Biochem* 208:547–557, 1992
12. Haller H, Lindschau C, Luft FC: Role of protein kinase C in intracellular signaling. *Ann N Y Acad Sci* 733:313–320, 1994
13. Li W, Mischak H, Yu JC, Wang LM, Mushinski JF, Heidaran MA, Pierce JH: Tyrosine phosphorylation of protein kinase C- δ in response to its activation. *J Biol Chem* 269:2349–2352, 1994
14. Soltoff SP, Toker A: Carbachol, substance P, and phorbol ester promote the tyrosine phosphorylation of protein kinase C δ in salivary gland epithelial cells. *J Biol Chem* 270:13490–13495, 1995
15. Braiman L, Sheffi-Friedman L, Bak A, Tennenbaum T, Sampson SR: Tyrosine phosphorylation of specific protein kinase C isoenzymes participates in insulin stimulation of glucose transport in primary cultures of rat skeletal muscle. *Diabetes* 48:1922–1929, 1999
16. Gschwendt M: Protein kinase C delta. *Eur J Biochem* 259:555–564, 1999
17. Schutze S, Nottrott S, Pfizenmaier K, Kronke M: Tumor necrosis factor signal transduction: cell-type-specific activation and translocation of protein kinase C. *J Immunol* 144:2604–2608, 1990
18. Prasanna G, Dibas A, Brown K, Yorio T: Activation of protein kinase C by tumor necrosis factor- α in human non-pigmented ciliary epithelium. *J Ocul Pharmacol Ther* 14:401–412, 1998
19. Miura A, Ishizuka T, Kanoh Y, Ishizawa M, Itaya S, Kimura M, Kajita K, Yasuda K: Effect of tumor necrosis factor- α on insulin signal transduction in rat adipocytes: relation to PKC β and zeta translocation. *Biochim Biophys Acta* 1449:227–238, 1999
20. Kellerer M, Mushack J, Mischak H, Haring HU: Protein kinase C (PKC) epsilon enhances the inhibitory effect of TNF α on insulin signaling in HEK293 cells. *FEBS Lett* 418:119–122, 1997
21. Cooper DR, de Ruiz G, Fanjul LF, Mojsilovic L, Standaert ML, Pollet RJ, Farese RV: Insulin but not phorbol ester treatment increases phosphorylation of vinculin by protein kinase C in BC3H-1 myocytes. *FEBS Lett* 214:122–126, 1987
22. Srinivasan M, Begum N: Stimulation of protein phosphatase-1 activity by phorbol esters: evaluation of the regulatory role of protein kinase C in insulin action. *J Biol Chem* 269:16662–16667, 1994
23. Bandyopadhyay G, Standaert ML, Galloway L, Moscat J, Farese RV: Evidence for involvement of protein kinase C (PKC)-zeta and noninvolvement of diacylglycerol-sensitive PKCs in insulin-stimulated glucose transport in L6 myotubes. *Endocrinology* 138:4721–4731, 1997
24. Chalfant CE, Ohno S, Konno Y, Fisher AA, Bisnauth LD, Watson JE, Cooper DR: A carboxy-terminal deletion mutant of protein kinase C beta II inhibits insulin-stimulated 2-deoxyglucose uptake in L6 rat skeletal muscle cells. *Mol Endocrinol* 10:1273–1281, 1996
25. Braiman L, Alt A, Kuroki T, Ohba M, Bak A, Tennenbaum T, Sampson SR: Protein kinase C delta mediates insulin-induced glucose transport in primary cultures of rat skeletal muscle. *Mol Endocrinol* 13:2002–2012, 1999
26. Braiman L, Alt A, Kuroki T, Ohba M, Bak A, Tennenbaum T, Sampson SR: Insulin induces specific interaction between insulin receptor and PKC in primary cultured skeletal muscle. *Mol Endocrinol* 15:565–574, 2001
27. Kellerer M, Mushack J, Seffer E, Mischak H, Ullrich A, Haring HU: Protein kinase C isoforms alpha, delta and theta require insulin receptor substrate-1 to inhibit the tyrosine kinase activity of the insulin receptor in human kidney embryonic cells (HEK 293 cells). *Diabetologia* 41:833–838, 1998
28. Paz K, Boura-Halfon S, Wyatt LS, LeRoith D, Zick Y: The juxtamembrane but not the carboxyl-terminal domain of the insulin receptor mediates insulin's metabolic functions in primary adipocytes and cultured hepatoma cells. *J Mol Endocrinol* 24:419–432, 2000
29. Hotamisligil GS, Peraldi P, Budavari A, Ellis R, White MF, Spiegelman BM: IRS-1-mediated inhibition of insulin receptor tyrosine kinase activity in TNF- α - and obesity-induced insulin resistance. *Science* 271:665–668, 1996
30. Hotamisligil GS: Mechanisms of TNF- α -induced insulin resistance [see comments]. *Exp Clin Endocrinol Diabetes* 107:119–125, 1999
31. Brodie C, Bak A, Shainberg A, Sampson SR: Role of Na-K ATPase in regulation of resting membrane potential of cultured rat skeletal myotubes. *J Cell Physiol* 130:191–198, 1987
32. Sampson SR, Brodie C, Alboim SV: Role of protein kinase C in insulin activation of the Na-K pump in cultured skeletal muscle. *Am J Physiol* 266:C751–C758, 1994
33. Brodie C, Brody M, Sampson SR: Characterization of the relation between sodium channels and electrical activity in cultured rat skeletal myotubes: regulatory aspects. *Brain Res* 488:186–194, 1989
34. Sampson SR, Brodie C, Alboim SV: Role of protein kinase C in insulin activation of the Na-K pump in cultured skeletal muscle. *Am J Physiol* 266:C751–C758, 1994
35. Miyake S, Makimura M, Kanegae Y, Harada S, Sato Y, Takamori K, Tokuda C, Saito I: Efficient generation of recombinant adenoviruses using adenovirus DNA-terminal protein complex and a cosmid bearing the full-length virus genome. *Proc Natl Acad Sci U S A* 93:1320–1324, 1996
36. Ohno S, Konno Y, Akita Y, Yano A, Suzuki K: A point mutation at the putative ATP-binding site of protein kinase C alpha abolishes the kinase activity and renders it down-regulation-insensitive: a molecular link between autophosphorylation and down-regulation. *J Biol Chem* 265:6296–6300, 1990
37. Hotamisligil GS, Murray DL, Choy LN, Spiegelman BM: Tumor necrosis factor alpha inhibits signaling from the insulin receptor. *Proc Natl Acad Sci U S A* 91:4854–4858, 1994
38. Osada S, Mizuno K, Saido TC, Suzuki K, Kuroki T, Ohno S: A new member of the protein kinase C family, nPKC theta, predominantly expressed in skeletal muscle. *Mol Cell Biol* 12:3930–3938, 1992
39. Cooper DR, Watson JE, Hernandez H, Yu B, Standaert ML, Ways DK, Arnold TT, Ishizuka T, Farese RV: Direct evidence for protein kinase C involvement in insulin-stimulated hexose uptake. *Biochem Biophys Res Commun* 188:142–148, 1992
40. Avignon A, Standaert ML, Yamada K, Mischak H, Spencer B, Farese RV: Insulin increases mRNA levels of protein kinase C-alpha and -beta in rat adipocytes and protein kinase C-alpha, -beta and -theta in rat skeletal muscle. *Biochem J* 308:181–187, 1995
41. Danielsen AG, Liu F, Hosomi Y, Shii K, Roth RA: Activation of protein kinase C alpha inhibits signaling by members of the insulin receptor family. *J Biol Chem* 270:21600–21605, 1995
42. Chin JE, Liu F, Roth RA: Activation of protein kinase C alpha inhibits insulin-stimulated tyrosine phosphorylation of insulin receptor substrate-1. *Mol Endocrinol* 8:51–58, 1994
43. Formisano P, Oriente F, Miele C, Caruso M, Auricchio R, Vigliotta G, Condorelli G, Beguinot F: In NIH-3T3 fibroblasts, insulin receptor interaction with specific protein kinase C isoforms controls receptor intracellular routing. *J Biol Chem* 273:13197–13202, 1998
44. Di Guglielmo GM, Drake PG, Baass PC, Authier F, Posner BI, Bergeron JJ: Insulin receptor internalization and signalling. *Mol Cell Biochem* 182:59–63, 1998
45. Sawai H, Okazaki T, Takeda Y, Tashima M, Sawada H, Okuma M, Kishi S, Umehara H, Domaie N: Ceramide-induced translocation of protein kinase C-delta and -epsilon to the cytosol: implications in apoptosis. *J Biol Chem* 272:2452–2458, 1997
46. Franz MG, Norman JG, Fabri PJ, Gower WRJ: Differentiation of pancreatic ductal carcinoma cells associated with selective expression of protein kinase C isoforms. *Ann Surg Oncol* 3:564–569, 1996
47. Brodie C, Kuperstein I, Acs P, Blumberg PM: Differential role of specific PKC isoforms in the proliferation of glial cells and the expression of the astrocytic markers GFAP and glutamine synthetase. *Brain Res Mol Brain Res* 56:108–117, 1998