

# Hypothalamic Pro-Opiomelanocortin mRNA Is Reduced By Fasting in *ob/ob* and *db/db* Mice, but Is Stimulated by Leptin

Tooru M. Mizuno, Steven P. Kleopoulos, Hugo T. Bergen, James L. Roberts, Catherine A. Priest, and Charles V. Mobbs

**Reduction in the activity of the  $\alpha$ -melanocyte-stimulating hormone ( $\alpha$ -MSH) system causes obesity, and infusions of  $\alpha$ -MSH can produce satiety, raising the possibility that  $\alpha$ -MSH may mediate physiological satiety signals. Since  $\alpha$ -MSH is coded for by the pro-opiomelanocortin (POMC) gene, we examined if POMC gene expression would be inhibited by fasting in normal mice or in models of obesity characterized by leptin insufficiency (*ob/ob*) or leptin insensitivity (*db/db*). In wild-type mice, hypothalamic POMC mRNA was decreased >60% after a 2-day fast and was positively correlated with leptin mRNA. Similarly, compared with controls, POMC mRNA was decreased by at least 60% in both *db/db* and *ob/ob* mice. POMC mRNA was negatively correlated with both neuropeptide Y (NPY) and melanin-concentrating hormone (MCH) mRNA. Finally, treatment of both male and female *ob/ob* mice with leptin stimulated hypothalamic POMC mRNA by about threefold. These results suggest that impairment in production, processing, or responsiveness to  $\alpha$ -MSH may be a common feature of obesity and that hypothalamic POMC neurons, stimulated by leptin, may constitute a link between leptin and the melanocortin system. *Diabetes* 47:294–297, 1997**

**T**he obesity of yellow  $A^y$  mice is thought to be due to ectopic expression of the product of the agouti gene (1), an antagonist of the action of  $\alpha$ -melanocyte-stimulating hormone ( $\alpha$ -MSH) acting through the melanocortin-4 (MC-4) receptor (2). Furthermore, disruption of the MCR-4 receptor (3) causes obesity. In addition,  $\alpha$ -MSH agonists and antagonists can promote satiety and feeding, respectively (4). Therefore  $\alpha$ -MSH is a plausible candidate as a physiological mediator of satiety. Obese  $A^y$  mice exhibit elevated expression of the gene coding for the hormone

leptin (5). Mutations in leptin cause obesity (6), experimental elevation of leptin reduces body weight and induces satiety (7), and expression of leptin is reduced during fasting (5). The elevation of leptin in obese  $A^y$  mice suggests that these mice are resistant to the satiety effects of leptin; in turn, since the obesity of  $A^y$  mice appears to be due to antagonism of  $\alpha$ -MSH, these data suggest the hypothesis that  $\alpha$ -MSH mediates effects of leptin. In contrast to the regulation of neuropeptide Y (NPY) mRNA, which is inhibited by leptin (8), these observations suggest that the synthesis and/or release of  $\alpha$ -MSH might be stimulated by leptin, and, conversely, inhibited in fasting mice and in *db/db* and *ob/ob* mice, which are resistant to and deficient in leptin, respectively.  $\alpha$ -MSH is produced by proteolytic cleavage and modification of pro-opiomelanocortin (POMC). Therefore, the present study assessed the hypothesis that in models characterized by low leptin (fasting and *ob/ob* mice) and leptin resistance (*db/db*), hypothalamic POMC mRNA would be reduced, and, conversely, that hypothalamic POMC mRNA would be elevated after leptin administration. Such a result would be consistent with the hypothesis that POMC gene products, in particular  $\alpha$ -MSH, may mediate some effects of leptin and other satiety factors.

## RESEARCH DESIGN AND METHODS

**Animals.** C57Bl/6J, CBA, *db/db* (along with control C57Bl/KsJ), and *ob/ob* (along with control C57Bl/6J) mice were obtained at 2 months of age from The Jackson Laboratory. Mice were individually housed with free access to feed and water under 12:12 h light-dark cycle (lights on at 0700). All studies had been approved by the appropriate Institutional Animal Review Board. To assess effects of nutritional status, mice were fasted for 48 h or fed ad lib ( $n = 4-12$  per group); in some cases, food was removed from cages of ad lib fed mice for 7 h during the day to ensure acute nutritional status was equivalent in all mice. To assess effects of leptin replacement, male *ob/ob* mice and wild-type controls were injected intraperitoneally with 3  $\mu$ g/g body wt daily for 5 days; in this study, POMC mRNA was assessed by Northern blots. In a second study, female *ob/ob* mice were either fed ad lib or fasted for 48 h; during the period of fasting, the ad lib fed mice and the fasting mice were injected every 12 h with saline or leptin (0.5  $\mu$ g/g body wt, i.p.) with the final, fifth injection 30 min before being killed; in this study POMC mRNA was assessed by in situ hybridization. Mice were killed just before lights out (except for the mice in the leptin replacement study, who were killed 2 h after lights on) under anesthesia by CO<sub>2</sub> narcosis. Brain blocks were removed, frozen on dry ice and stored at  $-70^{\circ}\text{C}$  until sectioning on a cryostat or preparation of RNA.

**Template production and probe labeling.** POMC template was prepared by amplifying a source plasmid DNA using polymerase chain reaction (PCR) with N-terminal primer: 5'-CCTGTGAAGGTGTACCCCAATGTC-3' and C-terminal primer: 5'-CAGTTCTTGATGATGGCGTTC-3'. NPY template was prepared in the same way with N-terminal primer: 5'-CTAGGTAACAACGAATGGGG-3' and C-terminal primer: 5'-CCACATGGAAGGGTCTTCAAG-3'. MCH template was prepared from mouse hypothalamic RNA by reverse transcription-polymerase chain reaction (RT-PCR) with N-terminal primer: 5'-CCAAGTCCATCAGGAACGTA-3' and C-terminal primer: 5'-TTGTGGACCAGCAGGTATCA-3'. Leptin template was prepared as described previously (5). The amplified fragments were gel-purified, diluted to 50 ng/ $\mu$ l and stored at  $-20^{\circ}\text{C}$ . Single-stranded internally labeled DNA

From the Fishberg Center for Neurobiology (T.M.M., S.P.K., C.A.P., J.L.R., C.V.M.), the Neurobiology of Aging Laboratories, and the Department of Geriatrics (T.M.M., S.P.K., C.V.M.), Mount Sinai School of Medicine, New York, New York; and the Department of Anatomy (H.T.B.), University of Manitoba, Winnipeg, Manitoba, Canada.

Address correspondence and reprint requests to Dr. Charles V. Mobbs, Neurobiology of Aging Laboratories, Box 1639, Mt. Sinai School of Medicine, 1425 Madison Ave., New York, NY 10029-6574. E-mail: mobbsc@alum.mit.edu.

Received for publication 2 October 1997 and accepted in revised form 11 November 1997.

$\alpha$ -MSH,  $\alpha$ -melanocyte-stimulating hormone; MCH, melanin-concentrating hormone; NPY, neuropeptide Y; POMC, pro-opiomelanocortin.

probes were produced by amplified primer extension labeling by using only C-terminal primer (5).

**Northern blot analysis.** RNA from mediobasal hypothalamus (microdissected as described previously [9]) was extracted in TRIzol (GIBCO BRL, Gaithersburg, MD) and 3 µg of total RNA (determined by UV absorbance corroborated by ethidium bromide-stained integrity gels) was subjected to Northern blot analysis to detect POMC mRNA and NPY mRNA in the hypothalamus, and leptin mRNA in adipose tissue from gonadal fat pad. Northern blot analysis was performed as described previously (5). To monitor RNA loading, membranes were reprobed and hybridized with <sup>32</sup>P-labeled probe encoding 18S ribosomal RNA to ensure equal loading in all lanes.

**In situ hybridization histochemistry.** Frozen coronal sections (10 µm thick) through the mouse hypothalamus were cut, fixed in 3% paraformaldehyde in 0.1 mol/l phosphate buffer (pH 7.0) containing 0.03% DEPC, dehydrated, and stored at -20°C until use. Sorting on the basis of histology was carried out to ensure sections were matched for anterior-posterior level. Sections were prehybridized in 2 X SSC, 5 mmol/l EDTA, 2.5 X Denhard's solution, 5 mmol/l DTT, 100 mg/ml hering sperm DNA, 100 mg/ml yeast tRNA, 5 mg/ml single-stranded calf thymus DNA and 50% deionized formamide for 2 h at 42°C. Hybridization was carried out in the same buffer containing 10% dextran sulfate and labeled probe (<sup>32</sup>P: 2 X 10<sup>6</sup> dpm per 20 µl per section) at 42°C overnight. Sections (two sections per matched A-P level) were washed twice in 1 X SSC for 15 min and in 0.1 X SSC overnight at room temperature, followed by a final wash in 0.1 X SSC for 1 h at 55°C. Slides were dehydrated, air dried, and apposed to autoradiography film.





**Data analysis.** The total integrated densities of hybridization signals were determined by computerized densitometric scanning (MCID System, St. Catherine's, Ontario, Canada) as described previously (10). Statistical analysis was performed by analysis of variance (ANOVA) followed by Tukey-Kramer pair-wise comparisons, or by linear regression, with criterion for significance as  $P < 0.05$ .

## RESULTS

As assessed by Northern blots in C57Bl/6J mice, a 48-h fast significantly reduced hypothalamic POMC mRNA by ~60% (Fig. 1A, D) and at the same time induced hypothalamic NPY mRNA by about fourfold (Fig. 1B, E), with no change in the 18S ribosomal RNA band (Fig. 1C, F). In these mice, leptin mRNA was reduced ~80% by the 48-h fast (FED: 100 ± 9%; FASTED: 16 ± 2%, expressed as percentage of fed controls). POMC mRNA was positively correlated with leptin mRNA ( $r = 0.89$ ,  $P < 0.01$ ), and negatively correlated with NPY mRNA ( $r = -0.88$ ,  $P < 0.01$ ). These data were corroborated in a separate study and using an RNase protection assay for POMC mRNA, which also demonstrated a 60% decrease by fasting (data not shown). The effects of fasting were also corroborated by in situ hybridization in a separate study in which fasting reduced POMC mRNA by over 80% throughout the arcuate nucleus (Fig. 1G). In the same mice, NPY was increased over 300% (Fig. 1H), and leptin mRNA was reduced by fasting over 90% (FED: 100 ± 11%; FASTED: 7.3 ± 2%).

Leptin resistance in nonfasted *db/db* mice was associated with significant reduction in POMC mRNA of ~80%, compared with nonfasted wild-type C57/KsJ mice (Fig. 2A-C). In wild-type KsJ mice, fasting (Fig. 2C) was associated with significant decreases in POMC mRNA. In *db/db* mice, fasting was associated with a further significant reduction in POMC mRNA (Fig. 2C). NPY mRNA was elevated in *db/db* mice compared with wild-type controls under all conditions, and, interestingly, fasting caused a further elevation in *db/db* mice (Fig. 2D). Similarly, MCH mRNA was significantly elevated in every circumstance in which POMC mRNA was reduced (Fig. 2E). POMC mRNA was significantly negatively correlated with both NPY mRNA ( $r = -0.79$ ,  $P < 0.001$ ) and MCH mRNA ( $r = -0.59$ ,  $P < 0.05$ ), while NPY mRNA and MCH mRNA were significantly positively correlated with each other ( $P = 0.46$ ,  $P < 0.05$ ).

As assessed by Northern blot analysis, hypothalamic POMC mRNA was reduced by ~90% in leptin-deficient nonfasted *ob/ob* male mice, compared with wild-type controls (Fig. 3A). Injection of leptin over 5 days significantly ele-



**FIG. 1.** Effect of a 48-h fast on hypothalamic POMC (A), NPY (B), and 18S RNA (C). Quantification of Northern blots shown in A-C: POMC (D), NPY (E), or 18S RNA (F) band, mean ± SE, expressed as percentage of fed controls. Quantification of POMC (G) or NPY (H) mRNA signal after in situ hybridization; in fed or fasted mice;  $n = 4-12$  per group. Abscissa in G and H indicates number of microns from beginning of arcuate nucleus, thus representing almost the entire extent on the nucleus in mice. \*Significant ( $P < 0.05$ ) effect of fast versus comparable fed.

vated hypothalamic POMC mRNA in *ob/ob* mice (Fig. 3A), although at these doses over this period of time, POMC mRNA was not restored completely to wild-type levels. As expected, NPY mRNA was elevated in *ob/ob* mice, but like POMC mRNA, at this dose and period of injection, NPY mRNA was only partially corrected (Fig. 3B). These results were corroborated in fed and fasted female *ob/ob* mice, in which leptin injection elevated hypothalamic POMC mRNA to about the same degree as observed in male mice as assessed by Northern blots (Fig. 3C).




FIG. 2. Effect of *db/db* genotype or fasting on POMC, NPY, and MCH mRNA. *A* and *B*: POMC mRNA, indicating in situ hybridization in groups as indicated. Quantification of POMC (*C*), NPY (*D*), or MCH mRNA (*E*), mean  $\pm$  SE,  $n = 6-8$ /group, expressed as percentage of fed, wild-type controls. Groups with different letters are statistically different ( $P < 0.05$ ).

**DISCUSSION**

In the present studies, two forms of genetic obesity (*db/db* and *ob/ob*) were associated with decreased hypothalamic POMC mRNA. Since mutations in the leptin gene have now been associated with obesity in humans (11), and mutations in the leptin receptor also produce obesity in rats as well as in *db/db* mice (12), it will be of interest to assess if hypothalamic POMC mRNA is also reduced in genetic obesity of these species. Mice in which the basic helix-loop-helix *Nhlh2* gene has been deleted become obese, and these mice also exhibit reduced hypothalamic POMC mRNA, whereas other hypothalamic gene products such as NPY, tyrosine hydroxylase, galanin, and arginine vasopressin were not influenced in these




FIG. 3. Effect of *ob/ob* genotype and leptin replacement on POMC (*A*) and NPY (*B*) mRNA levels, as assessed by Northern blot analysis in male mice, and POMC mRNA in fed and fasted *ob/ob* female mice (*C*). Mean  $\pm$  SE,  $n = 4-8$ /group, expressed as percentage of wild-type, saline-injected (Sal) mice (*A* and *B*), or percentage of fed *ob/ob* saline-injected mice (*C*). Groups with different letters are statistically different ( $P < 0.05$ , ANOVA, followed by Tukey-Kramer).

obese mice (13). Another form of genetic human obesity entails mutation of the PC-1 peptide processing enzyme which leads (among other impairments) to impaired ability to process the POMC precursor peptide into its components (14), presumably leading to impaired production of  $\alpha$ -MSH. Similarly, the obesity of the *fatty/fatty* mouse is due to a mutation in another processing enzyme, carboxypeptidase E (15), which leads (among other impairments) to impaired processing of the POMC precursor (16). Finally, the obesity of the yellow *A<sup>y</sup>* mouse entails overproduction (1) of an antagonist of the action of  $\alpha$ -MSH (2). Similarly, disruption of the MC-4 receptor through which  $\alpha$ -MSH acts also causes obesity (3).

Several forms of acquired obesity are also associated with decreased hypothalamic POMC mRNA. A syndrome of obesity produced by distemper-related virus is associated with reduced hypothalamic POMC mRNA (17). Finally, the process of aging, particularly in rats, is associated with the development of obesity, and hypothalamic POMC mRNA decreases during aging (18). Taken together, these studies indicate that impaired production, processing, or sensitivity to  $\alpha$ -MSH is a feature common to many forms of obesity.

The present study suggests that the effect of fasting to reduce hypothalamic POMC mRNA is robust, since results were similar in several strains of wild-type mice, consistent with previous studies in rats (19–20). The following evidence from the present study suggested that the effect of fasting to reduce POMC mRNA is mediated in part by a fall in leptin: 1) leptin decreased with fasting in normal mice, and POMC mRNA was positively correlated with leptin; 2) leptin resistance (in *db/db* mice) was associated with reduced hypothalamic POMC mRNA; 3) leptin insufficiency (in *ob/ob* mice) was associated with reduced POMC mRNA; and 4) leptin injection significantly stimulated hypothalamic POMC mRNA in *ob/ob* mice. The effect of leptin replacement was independent of effects on food intake and body weight, since the effects were comparable in fed and in fasted *ob/ob* mice, and leptin did not significantly reduce body weight or plasma glucose beyond the effect of fasting (not shown). The hypothesis that hypothalamic POMC gene products mediate some effects of leptin is consistent with other reports, including the following: 1) in agouti mice, in which the basis of the obesity is an antagonism of  $\alpha$ -MSH, leptin is elevated (5), suggesting a leptin resistance in agouti mice which has been recently directly demonstrated (21); 2) the peri-arcuate nuclear region, the main site of POMC synthesis (22) in the brain, is the area most sensitive to the satiety effects of leptin (23); and 3) leptin receptors colocalize with POMC-containing neurons (24). Nevertheless, since POMC mRNA was reduced (and NPY was elevated) by fasting in *db/db* mice, it is plausible that other factors, such as insulin and glucose, also stimulate POMC mRNA and inhibit NPY mRNA. The present studies, in which MCH mRNA was elevated in *db/db* and fasted mice, were also consistent with a role for MCH, which stimulates feeding and is elevated in *ob/ob* mice (25), in fasting and obesity. MCH can antagonize the effect of  $\alpha$ -MSH on behavior (26). Therefore, a plausible hypothesis is that products of the POMC gene and the MCH gene are inversely expressed and act in opposition (though possibly through different receptors) to regulate body weight. Such a hypothesis suggests that  $\alpha$ -MSH agonists may, like leptin (27), reverse or attenuate some effects of fasting.

#### ACKNOWLEDGMENTS

This work was supported by a grant from the National Institutes of Health (DK-50110) to C.V.M.

#### REFERENCES

- Bultman SJ, Michaud EJ, Woychik RP: Molecular characterization of the mouse agouti locus. *Cell* 71:1195–1204, 1992
- Lu D, Willard D, Patel IR, Kadwell S, Overton L, Kost T, Luther M, Chen W, Woychik RP, Wilkison WO, Cone RD: Agouti protein is an antagonist of the melanocyte-stimulating-hormone receptor. *Nature* 371:799–802, 1994
- Huszar D, Lynch CA, Fairchild-Huntress V, Dunmore JH, Fang Q, Berkemeier LR, Gu W, Kesterson RA, Boston BA, Cone RD, Smith FJ, Campfield LA, Burn P, Lee F: Targeted disruption of the melanocortin-4 receptor results in obesity in mice. *Cell* 88:131–141, 1997
- Fan W, Boston BA, Kesterson RA, Hruby VJ, Cone RD: Role of melanocortin-ergic neurons in feeding and the agouti obesity syndrome. *Nature* 385:165–168, 1997
- Mizuno TM, Bergen H, Funabashi T, Kleopoulos SP, Zhong YG, Bauman WA, Mobbs CV: Obese gene expression: reduction by fasting and stimulation by insulin and glucose in lean mice, and persistent elevation in acquired (diet-induced) and genetic (yellow agouti) obesity. *Proc Natl Acad Sci USA* 93:3434–3438, 1996
- Zhang Y, Proenca R, Maffei M, Barone M, Leopold L, Friedman JM: Positional cloning of the mouse obese gene and its human homologue. *Nature* 372:425–432, 1994
- Halaas JL, Gajiwala KS, Maffei M, Cohen SL, Chait BT, Rabinowitz D, Lallone RL, Burley SK, Friedman JM: Weight-reducing effects of the plasma protein encoded by the obese gene. *Science* 269:543–546, 1995
- Schwartz MW, Baskin DG, Bukowski TR, Kuijper JL, Foster D, Lasser G, Prunkard DE, Porte D Jr, Woods SC, Seeley RJ, Weigle DS: Specificity of leptin action on elevated blood glucose levels and hypothalamic neuropeptide Y gene expression in *ob/ob* mice. *Diabetes* 45:531–535, 1996
- Mobbs CV, Fink G, Pfaff DW: HIP-70: a protein induced by estrogen in the brain and LH-RH in the pituitary. *Science* 247:1477–1479, 1990
- Bergen HT, Mobbs CV: Ventromedial hypothalamic lesions produced by gold thioglucose do not impair induction of fasting. *Brain Res* 707:266–271, 1996
- Montague CT, Farooqi IS, Whitehead JP, Soos MA, Rau H, Wareham NJ, Sewter CP, Digby JE, Mohammed SN, Hurst JA, Cheetham CH, Earley AR, Barnett AH, Prins JB, O'Rahilly S: Congenital leptin deficiency is associated with severe early-onset obesity in humans. *Nature* 387:903–908, 1997
- Chua SC Jr, White DW, Wu-Peng XS, Liu SM, Okada N, Kershaw EE, Chung WK, Power-Keohoe L, Chua M, Tartaglia LA, Leibel RL: Phenotype of fatty due to *Gln269Pro* mutation in the leptin receptor (*Lepr*). *Diabetes* 45:1141–1143, 1996
- Good DJ, Porter FD, Mahon KA, Parlow AF, Westphal H, Kirsch IR: Hypogonadism and obesity in mice with a targeted deletion of the *Nhlh2* gene. *Nat Genet* 15:397–401, 1997
- Jackson RS, Creemers JWM, Ohagi S, Raffin-Sanson M-L, Sanders L, Montague CT, Hutton JC, O'Rahilly S: Obesity and impaired prohormone processing associated with mutations in the human prohormone convertase 1 gene. *Nature Genetics* 16:303–306, 1997
- Naggert JK, Fricker LD, Varlamov O, Nishina PM, Rouille Y, Steiner DF, Carroll RJ, Paigen BJ, Leiter EH: Hyperproinsulinaemia in obese fat/fat mice associated with a carboxypeptidase E mutation which reduces enzyme activity. *Nature Genetics* 10:135–142, 1995
- Cool DR, Normant E, Shen F, Chen HC, Pannell L, Zhang Y, Loh YP: Carboxypeptidase E is a regulated secretory pathway sorting receptor: genetic obliteration leads to endocrine disorders in *Cpefat* mice. *Cell* 88:73–83, 1997
- Nagashima K, Zabriskie JB, Lyons MJ: Virus-induced obesity in mice: association with a hypothalamic lesion. *J Neuropathol Exp Neurol* 51:101–109, 1992
- Lloyd JM, Scarbrough K, Weiland NG, Wise PM: Age-related changes in proopiomelanocortin (POMC) gene expression in the periaqueductal region of ovariectomized rats. *Endocrinology* 129:1896–1902, 1991
- Kim EM, Welch CC, Grace MK, Billington CJ, Levine AS: Chronic food restriction and acute food deprivation decrease mRNA levels of opioid peptides in arcuate nucleus. *Am J Physiol* 270:R1019–R1024, 1996
- Brady LS, Smith MA, Gold PW, Herkenham M: Altered expression of hypothalamic neuropeptide mRNAs in food-restricted and food-deprived rats. *Neuroendocrinology* 52:441–447, 1990
- Halaas JL, Boozer C, Blair-West J, Fidathusein N, Denton DA, Friedman JM: Physiological response to long-term peripheral and central leptin infusion in lean and obese mice. *Proc Natl Acad Sci USA* 94:8878–8883, 1997
- Gee CE, Chen CL, Roberts JL, Thompson R, Watson SJ: Identification of proopiomelanocortin neurones in rat hypothalamus by in situ cDNA-mRNA hybridization. *Nature* 306:374–376, 1983
- Satoh N, Ogawa Y, Katsuura G, Hayase M, Tsuji T, Imagawa K, Yoshimasa Y, Nishi S, Hosoda K, Nakao K: The arcuate nucleus as a primary site of satiety effect of leptin in rats. *Neurosci Lett* 224:149–152, 1997
- Cheung CC, Thornton JE, Clifton DK, Steiner RA: Coexpression of leptin receptor (OB-R) mRNA with POMC and NPY in the central nervous system of the rat (Abstract). *Endocr Soc Abstr* 79:379, 1997
- Qu D, Ludwig DS, Gammeltoft S, Piper M, Pellemounter MA, Cullen MJ, Mathes WF, Przypek R, Kanarek R, Maratos-Flier E: A role for melanin-concentrating hormone in the central regulation of feeding behavior. *Nature* 380:243–247, 1996
- Sanchez M, Baker BI, Celis M: Melanin-concentrating hormone (MCH) antagonizes the effects of  $\alpha$ -MSH and neuropeptide E-I on grooming and locomotor activities in the rat. *Peptides* 18:393–396, 1997
- Ahima RS, Prabakaran D, Mantzoros C, Qu D, Lowell B, Maratos-Flier E, Flier JS: Role of leptin in the neuroendocrine response to fasting. *Nature* 382:250–252, 1996