

# *FUT2* Nonsecretor Status Links Type 1 Diabetes Susceptibility and Resistance to Infection

Deborah J. Smyth, Jason D. Cooper, Joanna M.M. Howson, Pamela Clarke, Kate Downes, Trupti Mistry, Helen Stevens, Neil M. Walker, and John A. Todd

**OBJECTIVE**—*FUT2* encodes the  $\alpha(1,2)$  fucosyltransferase that determines blood group secretor status. Homozygotes (A/A) for the common nonsense mutation rs601338A>G (W143X) are nonsecretors and are unable to express histo-blood group antigens in secretions and on mucosal surfaces. This mutation has been reported to provide resistance to Norovirus and susceptibility to Crohn's disease, and hence we aimed to determine if it also affects risk of type 1 diabetes.

**RESEARCH DESIGN AND METHODS**—rs601338A>G was genotyped in 8,344 patients with type 1 diabetes, 10,008 control subjects, and 3,360 type 1 diabetic families. Logistic regression models were used to analyze the case-control collection, and conditional logistic regression was used to analyze the family collection.

**RESULTS**—The nonsecretor A/A genotype of rs601338A>G was found to confer susceptibility to type 1 diabetes in both the case-control and family collections (odds ratio for AA 1.29 [95% CI 1.20–1.37] and relative risk for AA 1.22 [95% CI = 1.12–1.32]; combined  $P = 4.3 \times 10^{-18}$ ), based on a recessive effects model.

**CONCLUSIONS**—Our findings linking *FUT2* and type 1 diabetes highlight the intriguing relationship between host resistance to infections and susceptibility to autoimmune disease. *Diabetes* 60:3081–3084, 2011

**T**ype 1 diabetes results from the complex interaction between genetic predisposition, the immune system, and the environment (1). The incidence of the autoimmune disease type 1 diabetes has increased dramatically over the last 25 years, which cannot be explained by genetic variation in the population. This increase in type 1 diabetes incidence is most likely due to the changing influence of many environmental factors—such as toxins, cow's milk, vitamin D, and microbial infections—acting on an underlying genetic susceptibility. One hypothesis is that our increasingly hygienic, healthy environment has altered the microbiomes of individuals at risk and the immune system's level of development and regulation such that in a higher proportion of children it is imbalanced away from regulation and toward autoimmunity (1). Accordingly,

From the Juvenile Diabetes Research Foundation/Wellcome Trust Diabetes and Inflammation Laboratory, Department of Medical Genetics, Cambridge Institute for Medical Research, University of Cambridge, Addenbrooke's Hospital, Cambridge, U.K.

Corresponding author: John A. Todd, john.todd@cimr.cam.ac.uk.

Received 13 May 2011 and accepted 28 July 2011.

DOI: 10.2337/db11-0638

This article contains Supplementary Data online at <http://diabetes.diabetesjournals.org/lookup/suppl/doi:10.2337/db11-0638/-DC1>.

D.J.S. and J.D.C. contributed equally to this work.

© 2011 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

See accompanying commentary, p. 2685.

one would expect to find genes that are involved directly in host resistance to infectious disease among type 1 diabetes susceptibility genes (1,2).

The *FUT2* gene on chromosome 19q13.33 codes for the  $\alpha(1,2)$  fucosyltransferase responsible for the synthesis of the H antigen, which is the precursor of the ABO histo-blood group antigens in body fluids and on the intestinal mucosa. Subjects homozygous for *FUT2* null alleles are nonsecretors (se) who do not express ABO antigens in saliva and the gastrointestinal tract. Nonsecretor status results from being homozygous for one of two nonsecretor variants: se<sup>428</sup>, which codes for a stop codon at position 143 (W143X rs601338A>G) in Europeans and Africans; and se<sup>385</sup> (I129F, rs1047781A>T) in South East and East Asians.

The H antigen secretion depends on additional glycosyltransferases, including the Lewis (*FUT3*) A/B enzymes, found in epithelial cells and erythrocytes. Both the Lewis and ABO genes are highly polymorphic, giving rise to null phenotypes, which are thought to be related to evolutionary changes for host-pathogen interactions because many pathogens use surface glycoproteins in host evasion (3).

The genetic inability to secrete ABO blood group antigens in body fluids has been associated with a variety of infectious diseases, in particular with susceptibility to *Candida albicans* (4,5) and *Streptococcus* infection (6). However, *FUT2* secretors are more susceptible to *Helicobacter pylori* (7). Additionally, the *FUT2* 143X/rs601338A allele has also been implicated in higher circulating serum vitamin B12 levels (8) and slower progression of HIV (9); and most intriguingly, it provides resistance to Norovirus infection (Norwalk virus) (10–12). Nonsecretors of *FUT2* have been shown to have almost complete protection to GGII Noroviruses, which is a major cause of acute gastroenteritis worldwide, because they do not express the H type-1 oligosaccharide ligand required for GGII Norovirus binding (10–12).

Recently, a genome-wide association (GWA) study (13) and a meta-analysis of GWA studies (14) identified the *FUT2* region as a Crohn's disease locus. In the GWA study, McGovern et al. (13) reported several single nucleotide polymorphisms (SNPs) in high linkage disequilibrium ( $r^2 > 0.80$ ) with evidence of association with Crohn's disease: rs492602, rs601338 (W143X), rs602662 (S258G), and rs485186. In the meta-analysis, Franke et al. (14) reported an association with rs281379 (odds ratio for allele A 1.07 [95% CI 1.04–1.11];  $P = 7.4 \times 10^{-12}$ );  $r^2$  between rs281379 and the nonsecretor SNP rs601338 was 0.80 (14).

The *FUT2* region also showed some evidence of association in a recent meta-analysis of type 1 diabetes GWA studies (15) ([www.tlbase.org](http://www.tlbase.org)). The most associated SNP was rs485186 T>C (multiplicative allelic effects  $P = 1.3 \times 10^{-5}$ , and genotype effects  $P = 3.3 \times 10^{-8}$ ), which is in linkage disequilibrium with rs601338 ( $r^2$  of 0.85 in 60 unrelated CEU parents [[www.1000genomes.org](http://www.1000genomes.org)]). Because the

*FUT2* SNP rs485186 T>C association was based on partially imputed genotypes and did not meet the criteria for follow-up in the type 1 diabetes GWA meta-analysis study (15) (multiplicative allelic effects  $P < 1 \times 10^{-6}$  [15]), and rs601338A>G (W143X) and provides the nonsecretor allele (A/A) in subjects of white European ancestry, we genotyped rs601338 in 8,344 case subjects, 10,008 control subjects, and 3,360 families (providing 5,182 trios) and tested for an association with type 1 diabetes.

**RESEARCH DESIGN AND METHODS**

Case subjects were diagnosed with type 1 diabetes before 17 years of age (mean age at diagnosis 7.8 years) and were from the Juvenile Diabetes Research Foundation/Wellcome Trust Diabetes and Inflammation Laboratory Genetic Resource Investigating Diabetes study ([www.childhood-diabetes.org.uk/grid.shtml](http://www.childhood-diabetes.org.uk/grid.shtml)). Control subjects were obtained from the British 1958 Birth Cohort ([www.cls.ioe.ac.uk/studies.asp?section=000100020003](http://www.cls.ioe.ac.uk/studies.asp?section=000100020003)) and the Wellcome Trust Case-Control Consortium U.K. Blood Service Common Control sample collection (15,16). All samples were of white European ancestry.

Families consisted of 211 multiplex families from Northern Ireland, 394 multiplex families from the Diabetes U.K. Warren collection, 52 multiplex families from Yorkshire (U.K.), 246 simplex/multiplex families from Romania, 299 multiplex families from the Human Biological Data Interchange (U.S.), 951 simplex/multiplex families from Finland, and 1,207 families made available through the Type 1 Diabetes Genetics Consortium (T1DGC) (<http://www-gene.cimr.cam.ac.uk/todd/dna-refs.shtml>; <http://www.t1dgc.org>). The T1DGC families comprised 579 families from across Europe, 481 from North America, 52 from the U.K., and 95 from the Asian-Pacific region.

**Genotyping.** A total of 8,344 type 1 diabetic subjects, 10,008 control subjects, and 3,360 type 1 diabetic families were genotyped at the *FUT2* SNP rs601338 and at two HLA tag SNPs rs2187668 and rs660895 using TaqMan 5' nuclease assay (Applied Biosystems, Warrington, U.K.) according to the manufacturer's protocol. Genotyping was performed blind to case-control status and double scored to minimize error. Over 3,000 samples from the family collections were double genotyped, and no discordances were found.

It has been reported that there is a SNP (rs1800459) adjacent to rs601338 (8), which could be problematic for TaqMan primer or probe binding efficiency. We resequenced the flanking region of rs601138/rs1800459 in 768 samples, a mix of case and control subjects, and found rs1800459 to be monomorphic. No additional SNPs were found in the probe or primer sequences, therefore, we do not agree with Hazra et al. (8). The sequenced genotypes for rs601138 were concordant with the TaqMan genotypes.

Primers and probes were as follows: forward primer GGGAGTACGTCC-GCTTCAC; reverse primer TGGCGGAGGTGGTGGTA; FAM probe CTGCTCCT-GGACCTT; and VIC probe CCTGCTCCTAGACCTT. *HLA-DRB1* and *HLA-DQB1* were genotyped at four digit resolution using Dynal RELI SSO assays (Invitrogen Ltd, Paisley, U.K.).

**Statistical analysis.** All statistical analyses were performed in either Stata ([www.stata.com](http://www.stata.com)) or R ([www.r-project.org](http://www.r-project.org)). The genotype frequencies of SNP rs601338 in the control subjects conformed to Hardy-Weinberg equilibrium (HWE;  $P = 0.62$ ). In contrast, we found deviation in the case subjects ( $P = 1.1 \times 10^{-5}$ ), affected offspring ( $P = 2.2 \times 10^{-4}$ ), and the unaffected parents ( $P = 0.0033$ ), all adjusted for population. Disease association can result in deviation from HWE in case subjects, affected offspring, and parents of affected offspring, who are not representative of the general population. The deviation is due to an excess of A/A homozygotes in the case subjects. In Crohn's disease case subjects, rs601338 genotype frequencies also deviated from HWE (13). Nevertheless, we performed a number of genotyping checks, including inspection of allele signal intensity plots, resequencing 3,000 samples from the family collections and resequencing the flanking regions of rs601338 to check for unknown polymorphisms. No inconsistencies were detected.

The case-control collection was analyzed using a logistic regression model, adjusted for 12 geographical regions within Great Britain (Southwestern, Southern, Southeastern, London, Eastern, Wales, Midlands, North Midlands, Northwestern, East and West Riding, Northern, and Scotland) to account for possible confounding by geographical variation in genotype frequency and disease incidence. These regions corresponded to the place of collection for case and control subjects. Because the case and control subjects were well matched for region, this stratification resulted in little loss of power (17). The family collection was analyzed using conditional logistic regression, after generating pseudo-controls (18).

When testing for an association between type 1 diabetes and rs601338, we performed a 1-degree of freedom (df) likelihood ratio test to determine whether a 1-df multiplicative allelic effects model was an appropriate approximation for the data or whether a 2-df genotype effects model (no specific mode of inheritance assumed) was required. We obtained evidence for nonmultiplicative allelic effects in both the case-control and family collections ( $P = 2.8 \times 10^{-3}$  and 0.025, respectively, for rejection of the multiplicative allelic effects model). The case-control and family genotype risks indicated a recessive predisposing effect (see genotype risks in Table 1), because two copies of the A allele (the nonsecretor allele) were required to confer susceptibility to type 1 diabetes. Consequently, we then performed an additional 1-df likelihood ratio test to determine whether a 1-df recessive effects model or a 2-df genotype effects model was more appropriate for the data. The scores and their variances were summed to pool case-control and family information. Importantly, there was no evidence of heterogeneity in the disease associations across different populations, regions or countries, despite there being significant frequency differences (Supplementary Tables 1–4).

A case subject-only (affected offspring only) test for age-at-diagnosis effects was performed by regressing age at diagnosis on SNP genotype (coded 0 or 1 for counts of the A/A genotype) on more than 13,500 case subjects. We adjusted for geographical region when analyzing case subjects from the case-control collections, and for collection when analyzing affected offspring from the family collections. We tested for sex effects in a similar manner using a logistic regression model with sex as the outcome variable.

To maximize power, interaction between HLA class II and *FUT2* was only tested in the case subjects and affected family members, which assumes

**TABLE 1**  
Association results for rs601338G>A (W143X) in the type 1 diabetes case-control and family collections

	Case subjects	Control subjects	OR (95% CI)	P values
Genotype model				2.94x10 <sup>-13</sup>
GG	1,849 (22.2)	2,435 (24.3)	1.00 (ref)	
AG	3,943 (47.3)	4,978 (49.7)	1.05 (0.97–1.45)	
AA	2,552 (30.6)	2,595 (25.9)	1.33 (1.29–1.73)	
Recessive model				7.28x10 <sup>-14</sup>
GG,GA	5,792 (69.4)	7,413 (74.1)	1.00 (ref)	
AA	2,552 (30.6)	2,595 (25.9)	1.29 (1.20–1.37)	
	Affected offspring	Parents	RR (95% CI)	
Genotype model				2.72x10 <sup>-5</sup>
GG	1,414 (27.4)	1,989 (29.6)	1.00 (ref)	
AG	2,447 (47.5)	3,165 (47.1)	1.04 (0.95–1.13)	
AA	1,295 (25.1)	1,566 (23.3)	1.26 (1.12–1.40)	
Recessive model				6.81x10 <sup>-6</sup>
GG,GA	3,861 (74.9)	5,154 (76.7)	1.00 (ref)	
AA	1,295 (25.1)	1,566 (23.3)	1.22 (1.12–1.32)	

Data are n (%) unless otherwise indicated. CI, confidence interval; OR, odds ratio; ref, reference genotype; RR, relative risk. The recessive model was found to be an appropriate approximation in both datasets (case-control  $P = 0.19$  and family  $P = 0.35$ , for rejection of the recessive effects model).

conditional independence in the control population (19). Under the null, the loci are assumed to interact multiplicatively, while under the alternative they interact nonmultiplicatively. Due to the number of genotypes present at *HLA-DRB1* and *HLA-DQB1*, the coding of the class II locus had to be simplified. Therefore, two SNPs, rs2187668 and rs660895, were used to tag the HLA-DRB1\*03 (DR3) and HLA-DRB1\*04 (DR4) class II alleles in up to 15,841 type 1 diabetic subjects and affecteds (20). The SNPs were used to code the HLA-DRB1\*03/04 genotype versus the remainder. The HLA-DRB1\*03/04 genotype was confined to those that were HLA-DQB1\*0302 positive, with the protective HLA-DRB1\*0403 and \*0407 included in the non-DR3/DR4 group where classical genotypes were available (12,476 subjects). The second coding adopted used six genotypes: HLA-DRB1\*03/03, HLA-DRB1\*03/04, HLA-DRB1\*04/04, HLA-DRB1\*04/X, HLA-DRB1\*03/X, HLA-DRB1\*X/X, where X represents any non-HLA-DRB1\*03 or non-HLA-DRB1\*04 allele, and included HLA-DRB1\*0403, HLA-DRB1\*0407, HLA-DQB1\*0301 where classical genotypes were available. Interaction was tested using multinomial logistic regression, with *FUT2* genotype as outcome variable and HLA genotype as predictors. Robust variance estimates were used to account for dependence between sibs, and geographical region or collection were included as strata.

The pseudo-control-case methods proposed by Cordell et al. (18) were used to test for parent-of-origin effects. Maternal and offspring genotypes were coded assuming a recessive model, and interaction of maternal genotype and offspring genotype tested in a conditional logistic regression model using a likelihood ratio test on 1-df.

## RESULTS

The wild-type/functional allele (secretor allele) of rs601338A>G is the G allele, which has a frequency of 49% in the control samples and 48% in the European CEU HapMap samples (hapmap.ncbi.nlm.nih.gov; Supplementary Table 1). In the analysis of rs601338A>G we used the G allele as reference, because the nonsecretor allele (A) confers susceptibility to Crohn's disease and homozygotes for the nonsecretor allele are resistant to the Norovirus. We found consistent and convincing evidence of a recessive association between type 1 diabetes and rs601338 in both collections (RESEARCH DESIGN AND METHODS; Table 1). In the case-control collection, the odds ratio for A/A against A/G and G/G was 1.29 (95% CI 1.20–1.37;  $P = 7.3 \times 10^{-14}$ ) and in the family collection, the relative risk for A/A against A/G and G/G was 1.22 (95% CI 1.12–1.32;  $P = 6.8 \times 10^{-6}$ ). The combined evidence for the case-control and family collections provided very convincing evidence for a type 1 diabetes locus at *FUT2* ( $P = 4.3 \times 10^{-18}$ ).

We found unconvincing evidence of age-at-diagnosis effect ( $P = 0.031$ ) and no evidence of a sex effect ( $P = 0.71$ ) in over 13,500 case subjects and affected offspring. In addition, there was no evidence of interaction between *FUT2* and the primary type 1 diabetes susceptibility locus, HLA class II ( $P > 0.60$ ; RESEARCH DESIGN AND METHODS). Given that the A/A genotype protects from Norovirus infection and confers susceptibility to type 1 diabetes, offspring of homozygous nonsecretor mothers who would not receive maternal antibodies to GGII Noroviruses, could have altered risk of type 1 diabetes distinct from the direct effect attributable to the *FUT2* genotype of the offspring. Furthermore, maternal-fetal ABO incompatibility has also been shown to be a risk factor for type 1 diabetes (21). Consequently, *FUT2* could also be associated with perinatal routes of infection or maternal-fetal immune dysregulation in the pathophysiology of type 1 diabetes. However, no evidence of a maternal-fetal interaction was obtained ( $P = 0.49$ ; RESEARCH DESIGN AND METHODS).

To test whether rs601338 was the sole causal variant in the 19q13.33 region, which includes the fucosyltransferase gene (*FUT1*; also involved in the synthesis of the A and B blood group antigens), we combined the rs601338 genotype data with the GWA data from Barrett et al. (15) (116 SNPs) and conducted a stepwise regression analysis in 3,419 case

subjects and 3,524 control subjects with complete genotype data (22). We found some unconvincing evidence for independent effects in the 19q13.33 region (data not shown). However, further analysis will be required on a dense SNP map of the 19q13.33 region in a much larger number of samples to map this region.

## DISCUSSION

We found convincing evidence that having two copies of the nonsecretor allele (A) at rs601338 confers susceptibility to type 1 diabetes. This is only the fourth type 1 diabetes locus with a recessive-like effect, the others being *INS/11p15.5*, *CCR5/3p21* (23), and *BACH2/6q15* (J.D.C., J.M.M.H., D.J.S., J.A.T., unpublished data). Previously, two copies of the nonsecretor allele at rs601338 were shown to provide resistance to some strains of Norovirus. Conversely, having a single functional copy of the *FUT2* gene and the subsequent Norovirus and/or some other infection (10–12) has a protective effect on type 1 diabetes.

It has previously been noted that the allele frequency of rs601338  $se^{428}$  and other *FUT2* polymorphisms vary between populations (Supplementary Table 1) (24), and this was also observed in our study (Supplementary Tables 2–4). The rs601338  $se^{428}$  nonfunctional allele (A) has increased in frequency in the European and Sub-Saharan African populations (Supplementary Table 1) and is more common than the ancestral allele (G), suggesting that the mutant A allele has been under selective pressure (24). Previous studies have failed to detect significant signatures of positive selection for *FUT2* (2,24), which can be explained by the possibility that the A allele is not a recent mutation in human populations allowing sufficient time for recombination to break down haplotypes, and the possibility of convergent evolution in different populations (24). The interpopulation differentiation statistic,  $F_{ST}$ , for rs601338 is 0.26 and, therefore, does provide some evidence supporting balancing selection as well as the existence of different evolutionary forces acting in different geographical areas (24).

Having nonfunctional alleles of *FUT2* protects against Norovirus and slows the progression of HIV (9), which are both beneficial effects, and is also associated with other bacterial and fungal infections, including reduced susceptibility to *H. pylori* infection (<http://geneticassociationdb.nih.gov/cgi-bin/tableview.cgi?table=allview&cond=GENE>). However, these nonfunctional alleles of *FUT2* increase susceptibility to type 1 diabetes and Crohn's disease, although in prewestern civilization, this is not expected to be a selective advantage since these immune diseases would have been very rare. The other possibility is that the mechanistic link between the enzyme *FUT2* and type 1 diabetes has nothing to do with Norovirus or other pathogens and it involves some other target of this  $\alpha(1,2)$  fucosyltransferase. We have recently found that the ABO gene is associated with positivity of autoantibodies to gastric parietal cells in type 1 diabetic patients (25). The *FUT2* enzyme is required for synthesis of the ABO blood group. The ABO SNP rs657152G>T was not associated with risk of type 1 diabetes itself, although there was weak evidence of *FUT2* with parietal cells positivity (25). Hence, *FUT2* could also be associated with susceptibility to autoimmune gastritis, which causes vitamin B12 malabsorption and deficiency, and pernicious anemia (8).

Our results justify follow-up functional investigations and highlight the intriguing relationship between the alleles of host defense genes and those of complex chronic

immune disorders, supporting a model in which alleles selected for diversity and resistance in host immune responses can now, in a modern environment predispose to autoimmune or immune-mediated chronic diseases (1,2). In this case, even one dose of a functional *FUT2* genotype provides about 30% protection against type 1 diabetes across different populations of European origin. It will also be interesting to test if the Asian-specific nonsecretor variant of *FUT2*, I129F (rs1047781) is associated with type 1 diabetes in eastern and south eastern populations.

#### ACKNOWLEDGMENTS

This work was funded by the Juvenile Diabetes Research Foundation International (JDRF), the Wellcome Trust, the National Institute for Health Research Cambridge Biomedical Centre, and the National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK). The Cambridge Institute for Medical Research is in receipt of a Wellcome Trust Strategic Award (079895).

No potential conflicts of interest relevant to this article were reported.

D.J.S. and J.A.T. researched data. D.J.S., J.D.C., and J.M.M.H. conducted analyses. D.J.S., J.D.C., and J.A.T. wrote the manuscript. N.M.W. managed the data. P.C., T.M., and H.S. organized the DNA samples. K.D. contributed to discussion. All authors reviewed, edited, and discussed the manuscript.

The authors thank all the patients, control subjects, and family members for their participation. They thank the U.K. Medical Research Council and Wellcome Trust for funding the collection of DNA for the British 1958 Birth Cohort (MRC grant G0000934, WT grant 068545/Z/02). DNA control samples were prepared and provided by S. Ring, R. Jones, and M. and W. McArdle of the University of Bristol; D. Strachan of the University of London; and P. Burton of the University of Leicester. The authors thank David Dunger and Barry Widmer of the University of Cambridge, and the British Society for Paediatric Endocrinology and Diabetes for the type 1 diabetes case collection. The authors acknowledge use of DNA from the U.K. Blood Services collection of Common Controls, which is funded by the Wellcome Trust grant 076113/C/04/Z, by the Wellcome Trust/JDRF grant 061858, and by the National Institute for Health Research of England. The collection was established as part of the Wellcome Trust Case-Control Consortium. The authors acknowledge use of DNA from the Human Biological Data Interchange and from Diabetes UK for the U.S. and U.K. multiplex families, respectively; D. Savage of the Belfast Health and Social Care Trust, C. Patterson and D. Carson of Queen's University Belfast, and P. Maxwell of Belfast City Hospital for the Northern Irish families; the Genetics of Type 1 Diabetes in Finland, J. Tuomilehto, L. Kinnunen, E. Tuomilehto-Wolf, V. Harjujalsalo, and T. Valle of the National Public Health Institute, Helsinki, for the Finnish families; and C. Guja and C. Ionescu-Tirgoviste of the Institute of Diabetes "N Paulescu," Romania, for the Romanian families. This research uses resources provided by the Type 1 Diabetes Genetics Consortium, a collaborative clinical study sponsored by the NIDDK, the National Institute of Allergy and Infectious Diseases, the National Human Genome Research Institute, the National Institute of Child Health and Human Development, and JDRF and supported by U01 DK062418. The authors also thank H. Stevens, P. Clarke, G. Coleman, S. Duley, D. Harrison,

S. Hawkins, M. Maisuria, T. Mistry, and N. Taylor for preparation of DNA samples.

#### REFERENCES

- Todd JA. Etiology of type 1 diabetes. *Immunity* 2010;32:457–467
- Barreiro LB, Quintana-Murci L. From evolutionary genetics to human immunology: how selection shapes host defence genes. *Nat Rev Genet* 2010; 11:17–30
- Marionneau S, Cailleau-Thomas A, Rocher J, et al. ABH and Lewis histo-blood group antigens, a model for the meaning of oligosaccharide diversity in the face of a changing world. *Biochimie* 2001;83:565–573
- Blackwell CC, Jónsdóttir K, Hanson M, et al. Non-secretion of ABO antigens predisposing to infection by *Neisseria meningitidis* and *Streptococcus pneumoniae*. *Lancet* 1986;2:284–285
- Thom SM, Blackwell CC, MacCallum CJ, et al. Non-secretion of blood group antigens and susceptibility to infection by *Candida* species. *FEMS Microbiol Immunol* 1989;1:401–405
- Blackwell CC, Jonsdottir K, Hanson MF, Weir DM. Non-secretion of ABO blood group antigens predisposing to infection by *Haemophilus influenzae*. *Lancet* 1986;2:687
- Ikehara Y, Nishihara S, Yasutomi H, et al. Polymorphisms of two fucosyltransferase genes (Lewis and Secretor genes) involving type I Lewis antigens are associated with the presence of anti-*Helicobacter pylori* IgG antibody. *Cancer Epidemiol Biomarkers Prev* 2001;10:971–977
- Hazra A, Kraft P, Selhub J, et al. Common variants of *FUT2* are associated with plasma vitamin B12 levels. *Nat Genet* 2008;40:1160–1162
- Kindberg E, Hejdeman B, Bratt G, et al. A nonsense mutation (428G→A) in the fucosyltransferase *FUT2* gene affects the progression of HIV-1 infection. *AIDS* 2006;20:685–689
- Carlsson B, Kindberg E, Buesa J, et al. The G428A nonsense mutation in *FUT2* provides strong but not absolute protection against symptomatic GII.4 Norovirus infection. *PLoS ONE* 2009;4:e5593
- Thorven M, Grahm A, Hedlund KO, et al. A homozygous nonsense mutation (428G→A) in the human secretor (*FUT2*) gene provides resistance to symptomatic norovirus (GGII) infections. *J Virol* 2005;79:15351–15355
- Lindesmith L, Moe C, Marionneau S, et al. Human susceptibility and resistance to Norwalk virus infection. *Nat Med* 2003;9:548–553
- McGovern DP, Jones MR, Taylor KD, et al.; International IBD Genetics Consortium. Fucosyltransferase 2 (*FUT2*) non-secretor status is associated with Crohn's disease. *Hum Mol Genet* 2010;19:3468–3476
- Franke A, McGovern DP, Barrett JC, et al. Genome-wide meta-analysis increases to 71 the number of confirmed Crohn's disease susceptibility loci. *Nat Genet* 2010;42:1118–1125
- Barrett JC, Clayton DG, Concannon P, et al.; Genome-wide association study and meta-analysis find that over 40 loci affect risk of type 1 diabetes. *Nat Genet* 2009;41:703–707
- Wellcome Trust Case Control Consortium. Genome-wide association study of 14,000 cases of seven common diseases and 3,000 shared controls. *Nature* 2007;447:661–678
- Clayton DG, Walker NM, Smyth DJ, et al. Population structure, differential bias and genomic control in a large-scale, case-control association study. *Nat Genet* 2005;37:1243–1246
- Cordell HJ, Barratt BJ, Clayton DG. Case/pseudocontrol analysis in genetic association studies: A unified framework for detection of genotype and haplotype associations, gene-gene and gene-environment interactions, and parent-of-origin effects. *Genet Epidemiol* 2004;26:167–185
- Umbach DM, Weinberg CR. Designing and analysing case-control studies to exploit independence of genotype and exposure. *Stat Med* 1997;16: 1731–1743
- Howson JM, Rosinger S, Smyth DJ, Boehm BO, the ADBW-END Study Group, Todd JA. Genetic analysis of adult-onset autoimmune diabetes. *Diabetes* 2011;60:2645–2653
- Dahlquist GG, Patterson C, Soltesz G. Perinatal risk factors for childhood type 1 diabetes in Europe. The EURODIAB Substudy 2 Study Group. *Diabetes Care* 1999;22:1698–1702
- Lowe CE, Cooper JD, Brusko T, et al. Large-scale genetic fine mapping and genotype-phenotype associations implicate polymorphism in the *IL2RA* region in type 1 diabetes. *Nat Genet* 2007;39:1074–1082
- Smyth DJ, Plagnol V, Walker NM, et al. Shared and distinct genetic variants in type 1 diabetes and celiac disease. *N Engl J Med* 2008;359:2767–2777
- Ferrer-Admetlla A, Sikora M, Laayouni H, et al. A natural history of *FUT2* polymorphism in humans. *Mol Biol Evol* 2009;26:1993–2003
- Plagnol V, Howson JMM, Smyth DJ, et al. Genome-wide association analysis of autoantibody positivity in type 1 diabetes cases. *PLoS Genet*, 2011;7:e1002216