

Barbara Szepietowska,¹ Tamas L. Horvath,² and Robert S. Sherwin¹

Role of Synaptic Plasticity and EphA5-EphrinA5 Interaction Within the Ventromedial Hypothalamus in Response to Recurrent Hypoglycemia

Hypoglycemia stimulates counterregulatory hormone release to restore euglycemia. This protective response is diminished by recurrent hypoglycemia, limiting the benefits of intensive insulin treatment in patients with diabetes. We previously reported that EphA5 receptor-ephrinA5 interactions within the ventromedial hypothalamus (VMH) influence counterregulatory hormone responses during acute hypoglycemia in nondiabetic rats. In this study, we examined whether recurrent hypoglycemia alters the capacity of the ephrinA5 ligand to activate VMH EphA5 receptors, and if so, whether these changes could contribute to pathogenesis of defective glucose counterregulation in response to a standard hypoglycemic stimulus. The expression of ephrinA5, but not EphA5 receptors within the VMH, was reduced by antecedent recurrent hypoglycemia. In addition, the number of synaptic connections was increased and astroglial synaptic coverage was reduced. Activation of VMH EphA5 receptors via targeted microinjection of ephrinA5-Fc before a hyperinsulinemic hypoglycemic clamp study caused a reduction in the glucose infusion rate in nondiabetic rats exposed to recurrent hypoglycemia. The increase in the counterregulatory response to insulin-induced

hypoglycemia was associated with a 150% increase in glucagon release ($P < 0.001$). These data suggest that changes in ephrinA5/EphA5 interactions and synaptic plasticity within the VMH, a key glucose-sensing region in the brain, may contribute to the impairment in glucagon secretion and counterregulatory responses caused by recurrent hypoglycemia.

Diabetes 2014;63:1140–1147 | DOI: 10.2337/db13-1259

Frequent episodes of acute hypoglycemia represent the principal obstacle to achieving optimal glycemic control during insulin treatment in patients with type 1 diabetes and long-standing type 2 diabetes (1,2). This problem is further magnified by the loss of an appropriate counterregulatory response to hypoglycemia that results as a consequence of frequent episodes iatrogenic insulin-induced hypoglycemia (3,4). The molecular mechanisms underlying this phenomenon remain uncertain but are likely to involve the key brain glucose-sensing region, the ventromedial hypothalamus (VMH) (5,6).

We have previously reported that local stimulation of VMH EphA5 receptors by microinjection of ephrinA5-Fc or ephrinA5 overexpression increased, whereas knock-down of VMH ephrinA5 reduced, counterregulatory

¹Yale University School of Medicine, Department of Internal Medicine and Endocrinology, New Haven, CT

²Program in Integrative Cell Signaling and Neurobiology of Metabolism, Section of Comparative Medicine, Yale University School of Medicine, New Haven, CT

Corresponding author: Robert S. Sherwin, robert.sherwin@yale.edu.

Received 15 August 2013 and accepted 6 November 2013.

© 2014 by the American Diabetes Association. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

responses to hypoglycemia. Furthermore, overexpression of VMH ephrinA5 transiently increased local glutamate concentrations, whereas ephrinA5 knockdown produced profound suppression of VMH interstitial fluid glutamine concentrations in the basal state and during hypoglycemia. These data suggest that the activation of VMH EphA5 receptors by ephrinA5 may act in concert with β -cell Eph receptor forward signaling to restore glucose homeostasis during acute hypoglycemia via alterations in glutamate/glutamine cycling (7,8).

Within the central nervous system, Eph receptors and their ligands, the ephrins, play a key role in cell-to-cell communication as well as in synaptic structure and function. Eph receptors function as transmembrane receptor tyrosine kinases and are divided by sequence similarity and ligand affinity into an A and a B subclass. Their ligands, the ephrins, are also divided into an A and a B subclass: the A subclass is tethered to the cell membrane by a glycosylphosphatidylinositol anchor, and members of the B subclass have a transmembrane domain and as a short cytoplasmic region. For the most part, A-type receptors bind to most or all A-type ligands, and B-type receptors bind to most or all B-type ligands (9). As for many other receptor tyrosine kinases, ligand-binding induces “forward signaling,” mostly through phosphotyrosine-mediated pathways. However, ephrins can also signal into their host cell—referred to as “reverse signaling” (10). Eph receptors and their ephrin ligands are present in the adult brain and are specifically enriched in glutamate excitatory synapses (11). Moreover, Eph receptor tyrosine kinases are mainly expressed in synaptic terminals where they influence synaptic plasticity via binding to ephrins found on astrocytic processes that surround the synapse or on neuronal synapses (12,13).

Several observations suggest that changes in hypothalamic synaptic plasticity may play a significant role in the regulation of energy balance. For example, peripheral signals, such as leptin, ghrelin, and estrogen, induce synaptic adaptations that serve as dynamic regulators of neuronal activity in the arcuate nucleus, the hypothalamic center for feeding control (14,15). Whether hypoglycemia per se induces local changes in the VMH affecting both neuronal synapses and/or surrounding glia cells is unknown, but such alterations could potentially modulate neurotransmission within VMH, thereby altering brain glucose sensing. This study tests whether recurrent hypoglycemia alters EphA5 receptor–ephrinA5 interactions within the VMH, which might contribute to diminished activation of counterregulatory responses to acute hypoglycemia.

RESEARCH DESIGN AND METHODS

Animals

Male Sprague-Dawley rats (Charles River, Wilmington, MA) weighing 300–350 g were individually housed in the Yale Animal Resource Center in temperature- (22–23°C)

and humidity-controlled rooms. Animals were fed rat chow (Prolab 3000; Agway, Syracuse, NY) and water ad libitum and were acclimatized to a 12-h light-dark cycle. The Yale University Institutional Animal Care and Use Committee approved the experimental protocols. Different sets of animals were used for the clamp and electron microscopy (EM) experiments described below.

Vascular and Stereotaxic Surgery

Animals were anesthetized ~7–10 days before study and underwent aseptic surgery in which vascular catheters were implanted into the left carotid artery for blood sampling and into the right jugular vein for infusion, as previously described (8). These catheters were tunneled subcutaneously and exteriorized at the back of the neck between the scapulae. Wound staples were used to close the incision. For stereotaxic surgery, animals were placed into a stereotaxic frame (David Kopf Instruments, Tujunga, CA), and stainless steel guide cannulas (Plastics One, Roanoke, VA) were inserted into the brain and secured in place with screws and dental acrylic (coordinates from Bregma: anteroposterior -2.6 mm, mediolateral ± 0.8 mm, and dorsoventral -8.0) for microinjection.

Recurrent Hypoglycemia Protocol

Rats received intraperitoneal injections of insulin (10 units/kg) for 3 consecutive days. After each injection, food was withheld to allow plasma glucose to fall into the hypoglycemic range (30–50 mg/dL); throughout, animals were monitored with tail vein glucose measurements every 30 min using the AlphaTRAK rodent glucometer (Abbott Animal Health, Chicago, IL) to ensure sustained hypoglycemia and to avoid glucose reduction sufficient to cause seizure activity. At the end of this period, the rats were given free access to food again. Control rats received an injection of 0.9% saline under the same conditions.

Microinjection of EphrinA5-Fc or Control-Fc

On the morning of the study, awake, overnight-fasted rats were connected to infusion pumps ~90 min before the start of the experiment and then left undisturbed to recover from handling stress. After the recovery period, 22-gauge microinjection needles (Plastics One), designed to extend 1 mm beyond the tip of the guide cannula, were inserted bilaterally through the guide cannula into each VMH region. Rats then received a microinjection of recombinant human ephrinA5-Fc or recombinant human IgG1-Fc (control-Fc protein; both R&D Systems, Minneapolis, MN; catalog number: 374-EA-200) in a concentration of 0.3 $\mu\text{g}/\mu\text{L}$ dissolved in artificial extracellular fluid delivered at a rate of (0.1 $\mu\text{L}/\text{min}$) over 60 min (dose: 1.8 μg for each side). After the microinjection, needles were left in place for 30 min before being removed. Immediately thereafter, a hyperinsulinemic hypoglycemic clamp study was performed. These compounds have been previously administered into the central nervous system in *in vivo* studies (16) and in

vitro in brain slices (17), without adverse effects. In addition, ephrinA5-Fc has been shown to specifically bind EphA5 receptors (18).

Hyperinsulinemic-Hypoglycemic Clamp

A primed continuous infusion of $20 \text{ mU} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ insulin (Humulin R; Eli Lilly, Indianapolis, IN) was given, and a variable infusion of 20% dextrose was adjusted at 5- to 10-min intervals based on glucose measurements (Analox Instruments, Lunenburg, MA) designed to maintain plasma glucose at 50 mg/dL from 30 to 90 min (19). Additional blood was drawn at baseline and at 30, 60, and 90 min for measurement of insulin, glucagon, epinephrine, and norepinephrine. Rats were killed at study termination, and the probe position was confirmed histologically.

Hormone and Neurotransmitter Analyses

Plasma catecholamine measurements were analyzed by high-performance liquid chromatography using electrochemical detection, and plasma insulin and glucagon concentrations were determined by radioimmunoassay (Linco, St. Charles, MO).

Immunoblot Analysis

Frozen tissue micropunches from VMH and control regions were homogenized in buffer containing 1% NP40, 150 mmol/L NaCl, 50 mmol/L Tris (pH 7.4), 1 mmol/L Na_3VO_4 , 1 mmol/L phenylmethylsulfonyl fluoride, and protease inhibitor (Roche Diagnostics) using a plastic pestle and ultrasonicator. Protein content was assessed with the Bradford protein assay. Protein samples were fractionated under reducing conditions on SDS-9% PAGE (Bio-Rad). After electrophoresis, proteins were electroblotted onto nitrocellulose membranes, blocked with 5% nonfat dry milk in PBS, probed with first antibody (α -tubulin; Cell Signaling, cat. 2125S; ephrin-A5; R&D Systems, cat. AF3743) and EphA5 receptor (EphA5; Sigma-Aldrich, cat. P8651), and incubated with the appropriate secondary antibody conjugated to peroxidase by horseradish peroxidase-linked protein A (1:2000; Sigma-Aldrich). The immunoblots were developed using an enhanced chemiluminescence detection system (Amersham Biosciences).

Electron Microscopy

Briefly, animals were perfused with paraformaldehyde fixative and ultrathin brain sections were cut on a Leica ultramicrotome, collected on Formvar-coated single-slot grids, and analyzed with a Tecnai 12 Biotwin EM (FEI). Glia coverage of the cell membrane of random VMH cells was performed using ImageJ software. EM photographs (original magnification $\times 11,500$) were used to first measure the perimeter of each VMH neuron analyzed, followed by determination of the amount of membrane covered by glia (in nanometers). Results are reported as glia coverage/perimeter of the VMH neurons.

Characteristics of synaptic contacts were defined, as previously described (20). They were collected from serial

sections of the cell membrane of random VMH neurons. Synapse characterization was performed at original magnification $\times 20,000$, and quantitative measurements were performed at original magnification $\times 11,500$. Results are reported as synapses number/perimeter of the VMH neurons (21).

Statistics

Data are expressed as the means \pm SEM. Analysis was performed by one-way ANOVA or the Student *t* test, as appropriate. Statistical analysis was then performed by two-way ANOVA for repeated measures, followed by post hoc analysis using GraphPad Prism 4.0 software (GraphPad Software, Inc., San Diego, CA). $P < 0.05$ was considered statistically significant.

RESULTS

Recurrent Hypoglycemia and EphrinA5 Expression

As shown in Fig. 1, rats exposed to recurrent hypoglycemia for 3 days exhibit a 25% reduction in ephrinA5 expression in the VMH (Fig. 1A and B). In contrast, no significant change in the expression of the EphA5 receptor in the VMH was detected (Fig. 1C and D).

Stimulation of VMH EphA5 Receptors in Rats Exposed to Recurrent Hypoglycemia

To assess the biological consequences of the reduction of ephrinA5 expression, we microinjected the Eph receptor agonist, ephrinA5-Fc, into the VMH before conducting a hyperinsulinemic-hypoglycemic clamp study in rats previously exposed to three episodes of insulin-induced hypoglycemia. Schematic representation of the experimental protocol is presented in Fig. 2A. Body weight and plasma levels of glucose, insulin, glucagon, epinephrine, and norepinephrine were indistinguishable at baseline and immediately after completion of the VMH microinjection of ephrinA5-Fc or control-Fc (Table 1). Subsequently, during the hypoglycemic clamp study, plasma glucose (Fig. 2B) and insulin (Fig. 2C) were indistinguishable between the two groups. EphrinA5-Fc delivery, however, significantly reduced within 15 min the glucose infusion rate required to maintain hypoglycemia (Fig. 2D). This was accompanied by a rapid 150% ($P < 0.001$) increase in glucagon release (Fig. 2E). As was observed in our previous study (8) in rats not exposed to antecedent hypoglycemia, neither plasma epinephrine (Fig. 2F) nor norepinephrine (Fig. 2G) responses to hypoglycemia were significantly altered by VMH delivery of ephrinA5-Fc compared with the control-Fc microinjection.

Effect of Recurrent Hypoglycemia on Glia Ensheathment and Synaptic Input Organization

Next, we assessed if recurrent hypoglycemia affected the VMH synaptic organization and glia ensheathment in rats exposed for 3 days to recurrent hypoglycemia compared with controls, a model we have previously shown suppresses hypoglycemic counterregulation (22). Figure 3A and B compares representative EMs of glia ensheathment




Figure 1—Effect of recurrent hypoglycemia (RH) on ephrinA5 and EphA5 expression. *A*: Representative data showing expression level of ephrinA5 in the VMH in rats after ~3 days of RH as determined by Western blot analysis. Tubulin served as the loading control. *B*: Relative VMH expression of ephrinA5 in control ($n = 8$) and RH ($n = 8$) rats. Data are presented as means \pm SEM. Statistical analysis by Student *t* test: $*P < 0.05$ vs. control. *C*: Representative data showing expression level of the EphA5 in VMH in rats after ~3 days of RH as determined by Western blot analysis. Tubulin served as the loading control. *D*: Relative VMH expression of EphA5 in control ($n = 7$) and RH ($n = 7$) rats. Data are presented as means \pm SEM. Statistical analysis by Student *t* test.

in random VMH neuron perikarya in control and recurrent hypoglycemic rats. The rats exposed to recurrent hypoglycemia exhibited reduced glial coverage of neurons ($P < 0.001$) (Fig. 3C) and, as a result, more total synaptic contacts in random VMH neurons ($P < 0.05$) (Fig. 3D).

DISCUSSION

The current study demonstrates that exposure of nondiabetic rats to recurrent hypoglycemia for 3 consecutive days diminishes ephrinA5 expression within the VMH in association with reduced glial coverage of VMH neurons and synaptic remodeling. In addition, targeted VMH delivery of the EphA5 receptor ligand ephrinA5-Fc enhances glucose counterregulation and glucagon release in rats exposed to recurrent hypoglycemia. These findings are consistent with the possibility that recurrent hypoglycemia diminishes EphA5 receptor forward signaling in the VMH, which in turn reduces the magnitude of glucagon secretion.

Signaling via the EphA/ephrinA receptor system has been reported to regulate neuron-astrocyte interactions that cause rapid changes in synaptic structural and functional plasticity (17,23). It has been proposed that the loss of ephrinA alters astrocytic-neuronal contacts (24), whereas application of ephrinA3-Fc or endogenous ephrin induces rapid growth of the astrocytes processes and growth of new filopodia (17). In addition, activation of EphA4 by ephrinA3 has been shown to induce spine

retraction (25). The current finding that recurrent hypoglycemia decreases ephrinA5 expression and produces diminished glial coverage and more synaptic connections within the VMH raises the question of a possible relationship. The fact that bypassing ephrinA5 using a targeted VMH ephrinA5-Fc microinjection can increase counterregulatory responses in animals exposed to recurrent hypoglycemia is in keeping with the hypothesis that reduced VMH ephrinA5 expression might induce alterations in VMH synaptic plasticity that in turn contribute to the development of disordered glucose counterregulation. However, a direct link between the observed changes in ephrinA5 and in glial coverage as well as synaptic connection rearrangements remains to be established in future studies.

Previous studies have reported rapid changes in synaptic network connectivity and glia morphology in the VMH in response to alterations in energy substrate bioavailability (15,26–30). Acute hypoglycemia has been shown to alter synaptophysin expression, findings consistent with a rapid alternation in synaptic morphology (31,32), whereas insulin-deficient diabetic rats display a decrease in the number of hypothalamic astrocytes as a consequence of increased death and decreased proliferation (33). Given that diabetes and recurrent glucose deprivation are both accompanied by impaired counterregulation (22), these observations are consistent with the hypothesis that synaptic connectivity and the




Figure 2—Effect of acute stimulation of VMH EphA5 receptors with ephrinA5-Fc on glucose counterregulation in rats exposed to recurrent hypoglycemia. The components are a schematic representation of the experimental protocol (A); plasma glucose (B); mean plasma insulin from 30, 60, 90 min time points (C); glucose infusion rate (D); plasma glucagon (E); plasma epinephrine (F); and plasma norepinephrine (G) during the hypoglycemic clamp study ($n = 7$ for ephrinA5-Fc and $n = 9$ for control-Fc). Data are presented as means \pm SEM. Statistical analysis used mixed-model ANOVA with the Bonferroni post hoc test: *** $P < 0.001$ vs. control-Fc. i.v., intravenous.

function of glia in the VMH play a significant role in supporting the neurotransmission required for proper counterregulatory responses to acute hypoglycemia.

It is noteworthy that the principal effect of VMH microinjection of the ephrinA5 ligand on hypoglycemia-induced counterregulatory hormone release was on glucagon, whereas recurrent hypoglycemia normally leads to a suppression of glucagon and epinephrine levels in nondiabetic animals. These findings are consistent with previous studies in mice showing that knockdown of VGLUT2 (vesicular glutamate transporter) selectively in SF1 VMH neurons predominately inhibited the secretion of glucagon in response to acute hypoglycemia (34). It should be noted, however, that alterations in EphA receptor/ephrinA signaling appear to influence epinephrine responses as well. Using a targeted gene expression manipulation approach to chronically alter VMH ephrinA5 expression, we observed that overexpression of ephrinA5 in the VMH stimulates, whereas targeted VMH knockdown of ephrinA5 inhibits, epinephrine as well as glucagon responses to acute hypoglycemia. These effects

also appear to be mediated by alterations in glutamate-glutamine cycling (8). In this study, we did not include a control group not subjected to recurrent hypoglycemia, and thus, the extent that acute delivery of the EphA receptor agonist restored the glucagon response to hypoglycemia in rats exposed to recurrent hypoglycemia cannot be determined. However, results from previous studies from our laboratory using a similar hypoglycemic clamp protocol show the effect of the EphA receptor agonist on glucagon responses appears to be at least as great as the suppressive effect of recurrent hypoglycemia on glucagon levels (8).

Given that the EphA5/ephrinA5 system is mainly localized on glutamatergic synapses (12,25), it is intriguing to speculate that the stimulation of glucagon produced by the EphA5 receptor agonist in the current study is most likely mediated by the augmented VMH glutamatergic neurotransmission. The maintenance of VMH glutamate neurotransmission during acute hypoglycemia is supported by the transport of glutamate into astrocytes, resulting in the production of glutamine for delivery to

Table 1—Characteristics for the rats with recurrent hypoglycemia at baseline and after microinjection of ephrinA5-Fc in the VMH

	Control-Fc (n = 9)		EphrinA5-Fc (n = 7)	
	Basal	60 min postinjection	Basal	60 min postinjection
Body weight (g)		305 ± 6.7		308 ± 5.8
Plasma glucose (mmol/L)	6.5 ± 0.2	6.5 ± 0.1	6.8 ± 0.3	6.8 ± 0.2
Insulin (μU/mL)	9.6 ± 1.8	10.1 ± 2.4	11.7 ± 4.1	8.0 ± 1.6
Glucagon (ng/L)	39 ± 1.9	39 ± 4	47 ± 8	49 ± 4
Epinephrine (pg/mL)	107 ± 43	148 ± 87	88 ± 45	80 ± 46
Norepinephrine (pg/mL)	212 ± 57	268 ± 64	215 ± 50	164 ± 41

Data are means ± SEM.

neurons and glutamate–glutamine cycle activation (35). Previous studies have shown that astrocyte synaptic coverage is linked to glutamate clearance and the activation of metabotropic glutamate receptors (36), and this has been proposed to alter synaptic and astroglia organization, and in turn, neurotransmission (37,38). Thus, the reduced glia coverage of the VMH neurons observed in the current study in rats exposed to recurrent hypoglycemia may have produced a deficit in astroglial function to support proper glutamate neurotransmission

during hypoglycemia. Interestingly, this was associated with more neuronal synaptic contacts in the VMH (Fig. 3D), which appeared to be in large part symmetric and thus potentially γ -aminobutyric acid (GABA) inhibitory in nature (21). An increase in the VMH GABA tone has been shown to be an important contributor to the development of impaired glucose counterregulation in response to recurrent hypoglycemia (39).

Taken together, our data demonstrate that recurrent hypoglycemia alters neuron-glia plasticity in VMH nuclei




Figure 3—Effect of recurrent hypoglycemia (RH) on glia ensheathment and synaptic input organization. Representative electron micrographs show random VMH neuron and glia ensheathment (white arrow) in a control rat (A) and in a rat after RH (B). C: Graph showing glia ensheathment in rats after repeated episodes of insulin-induced hypoglycemia and in controls. D: Graph showing the number of total synaptic connections in rats after repeated episodes of insulin-induced hypoglycemia and in controls. All data are expressed as means ± SEM. Student *t* test or one-way ANOVA. **P* < 0.05, ****P* < 0.001.

and diminishes ephrinA5 ligand expression within the VMH. It is thus possible decreased ephrin-induced activation of Eph receptors in VMH glutamate neurons may contribute to the impairment in glucose counterregulation in response to recurrent antecedent hypoglycemia.

Acknowledgments. The authors are grateful to Wanling Zhu, from R.S.S.'s laboratory, for performing animal surgeries; Aida Groszman, Maria Batsu, Codruta Todeasa, Ralph J. Jacob, from the Yale Center for Clinical Investigation Core Laboratory of the Yale School of Medicine, and Jan Czyzyk, from the University of Rochester Medical Center School of Medicine and Dentistry, for assisting with the protein expression experiments; and Erzsebet Borok, from the Yale Program in Integrative Cell Signaling and Neurobiology of Metabolism, for excellent technical support and assistance.

Funding. This work was supported by research grants from National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK) (20495), the Juvenile Diabetes Research Foundation, and the NIDDK-supported Diabetes Endocrinology Research Center.

Duality of Interest. No potential conflicts of interest relevant to this article were reported.

Author Contributions. B.S. designed the study, performed animal surgery and studies, analyzed data, and wrote the manuscript. T.L.H. designed EM experiments. R.S.S. designed the study, reviewed data, and revised the manuscript. B.S. and R.S.S. are the guarantors of this work, and, as such, had full access to all the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

Prior Presentation. This study was presented at the 70th Scientific Sessions of the American Diabetes Association, Orlando, FL, 25–29 June 2010.

References

- Amiel SA, Tamborlane WV, Simonson DC, Sherwin RS. Defective glucose counterregulation after strict glycemic control of insulin-dependent diabetes mellitus. *N Engl J Med* 1987;316:1376–1383
- Cryer PE, Gerich JE. Glucose counterregulation, hypoglycemia, and intensive insulin therapy in diabetes mellitus. *N Engl J Med* 1985;313:232–241
- Dagogo-Jack SE, Craft S, Cryer PE. Hypoglycemia-associated autonomic failure in insulin-dependent diabetes mellitus. Recent antecedent hypoglycemia reduces autonomic responses to, symptoms of, and defense against subsequent hypoglycemia. *J Clin Invest* 1993;91:819–828
- Davis SN, Mann S, Briscoe VJ, Ertl AC, Tate DB. Effects of intensive therapy and antecedent hypoglycemia on counterregulatory responses to hypoglycemia in type 2 diabetes. *Diabetes* 2009;58:701–709
- Borg WP, Sherwin RS, Doring MJ, Borg MA, Shulman GI. Local ventromedial hypothalamus glucopenia triggers counterregulatory hormone release. *Diabetes* 1995;44:180–184
- Levin BE, Dunn-Meynell AA, Routh VH. Brain glucosensing and the K(ATP) channel. *Nat Neurosci* 2001;4:459–460
- Konstantinova I, Nikolova G, Ohara-Imaizumi M, et al. EphA-Ephrin-A-mediated beta cell communication regulates insulin secretion from pancreatic islets. *Cell* 2007;129:359–370
- Szepietowska B, Zhu W, Czyzyk J, Eid T, Sherwin RS. EphA5-EphrinA5 interactions within the ventromedial hypothalamus influence counterregulatory hormone release and local glutamine/glutamate balance during hypoglycemia. *Diabetes* 2013;62:1282–1288
- Klein R. Bidirectional modulation of synaptic functions by Eph/ephrin signaling. *Nat Neurosci* 2009;12:15–20
- Murai KK, Pasquale EB. 'Eph'ective signaling: forward, reverse and crosstalk. *J Cell Sci* 2003;116:2823–2832
- Martone ME, Holash JA, Bayardo A, Pasquale EB, Ellisman MH. Immunolocalization of the receptor tyrosine kinase EphA4 in the adult rat central nervous system. *Brain Res* 1997;771:238–250
- Tremblay ME, Riad M, Bouvier D, et al. Localization of EphA4 in axon terminals and dendritic spines of adult rat hippocampus. *J Comp Neurol* 2007;501:691–702
- Murai KK, Pasquale EB. Eph receptors, ephrins, and synaptic function. *Neuroscientist* 2004;10:304–314
- Diano S, Farr SA, Benoit SC, et al. Ghrelin controls hippocampal spine synapse density and memory performance. *Nat Neurosci* 2006;9:381–388
- Horvath TL. Synaptic plasticity in energy balance regulation. *Obesity (Silver Spring)* 2006;14(Suppl. 5):228S–233S
- Gerlai R, McNamara A. Anesthesia induced retrograde amnesia is ameliorated by ephrinA5-IgG in mice: EphA receptor tyrosine kinases are involved in mammalian memory. *Behav Brain Res* 2000;108:133–143
- Nestor MW, Mok LP, Tulapurkar ME, Thompson SM. Plasticity of neuron-glia interactions mediated by astrocytic EphARs. *J Neurosci* 2007;27:12817–12828
- Noberini R, Rubio de la Torre E, Pasquale EB. Profiling Eph receptor expression in cells and tissues: a targeted mass spectrometry approach. *Cell Adhes Migr* 2012;6:102–112
- Saccà L, Cryer PE, Sherwin RS. Blood glucose regulates the effects of insulin and counterregulatory hormones on glucose production in vivo. *Diabetes* 1979;28:533–536
- Gray EG. Axi-somatic and axo-dendritic synapses of the cerebral cortex: an electron microscope study. *J Anat* 1959;93:420–433
- Pinto S, Roseberry AG, Liu H, et al. Rapid rewiring of arcuate nucleus feeding circuits by leptin. *Science* 2004;304:110–115
- Powell AM, Sherwin RS, Shulman GI. Impaired hormonal responses to hypoglycemia in spontaneously diabetic and recurrently hypoglycemic rats. Reversibility and stimulus specificity of the deficits. *J Clin Invest* 1993;92:2667–2674
- Richter M, Murai KK, Bourgin C, Pak DT, Pasquale EB. The EphA4 receptor regulates neuronal morphology through SPAR-mediated inactivation of Rap GTPases. *J Neurosci* 2007;27:14205–14215
- Nishida H, Okabe S. Direct astrocytic contacts regulate local maturation of dendritic spines. *J Neurosci* 2007;27:331–340
- Murai KK, Nguyen LN, Irie F, Yamaguchi Y, Pasquale EB. Control of hippocampal dendritic spine morphology through ephrin-A3/EphA4 signaling. *Nat Neurosci* 2003;6:153–160
- Horvath TL, Sarman B, García-Cáceres C, et al. Synaptic input organization of the melanocortin system predicts diet-induced hypothalamic reactive gliosis and obesity. *Proc Natl Acad Sci U S A* 2010;107:14875–14880
- Fuente-Martín E, García-Cáceres C, Granado M, et al. Leptin regulates glutamate and glucose transporters in hypothalamic astrocytes. *J Clin Invest* 2012;122:3900–3913
- Eberhart J, Barr J, O'Connell S, et al. Ephrin-A5 exerts positive or inhibitory effects on distinct subsets of EphA4-positive motor neurons. *J Neurosci* 2004;24:1070–1078
- Flanagan-Cato LM, Fluharty SJ, Weinreb EB, LaBelle DR. Food restriction alters neuronal morphology in the hypothalamic ventromedial nucleus of male rats. *Endocrinology* 2008;149:93–99
- LaBelle DR, Cox JM, Dunn-Meynell AA, Levin BE, Flanagan-Cato LM. Genetic and dietary effects on dendrites in the rat hypothalamic ventromedial nucleus. *Physiol Behav* 2009;98:511–516

31. Singh P, Heera PK, Kaur G. Expression of neuronal plasticity markers in hypoglycemia induced brain injury. *Mol Cell Biochem* 2003;247:69–74
32. Al-Noori S, Sanders NM, Taborsky GJ Jr, et al. Recurrent hypoglycemia alters hypothalamic expression of the regulatory proteins FosB and synaptophysin. *Am J Physiol Regul Integr Comp Physiol* 2008;295:R1446–R1454
33. Lechuga-Sancho AM, Arroba AI, Frago LM, et al. Reduction in the number of astrocytes and their projections is associated with increased synaptic protein density in the hypothalamus of poorly controlled diabetic rats. *Endocrinology* 2006;147:5314–5324
34. Tong Q, Ye C, McCrimmon RJ, et al. Synaptic glutamate release by ventromedial hypothalamic neurons is part of the neurocircuitry that prevents hypoglycemia. *Cell Metab* 2007;5:383–393
35. Magistretti PJ. Role of glutamate in neuron-glia metabolic coupling. *Am J Clin Nutr* 2009;90:875S–880S
36. Oliet SH, Piet R, Poulain DA. Control of glutamate clearance and synaptic efficacy by glial coverage of neurons. *Science* 2001;292:923–926
37. Theodosis DT, Poulain DA, Oliet SH. Activity-dependent structural and functional plasticity of astrocyte-neuron interactions. *Physiol Rev* 2008;88:983–1008
38. Oliet SH. Functional consequences of morphological neuroglial changes in the magnocellular nuclei of the hypothalamus. *J Neuroendocrinol* 2002;14:241–246
39. Chan O, Cheng H, Herzog R, et al. Increased GABAergic tone in the ventromedial hypothalamus contributes to suppression of counterregulatory responses after antecedent hypoglycemia. *Diabetes* 2008;57:1363–1370