

**Zinc, Not Insulin, Regulates the Rat Alpha Cell Response to Hypoglycemia *in vivo***

*Received for publication on 16 October 2006 and accepted in revised form on 26 November 2006*

Huarong Zhou<sup>1</sup>, Tao Zhang<sup>1</sup>, Jamie S. Harmon<sup>1</sup>, Joseph Bryan<sup>2</sup>,  
and R. Paul Robertson<sup>1,3</sup>

From the Pacific Northwest Research Institute<sup>1</sup>, Department of Molecular and Cellular Biology, Baylor College of Medicine, Houston, TX<sup>2</sup>, and the Departments of Medicine and Pharmacology at the University of Washington<sup>3</sup>, 720 Broadway, Seattle, WA 98122

## ABSTRACT

The intraislet insulin hypothesis proposes that the decrement in beta cell insulin secretion during hypoglycemia provides an activation signal for alpha cells to release glucagon. A more recent hypothesis proposes that zinc atoms suppress glucagon secretion via their ability to open alpha cell  $K_{ATP}$  channels. Since insulin binds zinc and zinc is co-secreted with insulin, we tested whether decreased zinc delivery to the alpha cell activates glucagon secretion. In streptozotocin-diabetic Wistar rats, we observed that switching off intrapancreatic artery insulin infusions *in vivo* during hypoglycemia greatly improved glucagon secretion (control =  $240 \pm 261$ ; expt'l =  $4346 \pm 1259$ ; n=5, AUC, pg/ml/90 min,  $p < 0.02$ ). Switching off pancreatic artery infusions of zinc chloride during hypoglycemia also improved the glucagon response (control =  $817 \pm 107$ ; expt'l =  $3445 \pm 573$ ; n=6, AUC, pg/ml/90 min,  $p < 0.01$ ). However, switching off zinc-free insulin infusions had no effect. Studies of glucose uptake in muscle and liver cell lines verified that the zinc-free insulin was biologically active. We conclude that zinc atoms, not the insulin molecule itself, provide the switch-off signal from the beta cell to the alpha cell to initiate glucagon secretion during hypoglycemia.

## INTRODUCTION

Glucagon secretion into the hepatic portal circulation is critical for counterregulation of hypoglycemia by virtue of its stimulatory effect on hepatic glycogenolysis, which releases glucose into the systemic circulation to return blood glucose to normal levels. The decrement in insulin release from beta cells has been proposed to signal the alpha cell to release glucagon (1-3). Recently, we demonstrated this phenomenon directly *in vivo* in streptozotocin-induced diabetic rats and *in vitro* using isolated islets from these animals. In the *in vivo* studies an insulin infusion into the pancreatic artery was switched off when the animals were made hypoglycemic by a jugular vein infusion of insulin (4). Glucagon secretion, absent in control diabetic animals, was fully restored by this maneuver. Glucagon release was not initiated by this maneuver if saline rather than insulin was used, or if insulin was infused but not switched off. The glucagon response was not observed during normoglycemic or hyperglycemic conditions; the switch-off signal was effective only in the setting of hypoglycemia. We corroborated these findings with *in vitro* islet perfusion studies (5).

Recent work from several laboratories has demonstrated that zinc atoms are capable of suppressing alpha cell function via activation of rat alpha cell  $K_{ATP}$  channels (6-9). Since zinc atoms are bound by and co-secreted with insulin, we hypothesized that zinc might provide the switch-off signal during hypoglycemia. To evaluate this hypothesis, we performed *in vivo* studies using intrapancreatic artery infusions of zinc-containing insulin, zinc chloride alone, or zinc-free insulin in streptozotocin-induced diabetic Wistar rats. Insulin and zinc chloride provided the switch-off signal to the alpha cell during hypoglycemia, but zinc-free insulin did not. These results lead us to conclude that decreased delivery of zinc, not insulin, provides the signal from the beta cell to the alpha cell to secrete glucagon during hypoglycemia in the rat.

## RESEARCH DESIGN AND METHODS

### Animals

Male Wistar rats (200-250g) were purchased from Charles River Laboratories. Animals were placed in rooms with 12 h light-dark cycle and constant temperature and given free access to food and water. All experiments were approved by the PNRI Institutional Animal Care and Use Committee. Diabetes was induced by intraperitoneal (i.p.) injection of streptozotocin (STZ; Sigma, St. Louis, MO) 120 mg/kg made up in freshly prepared citrate buffer. After STZ injection, blood glucose (BG) was measured daily. Rats were considered diabetic when BG exceeded 350 mg/dl on two sequential measurements. Animals that did not develop diabetes by 1 day were given a second dose of STZ. Insulin pellets (Linshin Canada Inc, Ontario, Canada) delivering approximately 1 U/24h were inserted under the skin. BG levels of insulin-treated diabetic rats were 100-250 mg/dl. Animals that were diabetic for at least 2 weeks were used for the experiments, at which time they weighed between 300-350 g.

### Surgical Procedure

On the day of the intrapancreatic artery infusions, non-fasted animals were anesthetized by inhalation of 2% isoflurane. The right jugular vein was cannulated with PE50 polyethylene tubing (Becton Dickinson company, Spark, MD) filled with heparin (500 U/ml), which was advanced to the superior vena cava. This cannula was used for intravenous insulin infusion and collection of blood samples. The upper abdomen was opened by a 3 cm vertical incision in the midline starting from the xiphoid process. The pancreas typically receives its blood supply from branches of the splenic and superior and inferior pancreaticoduodenal arteries. The hepatic artery was isolated, ligated superiorly, and punctured using a 25G needle. For infusion of insulin a micro-cannula (0.008 mm I.D., Biotime, Berkeley, CA) was inserted into the hepatic artery and advanced retrograde to the origin of the pancreatic artery which was ligated over the cannula. The positioning of the tip of the

cannula is at the take-off of the superior pancreaticoduodenal artery. This position does not interfere with normal blood flow coming from the celiac artery into the pancreatic arterial blood supply, which serves to carry the solutions delivered by infusion through the micro-cannula. After surgery, the opened abdomen was superfused with warm saline and covered with foil to prevent drying.

### Study protocols

After STZ treatment and establishment of hyperglycemia for at least 2 weeks, overnight fasted, anesthetized diabetic animals were first studied to assess whether they had intact glucagon responses to hypoglycemia. Insulin (0.5 U/ml, 50 $\mu$ L/min) was infused into the jugular vein to induce hypoglycemia and samples were collected before and at 0, 30, 60, and 90 min thereafter for blood glucose (BG), C-peptide, and glucagon measurements. Glucose levels were measured within 2 minutes of sample collection. At the end of the experiment, the jugular catheter was removed, the incision was closed, and the animals were allowed 2 weeks for recovery. Once the defective glucagon response in the diabetic animals had been established, the animals were randomly divided into groups to receive intrapancreatic artery infusions of regular insulin (Humulin), zinc chloride, or zinc-free insulin.

Approximately 2 weeks later, the surgical procedure to isolate the pancreaticoduodenal artery described above was carried out and the anesthetized animals were rested for 30 min. Insulin (0.5 U/ml at 50  $\mu$ L/min), or zinc-free insulin (0.5 U/ml at 50  $\mu$ L/min) in the case of the animals receiving zinc-free insulin pancreaticoduodenal artery infusions, was infused into the jugular vein to decrease BG to approximately 100 mg/dl. After a basal blood sample was collected, zinc chloride (4.5 or 30 mM), insulin (1.5 U/ml), or zinc-free insulin (1.5 U/ml) was infused at a rate of 50  $\mu$ L/min into the superior pancreaticoduodenal artery. The zinc concentrations were chosen because 30  $\mu$ M was used in the studies reported by

Franklin *et al* (9) and 4.5  $\mu\text{M}$  is 2 times greater than the  $\text{IC}_{50}$  for activation of alpha cell  $\text{K}_{\text{ATP}}$  channels (9) and equal to the amount of zinc found in 1.5 U/ml of Humulin R (10). 10 min later, another blood sample was collected. When BG fell to <60 mg/dl, the pancreatic artery infusions were switched off and blood was sampled at 0, 15, 30, 45, 60, and 90 min.

### Assays

Plasma glucose was measured immediately using a Roche Accu-check glucometer. Blood samples were collected into heparin-coated ice-chilled tubes containing 1,000 IU/ml trasylol to prevent degradation of glucagon. Plasma levels of C-peptide and glucagon were measured using rat C-peptide and glucagon radioimmunoassay kits (Linco Research, St. Charles, MO).

### Preparation of Zinc-free insulin

Zinc was removed from a commercial insulin preparation (Novalog, NovoNordsk) by passing over a column of Chelex 100 (BioRad Laboratories). The zinc concentration was 19.6  $\mu\text{g}/\text{ml}$  based on the manufacturer's data. We used ~12 mg of Chelex 100 per ml of insulin solution. This is approximately a twenty-fold excess of Chelex 100 resin based on the manufacturer's estimate of 0.4 meq/ml. The resin was washed with 20 volumes of phosphate-buffered saline at 4°C before loading the insulin solution. Biological activity was assayed by intraperitoneal injection into fed mice and produced hypoglycemia greater than or equal to zinc-containing insulin.

### Glucose uptake into L6 cells

Rat L6 myoblasts (American Type Cell Collection, ATCC catalog #CRL-1458) were maintained at subconfluence in DMEM (ATCC catalog #30-2002) supplemented with 10% fetal bovine serum (FBS) and 1X penicillin-streptomycin (Sigma P4333) at 37°C and 5%  $\text{CO}_2$ . Cells were plated in 6-well plates at a density of  $1 \times 10^6$  cells/well. Once the cells reached 90% confluence, the media was switched to DMEM with 2% FBS and 1X penicillin-streptomycin. They were

allowed to differentiate into myotubes for 7 days with media changes every 2<sup>nd</sup> day. After the 7<sup>th</sup> day, the differentiated cells were washed with Medium 199 (M199, Sigma) and then incubated for 3 h in M199 media containing 0.1% bovine serum albumin (BSA). The myotubes were then incubated for 30 min in glucose-free Krebs-Ringer Buffer (KRB) with 0.1% BSA plus 100 nM insulin (Humulin R, Lilly) or 100 nM zinc-free insulin or 102 nM zinc chloride, based on 0.04 mg zinc chloride/100 U of insulin (10). The buffer was then switched to KRB + 0.1% BSA containing insulin or zinc-free insulin or zinc chloride plus 0.5  $\mu\text{Ci}/\text{ml}$  of 2-deoxy-<sup>14</sup>C-D-glucose (300 mCi/mmol, American Radiolabeled Chemicals) for 5, 10, and 20 min. The cells were washed 3x with ice-cold phosphate buffered saline (PBS) and harvested in 400  $\mu\text{l}$  of lysis buffer (0.1% SDS, 0.2 M NaOH). Glucose uptake was determined by liquid scintillation counting. Non-stimulated glucose uptake was also measured in cells cultured with KRB + 0.1% BSA containing 0.5  $\mu\text{Ci}/\text{ml}$  of 2-deoxy-<sup>14</sup>C-D-glucose for each time point and subtracted from the stimulated uptake.

### Glucose uptake into HepG2 cells

Human HepG2 liver cells (ATCC catalog #HB-8065) were maintained in EMEM (ATCC catalog #30-2003) supplemented with 10% FBS and 1X penicillin-streptomycin at 37°C and 5%  $\text{CO}_2$ . Cells were plated in 6-well plates at a density of  $0.6 \times 10^6$  cells/well. After 48 h, the cells were washed with M199 and then incubated for 3 h in M199 media containing 0.1% BSA. Following a 15 min glucose washout period in glucose-free KRB + 0.1% BSA, the cells were incubated in KRB with 0.1% BSA plus 1  $\mu\text{M}$  insulin (Humulin R, Lilly) or 1  $\mu\text{M}$  zinc-free insulin or 1.02  $\mu\text{M}$  zinc chloride. The buffer was then switched to KRB + 0.1% BSA containing insulin or zinc-free insulin or zinc chloride plus 0.5  $\mu\text{Ci}/\text{ml}$  of 2-deoxy-<sup>14</sup>C-D-glucose plus 500  $\mu\text{M}$  cold deoxy-glucose for 5, 10, and 20 min. The cells were washed 3x with ice-cold PBS and harvested in 400  $\mu\text{l}$  of lysis buffer. Glucose uptake was then determined by liquid

scintillation counting. Non-stimulated glucose uptake was also measured in cells cultured with KRB + 0.1% BSA containing 0.5  $\mu\text{Ci/ml}$  of 2-deoxy- $^{14}\text{C}$ -D-glucose plus 500  $\mu\text{M}$  cold 2-deoxyglucose for each time point and subtracted from the stimulated uptake.

### Statistics

Data are presented as mean  $\pm$  SE. Results were analyzed using Students t-test, Wilcoxon matched pair signed-rank test, or ANOVA, as appropriate. A p value  $<0.05$  was considered statistically significant.

### RESULTS

Prior to abdominal surgery, the group of animals rendered diabetic by injection of streptozotocin had basal glucagon levels of  $39 \pm 5$  pg/ml and small but significant glucagon responses to insulin-induced hypoglycemia (area under the curve during hypoglycemia =  $549 \pm 145$ ; mean  $\pm$  SE, pg/ml/90 min,  $n=20$ ,  $p<0.001$ ).

On the day of the experiment two weeks later, fasted animals underwent anesthesia, laparotomy, and hepatic artery cannulation for infusion of insulin, zinc-free insulin, or zinc chloride. Hypoglycemia was induced by infusion of insulin or zinc-free insulin (for those animals receiving intrahepatic artery infusions of zinc-free insulin) into the jugular vein. Insulin and zinc-free insulin induced hypoglycemia, and switching off the pancreatic arterial infusions of insulin, zinc or zinc-free insulin had no effect on C-peptide levels, which were barely measurable (Fig. 1). Switching off the pancreatic arterial infusion of regular insulin in hypoglycemic animals produced a marked improvement in glucagon secretion compared to the control experiment performed 2 weeks earlier (control:  $240 \pm 261$ ; after switch off:  $4346 \pm 1259$ ,  $n=5$ , AUC, pg/ml/90 min,  $p<0.02$ ; Fig. 2). Switching off the arterial infusion of zinc chloride (30 or 4.5  $\mu\text{M}$ ) also produced a marked improvement in the glucagon response to hypoglycemia [30  $\mu\text{M}$  control:  $178 \pm 841$ ; after switch off:  $5020 \pm 2466$ ,  $n=3$ , AUC, pg/ml/90 min,  $p<0.05$  (Fig. 3); and 4.5  $\mu\text{M}$  control:  $817 \pm 107$ ; after

switch off:  $3445 \pm 573$ ,  $n=6$ , AUC, pg/ml/90 min,  $p<0.01$  (Fig. 4)]. However, the animals that received pancreatic artery infusions of zinc-free insulin did not have improved glucagon responses to hypoglycemia (control:  $632 \pm 164$ ; after switch off:  $938 \pm 296$ ,  $n=6$ , AUC, pg/ml/90 min,  $p=\text{ns}$ ; Fig 5). The incremental glucagon responses are summarized in Fig. 6.

The zinc-free insulin preparation we used caused hypoglycemia as effectively as zinc-containing insulin when infused into the jugular vein (Fig. 1). To further assess the biological activity of zinc-free insulin, we compared the uptake of [ $^{14}\text{C}$ ]-2-deoxyglucose into L6 myocyte and HepG2 hepatocyte cell lines stimulated by regular zinc-containing insulin, zinc chloride, or zinc-free insulin. We saw no significant difference in glucose uptake between zinc-free insulin and zinc-containing insulin in L6 myocytes; the effect in HepG2 hepatocytes was even greater with zinc-free insulin (Fig. 7). Zinc chloride had a small stimulatory effect on 2-deoxyglucose uptake in L6 myocytes and a larger effect in HepG2 cells.

### DISCUSSION

The intraislet insulin hypothesis proposes that insulin released from the beta cell tonically inhibits glucagon secretion and that the marked attenuation of insulin release during hypoglycemia signals the alpha cell to secrete glucagon (1-3). Although cessation of insulin secretion and initiation of glucagon secretion cannot be documented to occur at precisely the same glucose concentration when measuring systemic blood, finer discrimination of bidirectional hormonal regulation would be more evident in islet periportal blood if it could be measured. Our studies were designed to assess whether zinc atoms bound to and co-secreted with insulin contribute to this switch-off signal. Our *in vivo* study on streptozotocin-treated diabetic Wistar rats showed that during hypoglycemia decrements of both zinc-containing insulin and zinc chloride alone, but not zinc-free insulin, signaled the alpha cell to secrete glucagon. In these studies the control experiments were performed in the

same animals two weeks before the pancreaticoduodenal artery infusions, which allowed us to use a paradigm of paired comparisons for statistical analysis. In previous studies (4) we documented that infusion of saline by this route, followed by switching off the saline infusion, does not correct absent glucagon responses to hypoglycemia in this experimental model. Control studies of glucose uptake in L6 muscle cells and HepG2 liver cells documented the bioactivity of our zinc-free insulin preparation. This indicates that zinc atoms, not insulin molecules, provide the switch-off signal to activate the alpha cell during hypoglycemia. Previous work by others suggests multiple potential downstream effects of zinc that could participate in zinc action. For example, zinc has been reported to modulate many cellular functions including cell signal recognition, second messenger metabolism, protein kinase and phosphatase activities and transcription factor activities (11). Long term exposure of skeletal muscle cells to zinc has been shown to stimulate tyrosine phosphorylation of the insulin receptor and increase glucose uptake (12). We observed only a small increase in zinc-stimulated glucose uptake in the L6 cells, likely due to our short treatment period. Studies of the effect of zinc on hepatoma cells demonstrated an activation of the PI3K/Akt pathway (13). While insulin has been shown to stimulate glucose uptake in HepG2 cells (14,15), to our knowledge, our study is the first to demonstrate zinc-stimulated glucose uptake.

The intraislet insulin hypothesis has been debated, in part because of an early report demonstrating glucagon responses to hypoglycemia in type 1 diabetic subjects (16). However, the diabetic subjects in this study had been diagnosed for only 13-15 months, so it is possible that, although immeasurable in systemic venous blood, residual endogenous insulin secretion was still present in the islet periportal circulation. More recent support for the intraislet hypothesis was provided by a report that glucose-regulated glucagon secretion

requires the presence of insulin receptor expression (17). The hypothesis that zinc might be regulating glucagon secretion has been suggested most prominently by Wollheim's group (7,9) reporting that glucagon secretion is sensitive to suppression by zinc ( $EC_{50} = 2.7 \mu\text{mol/l}$  for  $\text{ZnCl}_2$ ), following earlier reports that the SUR1/ $K_{IR6.2}$ -type  $K_{ATP}$  channels present in islet alpha and beta cells are activated by zinc [6,8]. Zinc activation of  $K_{ATP}$  channels will hyperpolarize alpha cells and thus decrease the influx of cytosolic  $\text{Ca}^{2+}$  necessary for glucagon secretion. Alternatively, it has been suggested by Gopel et al. that closure of  $K_{ATP}$  channels could inactivate  $\text{Na}^+$  current, leading to hyperpolarization and cessation of  $\text{Ca}^{++}$  influx (18). A blunted glucagon response to hypoglycemia in Sur1KO mice lacking neuroendocrine-type  $K_{ATP}$  channels supports this concept (19). A report questioning a regulatory role of zinc for glucagon secretion in mice has recently appeared. This study observed that  $\text{ZnCl}_2$  did not suppress glucagon secretion from mouse islets (20), suggesting a significant species difference (mice vs rats) as one explanation for these negative findings. In this work the authors performed experiments using an alpha cell line to demonstrate decreased glucagon secretion when it was exposed to 10 rather than 0 mM glucose; a blocking effect of Wortmanin on insulin-mediated suppression of glucagon secretion; and the inability of Wortmanin to block glucose-suppression of glucagon secretion.

None of these previous studies examined hypoglycemia as the initiator of glucagon secretion or the role that insulin vs zinc might play as a regulator of glucagon secretion. Since the preparation of zinc-free insulin we used caused systemic hypoglycemia when injected into the jugular vein, and was active in stimulating glucose uptake in two cell lines, zinc is clearly not required for insulin action in these situations. Therefore, we reason that the zinc signaling effect on the alpha cell we observed may be specific to the setting of hypoglycemia<sup>1</sup>. That zinc, independent of insulin, can have insulin-

like effects is well-appreciated (21). Furthermore, other endogenous substances, such as intra-alpha cell L-glutamate directly or via stimulation of  $\gamma$ -aminobutyric acid secretion from beta cells, may also negatively regulate glucagon secretion (22).

In conclusion, we propose that during hypoglycemia the principal signal that initiates glucagon secretion is the detection by alpha cells of a sudden decrease in zinc paralleling the fall in insulin in the islet periportal circulation, and that this drop in concentration of zinc atoms, not insulin molecules themselves, closes alpha cell  $K_{ATP}$  channels, promotes entry of calcium, and stimulates glucagon secretion.

<sup>1</sup>Footnote: One of the Reviewers of this manuscript requested that we speculate what the pancreaticoduodenal artery blood zinc concentration might be during our infusions of zinc chloride. If one assumes blood volume of a rat is 24 cc, that blood zinc concentration

is normally 10  $\mu$ M, that circulation time is 3.3 seconds, that 1% of total blood volume or 0.24 cc passes through the artery every 10 seconds, then approximately 4.8 cc passes through every minute at a concentration of 10  $\mu$ M zinc. Since the zinc chloride infusion rate is 50  $\mu$ L or 0.050 cc/min, using a concentration of 4 mM would provide 1:100 (by volume) enrichment by 4000  $\mu$ M for blood containing 10 $\mu$ M zinc and flowing at 4.8 cc/min. This would result in a delta of 10  $\mu$ M to 40  $\mu$ M or a 4-fold step up in zinc concentration reaching the alpha cell. Of course, none of these calculations take into account the probable influence of % protein binding of zinc in blood, which begs the question re the concentration of free zinc ions. These issues will doubtless be more accurately approached using an in vitro methodology, such as islet perfusion and measurements of bound and unbound zinc in the perfusate.

## References

1. Samols E, Tyler J, Marks V: Glucagon-insulin interrelationships. In Glucagon, Molecular Physiology, Clinical and Therapeutic Implications. LeFebvre PJ, Unger RH, Eds. Elmsford, NY, Pergamon Press, 1972, 151-174.
2. Maruyama H, Hisatomi A, Orci L, Grodsky GM, Unger RH: Insulin within islets is a physiologic glucagon release inhibitor. *J Clin Invest* (1984) 74:2296-2299
3. Raju B, Cryer PE: Loss of the decrement in intraislet insulin plausibly explains loss of the glucagon response to hypoglycemia in insulin-deficient diabetes: documentation of the intraislet insulin hypothesis in humans. *Diabetes* 54:757-764, 2005
4. Zhou H, Tran PO, Yang S, Zhang T, LeRoy E, Oseid E, Robertson RP: Regulation of alpha-cell function by the beta-cell during hypoglycemia in Wistar rats: the "switch-off" hypothesis. *Diabetes* 53:1482-1487, 2004
5. Hope KM, Tran PO, Zhou H, Oseid E, Leroy E, Robertson RP: Regulation of alpha-cell function by the beta-cell in isolated human and rat islets deprived of glucose: the "switch-off" hypothesis. *Diabetes* 53:1488-1495, 2004.
6. Bloc A, Cens T, Cruz H, Dunant Y. Zinc-induced changes in ionic currents of clonal rat pancreatic  $\beta$ -cells : activation of ATP-sensitive  $K^+$  channels. *J. Physiol.* 529:723-734, 2000.
7. Ishihara H, Maechler P, Gjinovci A, Herrera P, Wollheim CB. Islet  $\beta$ -cell secretion determines glucagon release from neighbouring  $\alpha$ -cells. *Nat Cell Biol* 5:330-5, 2003
8. Prost A, Bloc A, Hussy N, Derand R, Vivaudou M. Zinc is both an intracellular and extracellular regulator of  $K_{ATP}$  channel function. *J Physiol.* 559:157-167, 2004
9. Franklin I, Gromada J, Gjinovci A, Theander S, Wollheim CB: Beta-cell secretory products activate alpha-cell ATP-dependent potassium channels to inhibit glucagon release. *Diabetes* 54:1808-1815, 2005
10. Mehanna AS. Insulin and oral antidiabetic agents. *Amer J Pharmaceutical Ed* 2005, 69:1-11, 2005.
11. Beyersmann D and Haase H. Functions of zinc in signaling, proliferation and differentiation of mammalian cells. *BioMetals* 14: 331-341, 2001.
12. Miranda ER and Dey CS. Effect of chromium and zinc on insulin signaling in skeletal muscle cells. *Biol. Trace Elem. Res.* 101 (1): 19-36, 2004.
13. Walter PL, Kampkotter A, Eckers A, Barthel A, Schmoll D, Sies H, and Klotz LO. Modulation of FoxO signaling in human hepatoma cells by exposure to copper or zinc ions. *Arch. Biochem. Biophys.* 454 (2):107-113, 2006.
14. Loizou CL, Ozanne SE, and Hales CN. The effect of insulin on  $\Delta^5$  desaturation in hepG2 human hepatoma cells and L6 rat muscle myoblasts. *Prostaglandins, Leukotrienes and Essential Fatty Acids.* 61(2): 89-95, 1999..
15. Martin KR, Failla ML, and Smith JC, Jr.  $\beta$ -Carotene and lutein protect HepG2 human liver cells against oxidant-induced damage. *J. Nutr.* 126(9): 2098-2106, 1996.
16. Bolli G, De Feo P, Perriello G, De Cosmo S, Compagnucci P, Santeusano F, Brunetti P, Unger RH: Mechanisms of glucagon secretion during insulin-induced hypoglycemia in man. Role of the beta cell and arterial hyperinsulinemia. *J Clin Invest* 73:917-922, 1984
17. Diao J, Asghar Z, Chan CB, Wheeler MB: Glucose-regulated glucagon secretion requires insulin receptor expression in pancreatic alpha-cells. *J Biol Chem* 280:33487-33496, 2005
18. Gopel SO, Kanno T, Barg S, Weng XG, Gromada J, Rorsman P. Regulation of glucagon release in mouse  $\alpha$ -cells by  $K_{ATP}$  channels and inactivation of TTX-sensitive  $Na^+$  channels. *J Physiol.* 528:509-20, 2000.
19. Hussain K, Bryan J, Christesen HT, Brusgaard K, Aguilar-Bryan L: Serum glucagon counterregulatory hormonal response to hypoglycemia is blunted in congenital hyperinsulinism. *Diabetes* 54:2946-2951, 2005

20. Ravier MA, Rutter GA: Glucose or insulin, but not zinc ions, inhibit glucagon secretion from mouse pancreatic alpha-cells. *Diabetes* 54:1789-1797, 2005
21. Sakurai H, Adachi Y: The pharmacology of the insulinomimetic effect of zinc complexes. *Biometals* 18:319-323, 2005
22. Uehara S, Muroyama A, Echigo N, Morimoto R, Otsuka M, Yatsushiro S, Moriyama Y: Metabotropic glutamate receptor type 4 is involved in autoinhibitory cascade for glucagon secretion by alpha-cells of islet of Langerhans. *Diabetes* 53:998-1006, 2004

## FIGURES AND LEGENDS

Figure 1. Levels of glucose and C-peptide before, during, and after switching off intrapancreatic artery infusions of insulin, zinc chloride, or zinc-free insulin during hypoglycemia in streptozotocin-induced diabetic rats. Results from the first blood sample collected during the experiments (Point A) corresponds to the glucose and C-peptide levels just before the jugular vein infusions were begun to induce hypoglycemia. Intrapancreatic artery infusions (represented by the black bar along the horizontal axis) of insulin, zinc chloride, or zinc-free insulin were begun when blood glucose reached 100-150 mg/dl. Point B corresponds to the levels of glucose and C-peptide 10 minutes after the pancreatic artery infusions started. The vertical dashed line at 0 time indicates the levels of glucose and C-peptide at the time that the infusions were switched off. The time elapsed between point A and the switching off of the pancreatic artery infusions varied depending on how much time it took (generally 30-60 minutes) for the animals to reach a glucose level < 60 mg/dl. In all cases glucose nadirs were <60 mg/dl and C-peptide was not measurable.

Figure 2. Levels of glucagon before, during, and after switching off an intrapancreatic artery infusion of regular insulin during hypoglycemia in streptozotocin-induced diabetic rats. See legend for Fig. 1 for infusion protocol. The glucagon response calculated as AUC was significantly greater ( $p < 0.05$ ) than the response in the same animals during insulin-induced hypoglycemia 2 weeks earlier.

Figure 3. Levels of glucagon before, during, and after switching off an intrapancreatic artery zinc chloride (30 $\mu$ M) infusion during hypoglycemia in streptozotocin-induced diabetic rats. See legend for Fig. 1 for infusion protocol. The glucagon response calculated as AUC was significantly greater ( $p < 0.05$ ) than the response in the same animals during insulin-induced hypoglycemia 2 weeks earlier.

Figure 4. Levels of glucagon before, during, and after switching off an intrapancreatic artery zinc chloride (4.5 $\mu$ M) infusion during hypoglycemia in streptozotocin-induced diabetic rats. See legend for Fig. 1 for infusion protocol. The glucagon response calculated as AUC was significantly greater ( $p < 0.02$ ) than the response in the same animals during insulin-induced hypoglycemia 2 weeks earlier.

Figure 5. Levels of glucagon before, during, and after switching off an intrapancreatic artery zinc-free insulin infusion during hypoglycemia in streptozotocin-induced diabetic rats. See legend for Fig. 1 for infusion protocol. The glucagon response calculated as AUC was not significantly different than the response in the same animals during insulin-induced hypoglycemia 2 weeks earlier.

Figure 6. Comparison of the glucagon responses expressed as AUCs in the 4 experimental groups vs the control experiment conducted 2 weeks previously.

Figure 7. Demonstration in studies of glucose uptake by L6 muscle cells (Top Panel) and HepG2 hepatic cell (Bottom Panel) lines that the zinc-free preparation used for infusions was as, if not more, biologically active as zinc-containing regular insulin. 1.0  $\mu$ M concentrations were used for insulin, zinc-free insulin, and zinc chloride. See Methods for details.

**Fig 1**




Fig 2




Fig 3




Fig 4





Fig 5



**Fig 6**




Fig 7




Fig 7

