

The TGF- β pathway is a common target of drugs that prevent experimental diabetic retinopathy

Running title: Drug targets in diabetic retinopathy

Chiara Gerhardinger,¹ Zeina Dagher,¹ Paola Sebastiani,² Yong Seek Park,¹ and Mara Lorenzi¹

¹Schepens Eye Research Institute and Department of Ophthalmology, Harvard Medical School, 20 Staniford Street, Boston, Massachusetts 02114, USA. ²Department of Biostatistics, Boston University School of Public Health, 715 Albany Street, Boston, Massachusetts 02118, USA

Correspondence:

Dr. Chiara Gerhardinger

email: chiara.gerhardinger@schepens.harvard.edu

Additional information for this article can be found in an online appendix at
<http://diabetes.diabetesjournals.org>

Submitted 24 July 2008 and accepted 2 April 2009.

This is an uncopyedited electronic version of an article accepted for publication in *Diabetes*. The American Diabetes Association, publisher of *Diabetes*, is not responsible for any errors or omissions in this version of the manuscript or any version derived from it by third parties. The definitive publisher-authenticated version will be available in a future issue of *Diabetes* in print and online at <http://diabetes.diabetesjournals.org>.

Objective. Prevention of diabetic retinopathy would benefit from availability of drugs that preempt the effects of hyperglycemia on retinal vessels. We aimed to identify candidate drug targets by investigating the molecular effects of drugs that prevent retinal capillary demise in the diabetic rat.

Research Design and Methods. We compared the gene expression profile of retinal vessels isolated from rats with 6 months of streptozotocin-diabetes with that of control rats. We then tested if the aldose reductase inhibitor sorbinil and aspirin, which have different mechanisms of action, prevented common molecular abnormalities induced by diabetes. The Affymetrix Rat Genome 230 2.0 array was complemented by RealTime-RT-PCR, immunoblotting, and immunohistochemistry.

Results. The retinal vessels of diabetic rats showed differential expression of 20 genes of the TGF- β pathway, in addition to genes involved in oxidative stress, inflammation, vascular remodeling, and apoptosis. The complete loop of TGF- β signaling, including Smad2 phosphorylation, was enhanced in the retinal vessels, but not in the neural retina. Sorbinil normalized the expression of 71% of the genes related to oxidative stress, and 62% of those related to inflammation. Aspirin had minimal or no effect on these two categories. The two drugs were instead concordant in reducing the upregulation of genes of the TGF- β pathway (55% sorbinil and 40% aspirin) and apoptosis (74% and 42%, respectively).

Conclusions. Oxidative and inflammatory stress is the distinct signature that the polyol pathway leaves on retinal vessels. TGF- β and apoptosis are however the ultimate targets in order to prevent the capillary demise of diabetic retinopathy.

The clinical manifestations of diabetic retinopathy are consequences of a microangiopathy where retinal capillaries develop thickened basement membranes, lose pericytes and endothelial cells by accelerated apoptosis, form microaneurysms, become abnormally permeable, and eventually are transformed into acellular tubes of basement membrane no longer perfused (1-3). When a critical number of capillaries are affected, retinal edema and/or ischemia ensue. In principle, prevention should be attainable with correction of the hyperglycemia; but is not being attained today because the means of treating diabetes are still imperfect and do not easily maintain normoglycemia. Thus, retinopathy and its sight-threatening features continue to occur even among patients treated intensively (4).

There is an outstanding need for drugs that complement antidiabetic treatment by limiting the effects of residual hyperglycemia on retinal vessels. No such drugs are available for clinical use, despite the fact that several have been tested in human diabetic retinopathy. The clinical trials performed to date may have failed because the drugs were given too late, for too short a period, and/or at ineffective doses (5); but also because the drug targets may have been irrelevant to human diabetic retinopathy. These considerations encourage the development of approaches that would limit the risk of both false negative and true negative trials in the future.

One such approach could be to identify the molecular processes that are causally linked to retinal microangiopathy in animal models, and then test if the processes are active in human diabetic retinopathy and represent therefore efficient targets for new drug trials. We tested this approach using the streptozotocin diabetic rat, which models human diabetic retinopathy more accurately

than the diabetic mouse (6). To capture comprehensively the effects of diabetes on retinal vessels we examined the genome-wide gene expression profile of the vessels, which had not been investigated previously. The results of gene profiling can be translated to human diabetes because changes in gene expression can be studied accurately in postmortem human retinas (7,8), at variance with biochemical and posttranslational changes (8). We studied rats with 6 month-duration of diabetes to capture events reflecting the ongoing injury induced by diabetes as well as the adaptive or maladaptive response to the injury that precedes capillary obliteration. The latter increases in the rat after 8-9 months of diabetes (3,9).

To identify the abnormalities with pathogenic importance, we compared the molecular effects of drugs well documented to prevent the histological features of the microangiopathy in the diabetic rat. We reasoned that pathways that are attenuated by more than one drug and have relevant biological consequences can be attributed a candidate pathogenic role, to be verified and pursued in successive studies. In this work, we studied the aldose reductase inhibitor (ARI) sorbinil and aspirin because they protect the retinal vessels of diabetic rats (10,11) likely through different mechanisms. Sorbinil inhibits the first enzyme of the polyol pathway, which is activated when glucose levels are high, and can damage cells by multiple mechanisms (12). Aspirin is a pleiotropic drug that protects the retinal vessels against diabetes in multiple species (11,13), through mechanisms that do not appear to include its antiplatelet effects (11), but are otherwise unsettled. There was a reasonable expectation that the effects of the two drugs on molecular pathways would show differences as well as similarities; and that the similarities, in turn, could identify

essential pathways to be silenced or attenuated in order to attain prevention of the vascular damage induced by diabetes.

RESEARCH DESIGN AND METHODS

Animals. Procedures involving animals were approved by the Animal Care and Use Committee of the Schepens Eye Research Institute. Sprague-Dawley male rats (five weeks of age, Taconic Farms, Hudson, NY) were randomly assigned to one of the following groups: control, diabetic, diabetic treated with sorbinil (65 mg/Kg/day, gift of Pfizer, Groton, CT), and diabetic treated with aspirin (30 mg/Kg/day). Induction of diabetes with streptozotocin, administration of sorbinil and aspirin with the food, and treatment of diabetic rats with maintenance insulin were as previously described (10,11). Diabetic rats and age-matched non-diabetic control rats were studied after 3, 6, and 8 months of diabetes. HbA1c, plasma salicylates levels, and retinal levels of sorbitol and fructose were measured as described (6,10,11).

Isolation of retinal vessels and RNA extraction. Rat retinal vessels were isolated by hypotonic lysis of the fresh retina, a method documented to yield the retinal vascular network free of glia and neural contamination (10). The two retinas of each rat were dissected and incubated in ice-cold sterile distilled water for 1 hour at 4°C. After incubation with DNase to dissolve DNA released from the lysed neural and glial cells, the vascular network was transferred to ice-cold PBS and cleaned of debris and remaining glial and neural elements by gentle pipetting. Total RNA free of contaminating genomic DNA was isolated using the RNeasy mini kit (Qiagen, Valencia, CA) following the on-column DNase digestion protocol according to the manufacturer instructions.

The retinal vascular network was also prepared by trypsin digestion (3) of retinas fixed in formalin immediately after sacrifice.

Oligonucleotide microarray hybridization and data analysis. Gene expression profiling of rat retinal vessels was performed using the GeneChip[®] Rat Genome 230 2.0 array (Affymetrix, Santa Clara, CA). Each sample subjected to microarray hybridization consisted of the RNA extracted from the vessels of both retinas of one individual rat. Because the amount of RNA was insufficient for the standard hybridization protocol, each RNA sample was subjected to a round of amplification (Ovation Biotin System, NuGEN Technologies, San Carlos, CA). GAGE R&R analysis documented that the amplification procedure did not introduce variance. All procedures for microarray analysis were carried out at the Harvard Medical School - Partners Healthcare Center for Genetics and Genomics Core Laboratory (<http://www.hpcgg.org/>). Gene expression data were preprocessed with MAS 5.0, and probes consistently labeled as absent in all samples were excluded from subsequent analysis.

Analysis of the microarray data was performed using BADGE (Bayesian Analysis of Differential Gene Expression, <http://genomethods.org/badge>), a Bayesian approach to identify differentially expressed genes across two experimental conditions and designed to yield high reproducibility at low sample size (14). The differential expression of each gene in two conditions is estimated by the fold change, and evidence of differential expression is measured by the probability that the fold change exceeds a fixed threshold, conditional on the data. To reduce the number of false positives due to multiple comparisons, the selection of significant changes of expression was based on a threshold on the posterior probability specified to achieve a preset false discovery rate as determined by SAM analysis (15).

To identify the genes whose expression is affected by diabetes, only the 19,054 probe sets "present" in all samples (8

diabetic and 9 controls) were included in the analysis. Probe sets with a probability >0.985 (equivalent to a false discovery rate < 0.01) were selected as differentially expressed in diabetes. To determine which of the diabetes-induced gene expression changes were prevented by the two drugs, we selected the probes identified as differentially expressed in diabetes and compared their expression in the sorbinil-treated diabetic group (n=5) versus the control group, and in the aspirin-treated diabetic group (n=9) versus the control group. Genes for which the posterior probability of differential expression versus control was > 0.95 were considered different from control and the diabetes-induced changes were thus classified as not corrected by the drug. Conversely, when the posterior probability of differential expression versus control was < 0.95 , the expression was classified as similar to control and thus normalized by the drug. The threshold was higher than that used to identify differential expression caused by diabetes in order to increase the specificity of the conclusions about the drug effects.

The probe sets differentially expressed in diabetes were annotated using NetAffx (www.affymetrix.com/analysis/), and the annotations were individually verified based on the latest UniGene cluster of the corresponding representative sequence. The differentially expressed genes identified through the annotation process were assigned to functional categories based on the information provided by Entrez Gene and additional information from the published literature.

Real-time RT-PCR. cDNA was synthesized (8) from one μg of total RNA isolated from retinal microvessels and total retina of rats different from those studied in the microarray experiments. RealTime PCR was performed as previously described (16). Primers and probe sets for rat TGF- β 1 and ADAMTS1 were from Applied Biosystems (Foster City, CA). Relative expression was

determined by the comparative $\Delta\Delta\text{C}_T$ method using β -actin as the endogenous control (16).

Western blotting. Fresh retinal microvessels and whole retinas were homogenized in RIPA buffer containing protease and phosphatase inhibitors (10). Immunoblotting was performed as described (10,16). The blots were sequentially probed for: TGF β 1, COX-1, and NCoA3. RasGAP (17) or β -actin were used to verify protein loading. For evaluation of Smad2 phosphorylation, blots were probed first for phosphorylated Smad2, stripped and reprobed for total Smad2. The primary antibodies used are listed in Supplemental Table 1 (available in the online appendix at <http://diabetes.diabetesjournals.org>).

TGF β 1 immunohistochemistry. Retinal trypsin digests were rehydrated in PBS and incubated in citrate buffer (pH 6.1, Dako, Carpinteria, CA) at 95°C for heat-mediated antigen retrieval. After permeabilization in 1% Triton X-100, quenching of endogenous peroxidase in 3% H_2O_2 , and blocking in 10% normal goat serum, endogenous biotin (unmasked by the antigen retrieval procedure) was blocked by incubation in avidin and biotin blocking solutions (Vector Laboratories, Burlingame, CA). The trypsin digests were then incubated overnight at 4°C with the TGF β 1 primary antibody (V-22, Santa Cruz) diluted to $1\mu\text{g}/\text{ml}$ in 2% goat serum/0.1% Triton X-100 in PBS. The antigen-antibody complexes were detected with the avidin-biotin-peroxidase method (ABC Elite, Vector), and visualized with diaminobenzidine. Each experiment included trypsin digests from each of the 4 groups of rats as well as a negative control obtained by substituting the primary antibody with an equivalent concentration of normal rabbit IgG. The trypsin digests were photographed at 5X (AxioVision Imaging System, Carl Zeiss, Gottingen, Germany) to reconstruct the entire vascular network from each retina (11). The networks were used to

identify for each rat 4 areas in the mid-retina with equivalent location and free of artifacts. The 4 areas were photographed at 20X magnification using the same exposure and light setting to allow for comparison among samples. For evaluation of TGF β RI immunostaining, the images were grouped in panels of 12 images each, with each panel containing images from control, diabetic, and diabetic treated rats. The staining intensity of the capillaries within each image was scored on a continuous scale from 0 (no staining) to 4 (highest staining) by four masked observers. The scores attributed to each retina by the four observers were averaged to obtain one individual final score for each rat. The scoring was consistent among the four observers, as indicated by an overall coefficient of variation of 17%.

Membrane attack complex (MAC) immunohistochemistry. Complement activation in rat retinal vessels was evaluated by MAC immunostaining on rat retinal sections as previously described (10).

Statistical analysis. The results of the effect of sorbinil and aspirin on the diabetes-induced gene-expression changes are summarized as number of genes whose expression was normalized by the drug, within each functional category. The significance of the deviation of this number from the null hypothesis of no gene being normalized was determined by means of a Fisher's exact test (2x2 contingency tables). Results of Western blotting and Real-Time RT-PCR are summarized as means \pm SD; the results in the diabetic and control group were compared by means of two-tailed unpaired t-test. Results of TGF β RI immunostaining are summarized with the mean \pm SD; the results in the four groups of rats were compared with ANOVA followed by Fisher's protected least significant differences test. The StatView 5.0 software (SAS Institute, Cary, NC) was used for all analyses.

RESULTS

The characteristics of the rats used in the microarray and related studies are presented in Table 1.

The microarray study was based on 31 expression profiles of retinal microvessels isolated from diabetic rats (n=8), diabetic rats treated with sorbinil (n=5), diabetic rats treated with aspirin (n=9), and control rats (n=9). The duration of diabetes was six months. When compared to the nondiabetic rats, the retinal vessels of the diabetic rats showed significantly different expression of 127 known genes (131 probe sets) and 229 ESTs or genes with similarity to known genes but not yet fully characterized. The majority of the 127 known genes (105 upregulated and 22 downregulated by diabetes) could be grouped into several functional categories (Table 2).

The TGF- β pathway in diabetic retinal vessels. There was a prominent effect of diabetes on the expression of genes involved in signaling by members of the TGF- β family or affected by such signaling (Figure 1 and Supplemental Table 2). Twenty genes could be attributed to this category, which made the TGF- β pathway the single functional pathway mostly affected by diabetes in retinal vessels. Diabetes increased the expression of the type I receptor activin receptor-like kinase 5 (ALK-5), the widely distributed TGF- β type I receptor that in endothelial cells signals inhibition of migration and proliferation (18). ALK-1, the TGF- β type I receptor that is expressed more selectively in endothelium and induces migration and proliferation (18), was not upregulated by diabetes. There was increased expression of syndecan-2 that is upregulated by TGF- β and, in turn, upregulates the TGF- β type I and II receptors (19); of CD44, also a target of TGF- β that facilitates the activation of TGF- β and signaling by the type I receptor (20); and of Ubc9, the key enzyme of the sumoylation reaction known to enhance TGF-

β signaling (21). The diabetic retinal microvessels showed enhanced expression of connective tissue growth factor (CTGF), the main TGF- β effector in the induction of fibrosis (22), and of other genes prominently regulated by TGF- β such as tissue inhibitor of metalloproteinase 1 (TIMP1), the serine (or cysteine) proteinase inhibitor HSP47, prostaglandin-endoperoxide synthase 1 (COX1), the histone acetyltransferase nuclear receptor co-activator 3 (NCoA3), and thioredoxin-interacting protein (Txnip) (22-25). For some of these molecules, increased expression was confirmed at the protein level (Supplemental Figure 1).

The diabetic retinal vessels showed changes in gene expression pointing to increased signaling also by bone morphogenetic proteins (BMPs) (Figure 1 and Supplemental Table 2). There was increased expression of BMP receptor type 1a, also named ALK-3 and utilized by BMP-2 (26); and of periostin and frizzled homolog 1, both responsive to vascular injury and known targets of BMP-2 (27,28). Facilitated BMP signaling was also suggested by the downregulation of the Hoxc-8 transcription factor, being Hoxc-8 a negative regulator of transcription of Smad-1 and other BMP-specific R-Smads (29). The array included probe sets for BMP-1 through BMP-7; only the BMP-2 transcript was detected in all samples, and its levels were not modified by diabetes.

Of the three TGF- β isoforms present in mammals, the array showed the TGF- β 2 and TGF- β 3 transcripts; the levels of neither were altered by diabetes. The TGF- β 1 mRNA was undetectable in all samples tested. When measured by the more sensitive real time RT-PCR, TGF- β 1 mRNA levels were increased 1.5 fold in the retinal microvessels of diabetic rats ($P=0.04$, Figure 2A). Increased levels of the TGF- β type I receptor ALK5 was confirmed at the protein level (Figure 2B). Because ALK5 signals through Smad2 and

Smad3, we sought evidence of ALK5 activation by measuring Smad2 phosphorylation in the retinal vessels of diabetic and control rats. We found increased Smad2 phosphorylation both after 6-month and 3-month duration of diabetes (Figure 2C). Taken together, the data indicate that the complete loop for TGF- β 1 signaling is enhanced in diabetic retinal vessels.

To learn if the diabetes-induced changes in the TGF- β pathway extended to non-vascular cells of the retina, we measured the levels of the TGF- β 1 transcript, TGF- β type I receptor, and Smad2 phosphorylation in the whole retina. We observed no changes induced by diabetes (Figure 3A-C).

Genes related to oxidative stress, inflammation, and remodeling in diabetic retinal vessels. Diabetes increased the expression of molecules that generate oxygen-derived free radicals, and decreased the expression of antioxidants (Supplemental Table 3). Similar to the aorta of diabetic rats (30), the retinal vessels showed prominent upregulation of Txnip, a negative regulator of the anti-oxidant thioredoxin. Diabetes also increased the expression of genes involved in inflammation and response to injury (Supplemental Table 4). The gene expression program was thus consistent with, and expanded, previous findings pointing to the occurrence of oxidative stress and inflammation in diabetic retinal vessels (9, 31, 32). Genes involved in tissue remodeling through formation of extracellular matrix, regulation of cell adhesion, and organization of the actin cytoskeleton were upregulated in the diabetic vessels (Supplemental Table 5). All changes related to actin were in the direction of increased polymerization and filament stability, which may contribute anti-proliferative effects (33). Additional gene expression changes induced by diabetes pointed to inhibitory effects on cell cycle progression and cell proliferation (Supplemental Table 6).

Pro-apoptotic pathways in diabetic retinal vessels. Multiple mechanisms of apoptosis appeared activated in diabetic vessels, many not suspected previously (Supplemental Table 7). Changes in the expression of six genes of the p53 pathway converged in the pro-apoptotic direction. Another potential mechanism for direct death signals was the upregulation of FAS. TGF- β can induce apoptosis by multiple mechanisms (34), which include upregulation of FAS and activation of JNK (34,35), the expression of which was also increased in the diabetic retinal vessels (Supplemental Table 7). A potential mechanism for apoptosis by neglect was the downregulation of the endothelial differentiation lysophosphatidic acid G-protein-coupled receptor 2 (Edg2), a survival factor for multiple cell types (36).

Concordant and discordant effects of the ARI sorbinil and aspirin on the diabetes-induced changes in gene expression in retinal vessels. The sorbinil and aspirin doses were previously documented to prevent the development of acellular capillaries in diabetic rats (10,11). In this study, the dose of sorbinil over-normalized sorbitol and reduced fructose accumulation by 96% in the retina of diabetic rats (Table 1), thus achieving almost complete inhibition of glucose flux through the polyol pathway. The aspirin dose, known to inhibit by 90% thromboxane B₂ formation during blood clotting in rats, and thus exert antiplatelet effect (11), resulted in serum salicylate levels (Table 1) 10-20-fold lower than those achieved by anti-inflammatory doses of aspirin (reviewed in Ref. 11). Sorbinil normalized the diabetes-induced expression changes in 56%, and aspirin in 32% of the known genes; the figures for the total number of probe sets (known genes and ESTs) were 48% and 47%, respectively.

Both sorbinil and aspirin attenuated significantly and to a similar extent the effects of diabetes on the TGF- β pathway (Figure 4),

preventing the expression changes of 55% and 40% of the genes, respectively. Of note, both drugs prevented the increased expression of TGF- β receptor 1 and CTGF. Likewise, sorbinil and aspirin reduced significantly and to a similar extent the effects of diabetes on pro-apoptotic pathways, with concordant correction of changes in the p53 pathway, and concordant restoration of survival signals. In contrast, sorbinil reduced the changes in gene expression relevant to oxidative stress and inflammation strikingly more than aspirin (Figure 4). Sorbinil led to normalization of expression of 71% of genes related to oxidative stress and 62% of those related to inflammation, whereas the figures for aspirin were 29% and 3%, respectively (effect of sorbinil vs aspirin on inflammation-related genes $P < 0.0001$). The effect of sorbinil and aspirin on the expression of individual genes is reported in Supplemental Tables 2-8.

The effects of sorbinil and aspirin on the diabetes-induced activation of the TGF- β pathway were also tested on trypsin digests prepared from the retinas of rats with 8 months of diabetes. These preparations afforded the opportunity of studying the retinal vascular network fixed immediately after sacrifice and not subjected to hypotonic lysis while fresh. Fig. 5 shows that the diffuse pattern of TGF β R1 immunoreactivity was increased in the retinal capillaries of diabetic rats as compared to control rats ($P=0.05$), and that both aspirin and sorbinil treatment prevented the increase ($P=0.03$ and $P<0.0001$ vs diabetes, respectively), in agreement with the microarray results (see Supplemental Table 2). Of note, while aspirin returned TGF β R1 immunoreactivity to control values ($P=0.62$ vs control), sorbinil suppressed staining intensity below control values ($P=0.015$ vs control). Phosphorylated Smad2 could not be detected in the fixed vascular preparations.

The discordant effect of sorbinil and aspirin on inflammation as shown by the

mRNA data was consistent with the discordant effect on the increased levels of intercellular adhesion molecule 1 in the diabetic rat retina (11), and was confirmed by data on complement activation in retinal vessels. Supplemental Figure 2 shows that aspirin, at variance with sorbinil (10), failed to prevent deposition of MAC in retinal vessels of diabetic rats.

The signature of diabetes on retinal vessels is tissue-specific. We compared the effects of diabetes on gene expression in retinal vessels to those in Müller cells, the principal glia of the retina that shows signs of activation/reactivity in both human (37) and experimental (6) diabetes. We had previously examined the gene expression profile of Müller cells in the same model used in this study (the rat with 6 month-duration of streptozotocin diabetes), and on a comparable platform (GeneChip Rat Genome RG-U34A, Affymetrix) (16). Supplemental Table 9 shows that the functional gene categories affected by diabetes in retinal vessels differed from those affected in Müller cells. Of the genes related to TGF- β that we found affected by diabetes in retinal vessels, 80% were represented in the RG-U34A array but only 6% of those were affected by diabetes in Müller cells. Of the genes related to apoptosis affected by diabetes in retinal vessels, 68% were represented in the RG-U34A array and 0% of those were affected by diabetes in Müller cells. Only the inflammation-related genes showed overexpression in both the vascular and glial cell types (76% of the genes affected by diabetes in the vessels were represented in the RG-U34A, and 55% were upregulated by diabetes in Müller cells).

DISCUSSION

The results of this work indicate that retinal microangiopathy in diabetic rats is the product of several molecular pathways that are interconnected but not of equal pathogenic importance. Attenuation of increased activity

of the TGF- β pathway and apoptosis, but not of oxidative stress and inflammation, appear to be required in order to prevent the microangiopathy. In view of its role in wound healing, extracellular matrix deposition, and fibrosis (22), TGF- β has been over the years an obvious candidate mechanism for the basement membranes thickening and matrix accumulation that are the hallmark of diabetes on blood vessels and vascular structures. However, the studies pointing to a role of excess TGF- β have been performed solely in relation to diabetic nephropathy (38).

Diabetic retinopathy has not been previously linked to excess TGF- β signaling, likely because the link has been more difficult to capture. We found increased TGF- β and TGF- β signaling selectively in microvessels, but not in cells more abundantly represented in the retina such as Müller glial cells (16) or neurons that would make an increase manifest in whole retinal extracts (this work). The few suggestions to date of a contribution of TGF- β to diabetic retinal microangiopathy have come from the observations that the retina of diabetic rats shows the presence of oncofetal fibronectin (39), an isoform typically stimulated by TGF- β during tissue healing and remodeling; and the retina of diabetic patients shows expression of CTGF in pericytes, a shift to the vascular compartment from the microglial location of CTGF in the retina of nondiabetic individuals (40). Our finding of increased Smad-2 phosphorylation indicates that increased TGF- β signaling does in fact occur in retinal vessels of diabetic rats; and the concordant attenuation of such signaling by drugs that protect the vessels from the effects of diabetes through different mechanisms, suggests that the increased signaling contributes to the vascular pathology. It may be argued that the diabetic retinal vessels showed overexpression of inflammation-related genes, while TGF- β is viewed, in general, as anti-inflammatory. In this respect, the increased TGF- β 1 expression

and activity in diabetes could even represent a compensatory response, and as such be protective for the retinal vessels. However, the fact that both sorbinil and aspirin attenuated the TGF- β pathway, but only sorbinil reduced inflammation, indicates that the TGF- β pathway was upregulated independently of the genes related to inflammation. This agrees with findings in Müller cells, which show in diabetic rats changes in the expression of genes related to inflammation but not to TGF- β (16). Fibrosis is not necessarily driven by inflammation, which might explain the general lack of efficacy of anti-inflammatory mediators in the treatment of fibrotic disease (41). In diabetes in particular, TGF- β upregulation could be induced and sustained by high glucose levels, which stimulate TGF- β promoter activity in vascular cells (42).

Comparison of the effects of sorbinil and aspirin led to the identification of the signature of the polyol pathway on retinal vessels. The ARI prevented or attenuated changes in all functional categories of genes, indicating that the polyol pathway mediates most molecular abnormalities induced by hyperglycemia in rat retinal vessels, including TGF- β overexpression. The role of the polyol pathway in mediating glucose-induced increase in TGF- β has been documented in cultured cells (43). Only the ARI, and not aspirin, prevented the pro-oxidant and pro-inflammatory changes in gene expression; consistent with the discordant effects of the two drugs on other indicators of diabetes-induced oxidative stress and inflammation (this work and Ref. 10,11). This finding, combined with the known pro-oxidant effects of the polyol pathway (44), and the fact that reactive oxygen species --induced by hyperglycemia or other stimuli-- beget inflammation (45), points to the combination of oxidative stress and inflammation as the distinct signature of the polyol pathway on retinal vessels.

The effects of aspirin indicated that attenuation of pro-oxidant and pro-inflammatory gene expression is not necessary in order to prevent retinal vascular cell apoptosis and vessel histopathology (11) in diabetes. We cannot exclude that aspirin may have attenuated aspects of inflammation not reflected in the gene expression data, but we note that the concentration of aspirin achieved in our rats (two orders of magnitude lower than those exerting typical anti-inflammatory effects, reviewed in Ref. 11) also failed to attenuate complement deposition in diabetic retinal vessels. However, it remains conceivable that attenuation of oxidative stress and/or inflammation may be sufficient to prevent retinopathy if these processes in diabetes are upstream of, and contribute to, apoptosis. What the effect of aspirin highlights is that in diseases resulting from the contribution of multiple pathogenic events, the attenuation of some of the events is sufficient to prevent the ultimate phenotype despite persistence of the other known contributors. An example of such interplay was recently documented in a model of Alzheimer's disease (46). The mechanism(s) whereby aspirin attenuated the effects of diabetes on the TGF- β and pro-apoptotic pathways in retinal vessels remain speculative. The dose effective in our diabetic rats was 20-fold lower than the dose reported to decrease the excess TGF- β and CTGF expression induced by diabetes in the rat kidney and by high glucose in cultured mesangial cells (47). Comparison with the selective anti-platelet agent clopidogrel had suggested that our dose of aspirin protects retinal vessels in diabetes by mechanisms other than anti-platelet activity (11). Aspirin could have worked by inhibiting the activity of COX-1 in vascular cells, which expressed higher levels of the enzyme in diabetes. However, it is not known whether increased COX-1 is harmful to vascular cells, and the fact that aspirin prevented the overexpression

of COX-1 (not a previously known effect of aspirin), suggests that the effects on COX-1 were downstream of some other beneficial effect. At concentrations consistent with those achieved in our experiments, aspirin induces NO release from endothelial cells (48).

Our findings have several translational implications. Knowledge that the combination of oxidative stress and inflammation is the distinct signature of the polyol pathway on retinal vessels makes possible to seek such signature in human diabetes to help determine whether there is a rationale for adjunct therapy with ARIs. The action of aspirin, that prevents the vascular histopathology resulting from a complex process by attenuating only selected pathways, establishes a valuable paradigm for development of diversified drug approaches to the complications of diabetes. The observation that increased TGF- β signaling begins early in diabetic retinal vessels, proposes that capillary remodeling also begins early. Early onset of pathological remodeling can be one of the reasons for the “memory” that retinal vessels carry of exposure to the diabetic milieu (5, 49), which translates into poor reversibility of even the initial lesions of diabetic retinopathy and justifies emphasis on prevention (5). Targeting the TGF- β pathway may become a rational strategy in the prevention of the two main microvascular complications of diabetes, retinopathy and nephropathy. In view of the homeostatic importance of TGF- β , the targeting modalities would need not to interfere with physiological activity. Partial attenuation of overactivity appears to be sufficient to limit the consequences on tissues. Phenotypes due to increased TGF- β signaling can be rescued by inactivation of a single

TGF- β 1 allele (50), and in our study prevention of retinal diabetic microangiopathy did not require complete normalization of the gene expression changes in the TGF- β pathway. Angiotensin II blockers may prove effective anti-TGF- β drugs, as they were recently reported to reduce the progression of vascular pathologies caused by excessive TGF- β signaling (51). This effect of angiotensin II blockers, coupled with our findings of an overactive TGF- β pathway in diabetic retinopathy, may need to be taken into account in interpreting the encouraging results obtained with candesartan in the primary prevention of retinopathy in normotensive Type 1 diabetic patients (52). Orally active inhibitors of TGF- β signaling are in development (53), and would add options for highly targeted interventions. The next steps in these studies are to define the effects of selective normalization of TGF- β signaling in retinal vessels, and ascertain whether the TGF- β pathway is overactive in human diabetic retinopathy.

ACKNOWLEDGMENTS

We thank Drs. P.J. Oates for providing the sorbinil and A. Kazlauskas for the anti-RasGAP antibodies. This work was supported by grants from the National Institutes of Health (R01EY016206 to C.G., R01EY017637 to C.G. and M.L., and R01HG003354 to P.S.), the Juvenile Diabetes Research Foundation (JDRF) Center for Diabetic Retinopathy at the Schepens Eye Research Institute, the Massachusetts Lions Eye Research Fund, and the George and Frances Levin Endowment to M.L. Y.S.P. was the recipient of a JDRF Fellowship.

REFERENCES

1. Davis M: Diabetic retinopathy-A clinical overview. *Diabetes Care* 15:844-1874, 1992
2. Roy S, Maiello M, Lorenzi M: Increased expression of basement membrane collagen in human diabetic retinopathy. *J Clin Invest* 93:438-442, 1994
3. Mizutani M, Kern TS, Lorenzi M: Accelerated death of retinal microvascular cells in human and experimental diabetic retinopathy. *J Clin Invest* 97:2883-2890, 1996
4. The Writing Team for the Diabetes Control and Complications Trial/Epidemiology of Diabetes Interventions and Complications Research Group: Effect of intensive therapy on the microvascular complications of type 1 diabetes mellitus. *JAMA* 287:2563-2569, 2002
5. Lorenzi M: Mechanisms and strategies for prevention in diabetic retinopathy. *Curr Diab Rep* 6:102-107, 2006
6. Asnaghi V, Gerhardinger C, Hoehn T, Adeboje A, Lorenzi M: A role for the polyol pathway in the early neuroretinal apoptosis and glial changes induced by diabetes in the rat. *Diabetes* 52:506-511, 2003
7. Gerhardinger C, Brown LF, Roy S, Mizutani M, Zucker C, Lorenzi M: Expression of vascular endothelial growth factor in the human retina and in nonproliferative diabetic retinopathy. *Am J Pathol* 152:1453-1462, 1998
8. Gerhardinger C, McClure KD, Romeo G, Podestà F, Lorenzi M: IGF-I mRNA and signaling in the diabetic retina. *Diabetes* 50:175-183, 2001
9. Romeo G, Liu W-H, Asnaghi V, Kern TS, Lorenzi M: Activation of nuclear factor- κ B induced by diabetes and high glucose regulates a proapoptotic program in retinal pericytes. *Diabetes* 51:2241-2248, 2002
10. Dagher Z, Park YS, Asnaghi V, Hoehn T, Gerhardinger C, Lorenzi M: Studies of rat and human retinas predict a role for the polyol pathway in human diabetic retinopathy. *Diabetes* 53:2404-2411 2004
11. Sun W, Gerhardinger C, Dagher Z, Hoehn T, Lorenzi M: Aspirin at low-intermediate concentrations protects retinal vessels in experimental diabetic retinopathy through non-platelet-mediated effects. *Diabetes* 54:3418-3426, 2005
12. Chung SSM, Chung SK: Aldose reductase in microvascular complications. *Curr Drug Targets* 6:475-486, 2005
13. Kern TS, Engerman RL: Pharmacological inhibition of diabetic retinopathy. Aminoguanidine and aspirin. *Diabetes* 50:1636-1642, 2001
14. Klings ES, Safaya S, Adewoye AH, Odhiambo A, Frampton G, Lenburg M, Gerry N, Sebastiani P, Steinberg MH, Farber HW: Differential gene expression in pulmonary artery endothelial cells exposed to sickle cell plasma. *Physiol Genomics* 21:293-298, 2005
15. Tusher VG, Tibshirani R, Chu G: Significance analysis of microarrays applied to the ionizing radiation response. *Proc Natl Acad Sci USA*,98:5116-5121, 2001
16. Gerhardinger C, Biarnes Costa M, Coulombe MC, Toth I, Hoehn T, Grosu P: Expression of acute-phase response proteins in retinal Müller cells in diabetes. *Invest Ophthalmol Vis Sci* 46:349-357, 2005
17. Im E, Venkatakrisnan A, Kazlauskas A: Cathepsin B regulates the intrinsic angiogenic threshold of endothelial cells. *Mol Biol Cell* 16:3488-3500, 2005
18. ten Dijke P, Hill CS: New insights into TGF- β -smad signaling. *Trends Biochem Sci* 29:265-273, 2004
19. Chen L, Klass C, Woods A: Syndecan-2 regulates transforming growth factor- β signaling. *J Biol Chem* 279:15715-15718, 2004

20. Rouschop KM, Sewnath ME, Claessen N, Roelofs JJTH, Hoedemaeker I, van der Neut R, Aten J, Pals ST, Weening JJ, Florquin S: CD44 deficiency increases tubular damage but reduces renal fibrosis in obstructive nephropathy. *J Am Soc Nephrol* 15:674-686, 2004
21. Lin X, Liang M, Liang Y, Brunicardi FC, Melchior F, Feng X: Activation of transforming growth factor- β signaling by SUMO-1 modification of tumor suppressor Smad4/DPC4. *J Biol Chem* 278:18714-18719, 2003
22. Leask A, Abraham DJ: TGF- β signaling and the fibrotic response. *FASEB J* 18:816-827, 2004
23. Akool E, Doller A, Müller R, Gutwein P, Xin C, Huwiler A, Pfeilschifter J, Eberhardt W: Nitric oxide induces TIMP-1 expression by activating the transforming growth factor β -Smad signaling pathway. *J Biol Chem* 280:39403-39416, 2005
24. Akiyama N, Matsuo Y, Sai H, Noda M, Kizaka-Kondoh S: Identification of a series of transforming growth factor β -responsive genes by retrovirus-mediated gene trap screening. *Mol Cell Biol* 20:3266-3273, 2000
25. Han SH, Jeon JH, Ju HR, Jung U, Kim KY, Yoo HS, Lee YH, Song KS, Hwang HM, Na YS, Yang Y, Lee KN, Choi I: VDUP1 upregulated by TGF- β 1 and 125-dihydroxyvitamin D3 inhibits tumor cell growth by blocking cell-cycle progression. *Oncogene* 22:4035-4046, 2003
26. Goto K, Kamiya Y, Imamura T, Miyazono K, Miyazawa K: Selective inhibitory effects of Smad6 on bone morphogenetic protein type I receptors. *J Biol Chem* 282:20603-20611, 2007
27. Lindner V, Wang Q, Conley BA, Friesel RE, Vary CPH: Vascular injury induces expression of periostin. Implications for vascular cell differentiation and migration. *Arterioscler Thromb Vasc Biol* 25:77-83, 2005
28. Yang L, Yamasaki K, Shirakata Y, Dai X, Tokumaru S, Yahata Y, Tohyama M, Hanakawa Y, Sayama K, Hashimoto K: Bone morphogenetic protein-2 modulates Wnt and frizzled expression and enhances the canonical pathway of Wnt signaling in normal keratinocytes. *J Dermatol Sci* 42:111-119, 2006
29. Liu Z, Shi W, Ji X, Sun C, Jee WSS, Wu Y, Mao Z, Nagy TR, Li Q, Cao X: Molecules mimicking Smad1 interacting with Hox stimulate bone formation. *J Biol Chem* 279:11313-11319, 2004
30. Schulze PC, Yoshioka J, Takahashi T, He Z, King GL, Lee RT: Hyperglycemia promotes oxidative stress through inhibition of thioredoxin function by thioredoxin-interacting protein. *J Biol Chem* 279:30369-30374, 2004
31. Zhang J, Gerhardinger C, Lorenzi M: Early complement activation and decreased levels of glycosylphosphatidylinositol-anchored complement inhibitors in human and experimental diabetic retinopathy *Diabetes* 51:3499-3504, 2002
32. Zheng L, Szabo C, Kern TS: Poly(ADP-ribose) polymerase is involved in the development of diabetic retinopathy via regulation of nuclear factor- κ B. *Diabetes* 53:2960-2967, 2004
33. Moulding DA, Blundell MP, Spiller DG, White MRH, Cory GO, Calle Y, Kempinski H, Sinclair J, Ancliff PJ, Kinnon C, Jones GE, Thrasher AJ: Unregulated actin polymerization by WASp causes defects of mitosis and cytokinesis in X-linked neutropenia. *J Exp Med* 204:2213-2224, 2007
34. Moustakas A, Heldin C: Non-Smad TGF- β signals. *J Cell Sci* 118:3573-3584, 2005
35. Perlman R, Schiemann WP, Brooks MW, Lodish HF, Weinberg RA: TGF- β -induced apoptosis is mediated by the adapter protein Daxx that facilitates JNK activation. *Nat Cell Biol* 3:708-714, 2001

36. Takuwa Y, Takuwa N, Sugimoto N: The Edg family G protein-coupled receptors for lysophospholipids: their signaling properties and biological activities. *J Biochem* 131:767-771, 2002
37. Mizutani M, Gerhardinger C, Lorenzi M: Müller cell changes in human diabetic retinopathy. *Diabetes* 47:445-449, 1998
38. Ziyadeh FN, Hoffman BB, Han DC, Iglesias-de la Cruz MC, Hong SW, Isono M, Chen S, McGowan TA, Sharma K: Long-term prevention of renal insufficiency excess matrix gene expression and glomerular mesangial matrix expansion by treatment with monoclonal antitransforming growth factor- β antibody in *db/db* diabetic mice. *Proc Natl Acad Sci USA*,97:8015-8020, 2000
39. Khan ZA, Cukiernik M, Gonder JR, Chakrabarti S: Oncofetal fibronectin in diabetic retinopathy. *Invest Ophthalmol Vis Sci* 45:287-295, 2004
40. Kuiper EJ, Witmer AN, Klaassen I, Oliver N, Goldschmeding R, Schlingemann RO: Differential expression of connective tissue growth factor in microglia and pericytes in the human diabetic retina. *Br J Ophthalmol* 88:1082-1087, 2004
41. Wynn TA: Common and unique mechanisms regulate fibrosis in various fibroproliferative diseases. *J Clin Invest* 117:524-529, 2007
42. Weigert C, Sauer U, Brodbeck K, Pfeiffer A, Häring HU, Schleicher ED: AP-1 proteins mediate hyperglycemia-induced activation of the human TGF- β 1 promoter in mesangial cells. *J Am Soc Nephrol* 11:2007-2016, 2000
43. Ishii H, Tada H, Isogai S: An aldose reductase inhibitor prevents glucose-induced increase in transforming growth factor- β and protein kinase C activity in cultured human mesangial cells. *Diabetologia* 41:362-364, 1998
44. Obrosova IG, Pacher P, Szabó C, Zsengeller Z Hirooka H, Stevens MJ, Yorek MA: Aldose reductase inhibition counteracts oxidative-nitrosative stress and poly(ADP-ribose) polymerase activation in tissue sites for diabetes complications. *Diabetes* 54:234-242, 2005
45. Esposito K, Nappo F, Marfella R, Giugliano G, Giugliano F, Ciotola M, Quagliaro L, Ceriello A, Giugliano D: Inflammatory cytokine concentrations are acutely increased by hyperglycemia in humans: role of oxidative stress. *Circulation* 106:2067-2072, 2002
46. Roberson ED, Scarce-Levie K, Palop JJ, Yan F, Cheng IH, Wu T, Gerstein H, Yu G, Mucke L: Reducing endogenous tau ameliorates amyloid β -induced deficits in an Alzheimer's disease mouse model. *Science* 316:750-754, 2007
47. Makino H, Mukoyama M, Sugawara A, Mori K, Suganami T, Yahata K, Fujinaga Y, Yokoi H, Tanaka I, Nakao K: Roles of connective tissue growth factor and prostanoids in early streptozotocin-induced diabetic rat kidney: the effect of aspirin treatment. *Clin Exp Nephrol* 7:33-40, 2003
48. Taubert D, Berkels R, Grosser N, Schröder H, Gründemann D, Schömig E: Aspirin induces nitric oxide release from vascular endothelium: a novel mechanism of action. *Br J Pharmacol* 143:159-165, 2004
49. Kowluru RA, Kanwar M, Kennedy A: Metabolic memory phenomenon and accumulation of peroxynitrate in retinal capillaries. *Exp Diabetes Res* 2007:21976, 2007
50. Zacchigna L, Vecchione C, Notte A, Cordenonsi M, Dupont S, Maretto S, Cifelli G, Ferrari A, Maffei A, Fabbro C, Braghetta P, Marino G, Selvetella G, Aretini A, Colonnese C, Bettarini U, Russo G, Soligo S, Adorno M, Bonaldo P, Volpin D, Piccolo S, Lembo G, Bressan GM: Emilin1 links TGF- β maturation to blood pressure homeostasis. *Cell* 124:929-942, 2006

51. Brooke BS, Habashi JP, Judge DP, Patel N, Loeys B, Dietz HC: Angiotensin II blockade and aortic-root dilation in Marfan's syndrome. *N Engl J Med* 358:2787-2795, 2008
52. Chaturvedi N, Porta M, Klein R, Orchard T, Fuller J, Parving HH, Bilous R, Sjølie AK; DIRECT Programme Study Group. Effect of candesartan on prevention (DIRECT-Prevent 1) and progression (DIRECT-Protect 1) of retinopathy in type 1 diabetes: randomised, placebo-controlled trials. *Lancet* 372:1385-1393, 2008
53. Fu K, Corbley MJ, Sun L, Friedman JE, Shan F, Papadatos JL, Costa D, Lutterodt F, Sweigard H, Bowes S, Choi M, Boriack-Sjodin PA, Arduini RM, Sun D, Newman MN, Zahng X, Mead JN, Chuaqui CE, Cheung HK, Zhang X, Cornebise M, Carter MB, Josiah S, Sing J, Lee WC, Gill A, Ling LE: SM16 an orally active TGF- β type I receptor inhibitor prevents myofibroblast induction and vascular fibrosis in the rat carotid injury model. *Arterioscler Thromb Vasc Biol* 28:665-671, 2008

Table 1: Characteristics, retinal polyol pathway activity, and serum salicylates levels of study rats.

	n	Body weight (g)	HbA1c (%)	Retinal Sorbitol nmol/mg protein*	Retinal Fructose nmol/mg protein*	Serum salicylates mg/l [†]
Control	50	629 ± 77	5.8 ± 1.5	1.17 ± 0.40	3.85 ± 1.45	not tested
Diabetes	47	384 ± 50 [‡]	13.3 ± 1.9 [‡]	12.54 ± 8.64 [§]	65.32 ± 13.16 [‡]	Undetected
Diabetes with sorbinil	15	382 ± 48 [‡]	12.3 ± 1.4 [‡]	0.56 ± 0.25	6.40 ± 2.14	not tested
Diabetes with aspirin	15	370 ± 43 [‡]	12.9 ± 2.9 [‡]	not tested	not tested	13.5 ± 7.8

Values are mean ± SD. Statistical analysis was performed with ANOVA followed by Fisher's PLSD test. * Measurements were performed in 4 rats from each group. † Measurements were performed in 3 diabetic rats treated with aspirin and 1 untreated diabetic rat. ‡ $P < 0.0001$, § $P < 0.01$ compared with control rats.

Table 2: Functional clustering of the known genes differentially expressed in retinal vessels in diabetes.

Functional category *	Number of genes	Known function	upregulated	downregulated
TGF-β/BMPs pathway	20	TGF- β activation	16	
		BMPs activation	3	1
Oxidative stress	7	pro-oxidant	3	
		anti-oxidant	2	1
		pro/anti-oxidant	1	
Inflammation and response to injury	29	inflammatory mediators	5	
		acute-phase response proteins	9	
		IFN γ pathway	8	
		antigen presentation	3	1
		anti-inflammatory	3	
Matrix and vascular remodeling	35	matrix	4	1
		cell adhesion	7	1
		actin organization	3	2
		vascular remodeling	17	
Cell cycle	7	cell cycle progression	1	2
		cell cycle arrest	1	1
		proliferation inhibition	2	
Apoptosis	19	pro-apoptotic p53 pathway	5	2
		pro-apoptotic	7	1
		anti-apoptotic	2	2
Other	39	metabolism	8	1
		miscellaneous [†]	6	2
		function unknown [‡]	15	7

* The 127 known genes differentially expressed in diabetes were assigned to functional categories based on review of information from NCBI Entrez Gene integrated with data from published literature. Individual genes are listed in all pertinent categories based on their known functions. The complete list of the genes included in each functional category is presented in supplemental tables 2-8. [†] Includes genes related to functions that cannot be assigned to any of the major category; [‡] as stated in bibliographical sources.

FIGURE LEGENDS

Figure 1. Diabetes alters the expression of multiple genes of the TGF- β pathway (left side of the panel) and BMPs pathway (right side) in rat retinal vessels. Upregulated genes are shown in red, downregulated in blue. The expected effect of overexpression of genes on Smad signaling is shown by arrowed lines (stimulation) or blunt lines (inhibition). The expected effect of downregulation of Hoxc8 is release of inhibition, and thus activation of signaling (dotted blunt line).

Figure 2. Upregulation of the TGF- β 1 signaling loop in retinal vessels of diabetic rats. Total RNA and protein lysates were prepared from retinal vessels isolated by hypotonic lysis from the retina of diabetic (D) and age-matched control (C) rats. **(A)** TGF- β 1 mRNA levels assayed by quantitative RealTime PCR. Relative expression of TGF- β 1 mRNA was calculated by the comparative C_T method using β -actin as endogenous control. Values are mean \pm SD of = 6 rats per group. * $P = 0.04$.

(B) Representative Western blot of TGF β R1 (ALK5) and bar plot of the quantitative analysis. Blots were probed first for TGF β R1 followed by RasGAP as control for loading. Values are mean \pm SD of 3-5 rats per group. * $P < 0.04$. **(C)** Representative Western blots of phosphorylated Smad2 (P-Smad2) and total Smad2 (Smad2) and bar plot of the P-Smad2/Smad2 ratio. The P-Smad2/Smad2 ratio was examined in the retinal vessels of rats with 6 months (left panel) and 3 months of diabetes (right panel). The 58-60 kDa bands detected in both retinal microvessels and the + control (TGF- β -treated HepG2 cells) corresponds to Smad2, the 50 kDa band present in the + control only corresponds to Smad3. The bar plot presents the pooled results obtained in rats with 6 months and 3 months of diabetes. Values are mean \pm SD of 5-9 rats per group. * $P = 0.05$.

Figure 3. TGF- β 1 signaling is not increased in the neural retina of diabetic rats. Total RNA and protein lysates were prepared from the whole retina of diabetic (D) and age-matched control (C) rats. **(A)** TGF- β 1 mRNA levels assayed by quantitative RealTime PCR. Values are mean \pm SD of 4-5 rats per group. **(B)** Representative Western blot of TGF β R1 (ALK5) and bar plot of the quantitative analysis. Blots were probed first for TGF β R1 followed by β -actin as control for loading. Values are mean \pm SD of 5 rats per group. **(C)** Representative Western blots of phosphorylated Smad2 (P-Smad2) and total Smad2 (Smad2) and bar plot of the P-Smad2/Smad2 ratio. Values are mean \pm SD of 8-14 rats per group.

Figure 4. Effects of sorbinil and aspirin treatments on the gene expression changes induced by diabetes in rat retinal vessels. The white bars represent the number of genes differentially expressed in the retinal vessels of diabetic rats (untreated or treated) when compared to control rats. The black bars represent the number of genes whose expression is normalized by the drug (i.e. the expression in the retinal vessels of treated diabetic rats was not different from that in control rats). Each functional category includes all genes pertinent to that category as indicated in Table 2. Data were analyzed by Fisher's exact test. ns = non significant. D, diabetic rats; D+Sor, diabetic rats treated with sorbinil; D+Asa, diabetic rats treated with aspirin.

Figure 5. The increased TGF β RI immunoreactivity in retinal vessels of diabetic rats is prevented by sorbinil and aspirin treatments. TGF β RI was detected by immunohistochemistry in retinal trypsin digests from diabetic rats (8 months of diabetes

duration), diabetic rats treated with sorbinil or aspirin, and age-matched control rats. The effect of diabetes and of the two drugs on TGFβRI immunostaining was quantitated by four masked observers. **(A-C)** Representative photographs of mid-retina fields showing TGFβRI immunostaining of retinal capillaries. **(A)** control, score 2.0 ± 0.7 (mean \pm SD of scores by the different masked observers); **(B)** diabetes, score 3.8 ± 0.04 ; **(C)** diabetes treated with sorbinil, score 1 ± 0.0 ; **(D)** diabetes treated with aspirin, score 2.3 ± 0.7 . **(E)** Negative control. Scale bar = 100 micron. **(F)** Bar plot of the quantitative analysis of staining intensity; values are mean \pm SD of the final scores computed for each individual rat. C, control rats, n=11; D, diabetic rats, n=10; D+Sor, diabetic rats treated with sorbinil, n=6; D+Asa, diabetic rats treated with aspirin, n=6. * P=0.05 vs C, † P< 0.02 vs C, D, and D+Asa, ‡ P<0.04 vs D.

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5

