

Mitochondrial respiration is decreased in skeletal muscle of patients with type 2 diabetes

Running title: Mitochondrial dysfunction in type 2 diabetes

Received for publication 14 July 2006 and accepted in revised form 26 February 2007.

Martin Mogensen¹, Kent Sahlin^{1,2,3}, Maria Fernström^{2,3}, Dorte Glinborg⁴, Birgitte F. Vind⁴, Henning Beck-Nielsen⁴, Kurt Højlund⁴.

¹Institute of Sports Science and Clinical Biomechanics, University of Southern Denmark, Odense, Denmark

²Stockholm University College of Physical Education and Sports, GIH, Stockholm, Sweden

³Department of Physiology and Pharmacology, Karolinska Institutet, Stockholm, Sweden

⁴Diabetes Research Centre, Department of Endocrinology, Odense University Hospital, Denmark

Corresponding author:

Kurt Højlund

MD, PhD

Diabetes Research Centre, Department of Endocrinology, Odense University Hospital

Kloevervaenget 6, 3

DK-5000 Odense C

Denmark

E-mail: k.hojlund@dadlnet.dk

Abstract

We tested the hypothesis of a lower respiratory capacity per mitochondrion in skeletal muscle of type 2 diabetic (T2D) patients compared with obese subjects. Muscle biopsies obtained from 10 obese T2D and 8 obese nondiabetic male subjects were used for assessment of 3-hydroxy-Acyl-CoA-dehydrogenase (HAD) and citrate synthase (CS) activity, uncoupling protein 3 (UCP3) content, oxidative stress measured as 4-hydroxy-2-nonenal (HNE), fiber type distribution and respiration in isolated mitochondria. Respiration was normalized to CS activity (mitochondrial content) in isolated mitochondria. Maximal ADP-stimulated respiration (state 3) with pyruvate+malate and respiration through the electron transport chain (ETC) were reduced in T2D patients, and the proportion of type 2X fibers were higher in T2D patients compared with obese subjects (all $P < 0.05$). There were no differences in respiration with palmitoyl-L-carnitine+malate, CS activity, HAD activity, UCP3 content or oxidative stress measured as HNE between the groups. In the whole group, state 3 respiration with pyruvate+malate and respiration through ETC were negatively associated with HbA1c, and the proportion of type 2X fibers correlated with markers of insulin resistance ($P < 0.05$). In conclusion, we provide evidence for a functional impairment in mitochondrial respiration and increased amount of type 2X fibers in muscle of T2D patients. These alterations may contribute to the development of type 2 diabetes in humans with obesity.

Type 2 diabetes (T2D) is characterized by insulin resistance in major metabolic tissues such as skeletal muscle, liver and adipose tissue, and failure of the pancreatic β -cells to compensate for this abnormality (1). Skeletal muscle is the major site of glucose disposal in response to insulin, and correspondingly, the major site of insulin resistance in T2D (1,2). Despite extensive research, the mechanisms underlying insulin resistance are not fully understood. Indeed, several abnormalities have been identified in insulin resistant muscle including impaired insulin activation of glycogen synthase (1,3), impairment of the proximal components of the insulin signaling cascade (2), and increased intramuscular triglyceride content (4,5). Another important component of insulin resistance appears to be a decreased ability of insulin to regulate fuel utilization (6-8). In insulin resistant subjects this impaired ability to switch from lipid to carbohydrate oxidation in response to insulin has been described as “metabolic inflexibility” of skeletal muscle (8).

Being the site of fuel oxidation the mitochondrion has gained increasing interest in T2D research during the last decade. Several studies have indicated a role for mitochondrial dysfunction in the pathogenesis of insulin resistance and T2D. This includes reports of a decreased leg lipid oxidation in T2D and obesity (9) and a strong negative correlation between leg lipid oxidation and insulin resistance (10-13). Furthermore, microarray analysis has revealed a coordinated downregulation of genes involved in mitochondrial oxidative phosphorylation in T2D patients and high-risk individuals (14-16), and proteomic analysis has demonstrated a decreased content of the ATP synthase β -subunit in T2D patients (17). These findings are supported by reports of a decreased activity of the electron transport chain (ETC) in muscle mitochondria in T2D and obesity (18,19) and in vivo studies using magnetic resonance spectroscopy showing impaired mitochondrial ATP production in insulin resistant offspring of T2D patients (20).

Nevertheless, it remains to be fully clarified whether mitochondrial dysfunction in T2D and obesity is caused by a reduced content of mitochondria, a reduced functional capacity, or a combination of both. In a study by Kelley et al it was reported that the ratio between the activities of NADH: O_2 oxidoreductase and citrate synthase (CS) (a marker of mitochondria content) in skeletal muscle did not differ between lean and obese non-diabetic subjects and T2D patients (18). In a more

recent study from the same research group, ETC activity was measured as the activity of succinate oxidase in the subsarcolemmal and intermyofibrillar subpopulations of muscle mitochondria (19). Total ETC activity corrected for mitochondrial content (mitochondrial DNA) was reduced in both T2D and obesity compared with lean subjects. Moreover, ETC activity in the subsarcolemmal mitochondria fractions was reduced in T2D patients compared with obese subjects, but this was not corrected for mitochondrial content (19). Thus, it is still unclear whether the respiratory capacity per mitochondrion is reduced in T2D patients compared with obese non-diabetic subjects.

An essential feature of the mitochondrion is the coupling between substrate oxidation and phosphorylation of ADP through generation and utilization of the proton gradient over the inner mitochondrial membrane. Previous studies have suggested a possible link between insulin resistance and the coupling of these systems (21). The functionality of the mitochondrion can be assessed by measuring respiration in isolated mitochondria from human skeletal muscle under controlled conditions without the influence of circulating hormones and substrates (22). Here the intactness of the inner membrane is sustained enabling an interaction between substrate oxidation and phosphorylation. By using different substrate combinations different enzymatic pathways can be included and excluded. The mitochondrial respiratory rate will thereby describe the functionality and interaction between these pathways. By relating the mitochondrial respiration to CS activity the respiratory capacity per mitochondrion can be assessed. To our knowledge, no previous studies have investigated the respiratory function of the mitochondrion in skeletal muscle of T2D patients.

The present study was undertaken to investigate the role of mitochondrial dysfunction in insulin resistance in skeletal muscle of patients with T2D. To account for the effects of fiber type composition, mitochondrial content, obesity and oxidative stress induced by hyperglycemia, we measured the respiratory capacity per mitochondrion, oxidative stress as 4-hydroxy-2-nonenal (HNE), and fiber type distribution in skeletal muscle of T2D patients and well-matched obese, healthy subjects.

Method
Subjects

Ten patients with T2D and 8 healthy control male subjects, matched according to age and body mass index (BMI) participated in the study (Table 1). All subjects were obese and sedentary and did not participate in any kind of physical training. T2D patients were treated with either diet alone or diet in combination with sulfonylurea, metformin or insulin. Medication was withdrawn 1 week prior to the study. The patients were all GAD65-antibody negative and without signs of diabetic retinopathy, neuropathy, nephropathy or macrovascular complications. The control subjects had normal glucose tolerance and no family history of diabetes. All subjects had normal results on screening blood tests of hepatic and renal function. All subjects gave written informed consent and the study was approved by the local ethics committee of Funen and Vejle County and was performed in accordance with the Helsinki declaration. Participants were instructed to avoid vigorous exercise for 48 h before the study, which was carried out after a 10-h overnight fast. Fasting blood samples were analyzed for glucose (Glucose Analyzer II; Beckman Instruments, Fullerton, CA), glycosylated haemoglobin (HbA1c), free fatty acids (FFA) (Wako Chemicals GmbH, Neuss, Germany), triglycerides, total cholesterol, insulin, and C-peptide (Wallac Oy, Turku, Finland). Homeostasis model assessment of insulin resistance (HOMA-IR) was calculated from fasting values of serum insulin and plasma glucose (23).

Muscle biopsies and preparation of mitochondria

Muscle biopsies were taken from m. vastus lateralis using a modified Bergström needle with suction under local anaesthesia (5 ml of lidocaine 2%). A small part of the biopsy (~50-100 mg) was rapidly frozen in liquid nitrogen and stored at -80°C for later determination of oxidative enzyme activities, UCP3 protein content, fiber type distribution, and oxidative stress (HNE). The major part (~100-150 mg) of the biopsy was used for isolation of mitochondria as described previously (24). Using this method both the subsarcolemmal and intramyofibrillar mitochondrial fractions are isolated. Due to transportation (from Odense University Hospital to University of Southern Denmark, Odense) muscle samples was on average stored for 45 min (all < 1-h) in ice-cold isolation buffer before initiation of the isolation procedure. The major part of the isolated mitochondria was used for assessment of respiratory function. The remaining part was frozen in liquid nitrogen and stored at -80°C for later assessment of CS activity and maximal respiration through ETC.

Mitochondrial respiratory activity

Mitochondrial oxygen consumption was measured polarographically using a Clark-type electrode (DW1 oxygraph, Hansatech Instruments, Norfolk, England). The electrode was surrounded by a temperature controlled water-jacketed glass chamber maintaining the temperature at 25°C. The oxygraph was equipped with magnetic stirring and a gas tight plunger ensuring a minimum of oxygen diffusion between the chamber solution and the surroundings. For data collection, the electrode was connected to a computer. The measurement was carried out in an oxygraph medium containing 225 mmol/l mannitol, 75 mmol/l sucrose, 10 mmol/l Tris, 10 mmol/l KCl, 10 mmol/l K₂HPO₄, 0.1 mmol/l EDTA, 0.8 mmol/l MgCl₂·(6H₂O), and pH 7.0. 5 mmol/l pyruvate + 2 mmol/l L-malate and 10 μmol/l palmitoyl-L-carnitine + 2 mmol/l L-malate was used as substrate combinations. Mitochondrial respiration was initiated by the addition of mitochondrial rich suspension medium. After reaching a stable rate maximal respiration was initiated by adding K-ADP (final concentration 0.3 mmol/l). After determination of the maximal ADP-initiated respiration (termed V_{max} or state 3) and respiration without ADP (termed non-coupled respiration or state 4) the oxygen tension was regained and the submaximal respiration was initiated by low rate ADP infusion. This was accomplished by the use of a microdialysis pump (CMA/102, CMA/Microdialysis, Solna, Sweden) with a pumping rate ranging from 0.1 – 20 μl/min. At about 50% of state 3 respiration a sample was withdrawn from the respiratory chamber for calculation of ADP sensitivity ([ADP]_{1/2V_{max}}) as described previously (25).

Maximal respiration through ETC was determined in permeabilized mitochondria. Permeabilization was accomplished by addition of alamethicin to the respiratory medium (final concentration 6 μg/ml) (18). Maximal respiration was determined in the presence of 2 μmol/l cytochrome c and initiated by addition of 450 μmol/l NADH. The measurement of respiration through ETC in mitochondria from one control subject was excluded from further analysis due to signs of mitochondrial degeneration. This was confirmed by reevaluation of CS activity, which was only 25% of the expected value in the particular sample.

All respiratory measurements were related to CS activity in mitochondrial suspension (26). Respiratory control index (RCI = state 3/state 4), P/O ratio (mitochondrial efficiency) and P/O ratio

during submaximal respiration (33% of state 3) was calculated. RCI describes the coupled state of the mitochondrion and P/O ratio describes the efficiency of the mitochondrion, calculated as the amount of ADP consumed per oxygen being reduced. Both values were used to evaluate the success of the isolation procedure and the proton leak across the inner membrane. The P/O ratio at submaximal respiration was calculated to evaluate the mitochondrial efficiency while sustaining a membrane potential (27).

Enzyme activity, fiber type distribution and UCP3 content

The activity of 3-hydroxy-Acyl-CoA-dehydrogenase (HAD) and CS was determined spectrophotometrically at 25°C as described (28,29). CS and HAD activity was determined as unit per mg muscle protein per minute (Pierce, BCA™ Protein Assay Kit, Rockford, USA). Fiber type distribution was determined by electrophoresis as different isoforms of myosin heavy chain (type 1 fibers, type 2A fibers and type 2X fibers) as previously described (30). Determination of percentage of type 1 fibers by either this method or the histochemical method correlates significantly ($r=0.96$) (31). Determination of UCP3 protein content was analyzed in duplicate as described in detail previously (32).

Oxidative stress

4-Hydroxy-2-nonenal (HNE) is a major product of endogenous lipid peroxidation that reacts with several functional groups in muscle proteins to form thioesters and Michael adducts. Protein modification by HNE was performed by immunoblotting as described (33). Freeze-dried muscle samples were homogenized for 5 min in cold lysing buffer in 1:70 weight/volume, extracted for 15 min at 4°C with slow mixing and centrifuged at 1000g for 10 min. Protein concentration of the supernatant was determined with a protein assay kit (BCA™ Pierce). Equal amounts of proteins (15 µg) were loaded and separated on SDS-polyacrylamide gels (BioRad), and transferred to polyvinylidene difluoride membranes. After blocking in Tris-buffered saline with 5% non-fat milk, the membranes were incubated with primary polyclonal antibodies to the HNE adducts (Alpha Diagnostics) followed by incubation with horseradish peroxidase conjugated secondary antibody (anti-goat, Santa Cruz Bio Technology). The membrane was incubated with chemiluminescence detection reagent ECL (Amersham). Actin protein content was determined with primary antibodies to actin (anti-rabbit, Sigma) on the same membrane after

stripping the membrane in stripping buffer (Pierce). Quantification was performed with Quantity One 1-D Analysing software (Bio-Rad).

Statistics

Data are presented as mean ± SEM. Differences between T2D and control subjects were tested for statistical significance with the Students unpaired t-test. Correlation between variables was tested with Pearson Correlation analysis. Statistical significance was accepted at $P < 0.05$.

Results

Clinical, metabolic and muscle characteristics

T2D patients had significant higher concentrations of fasting plasma glucose and HbA1c compared with obese subjects (Table 1). There were no differences between groups in the circulating concentrations of C-peptide, insulin, FFA, total cholesterol, or triglycerides. Insulin resistance determined as HOMA-IR was significantly higher in diabetic patients compared with obese subjects.

Analyses of CS and HAD activity revealed no significant differences between T2D patients and obese subjects (Table 1). T2D patients had a significant higher proportion of type 2X fibers, but no differences in type 1 or type 2A fibers were found between the groups (Fig. 1A). Furthermore, there were no differences in muscle oxidative stress measured as HNE (Fig. 1B), or in UCP3 content measured per mg protein or per mitochondrion (CS activity) between the groups.

Mitochondrial characteristics

Maximal respiration through ETC was significantly reduced in T2D patients compared with obese subjects (Fig. 2A). Maximal ADP-stimulated respiration (state 3) with pyruvate+malate was also significantly reduced in T2D patients compared with obese subjects, whereas state 3 respiration with palmitoyl-L-carnitine+malate did not differ between the groups (Fig. 2B-C). In T2D patients RCI was significantly lower than in obese subjects, but again only when using pyruvate+malate as substrates (Table 2). T2D patients had a slightly lower P/O ratio using pyruvate+malate, but this did not reach statistical significance. There were no significant differences in P/O ratio during submaximal respiration or ADP sensitivity between the groups regardless of which substrates were used.

Correlation analysis

To explore potential mechanisms underlying reduced state 3 respiration and respiration through

ETC we examined the relationship between these measures of mitochondrial respiration and muscle and metabolic characteristics. In the whole study group (n=18) state 3 respiration with pyruvate+malate correlated negatively with HbA1c ($r = -0.51$, $P = 0.03$) (Fig. 3A) and tended to correlate with plasma glucose ($r = -0.46$, $P = 0.052$). Respiration through ETC was also inversely correlated with HbA1c ($r = -0.49$, $P = 0.048$) (Fig. 3B). State 3 respiration with palmitoyl-L-carnitine+malate was positively correlated with the proportion of type 1 fibers ($r = 0.50$, $P = 0.04$) and tended to correlate with HbA1c ($r = -0.46$, $P = 0.06$). Neither respiration through ETC or state 3 respiration correlated with the amount of type 2X fibers or HNE (oxidative stress).

We also studied the relationship between the amount of type 2X fibers and the other parameters measured. There was a significant positive correlation between the proportion of type 2X fibers and state 4 respiration using both pyruvate+malate ($r = 0.55$, $P = 0.02$) and palmitoyl-L-carnitine+malate ($r = 0.54$, $P = 0.02$) as substrates. Moreover the amount of type 2X fibers correlated positively with plasma glucose ($r = 0.48$, $P = 0.04$), FFA ($r = 0.53$, $P = 0.02$), insulin ($r = 0.55$, $P = 0.01$), C-peptide ($r = 0.66$, $P = 0.003$), and HOMA IR ($r = 0.76$, $P < 0.001$) (Fig. 3C-D).

Discussion

Several studies have implicated a role for mitochondrial dysfunction in the pathogenesis of skeletal muscle insulin resistance (6,7,14-20). However, evidence for a difference in mitochondrial function between obese non-diabetic subjects and T2D patients under basal conditions is scarce (19), and to which extent this is caused by a lower mitochondrial content or an impaired functional capacity remains unknown. The most important finding of the present study was a reduced respiratory function per mitochondrion in skeletal muscle of T2D patients compared with obese non-diabetic subjects. Moreover, in the whole study group the measures of mitochondrial respiration were negatively associated with HbA1c. However, muscle oxidative stress measured as HNE was similar in T2D patients and obese subjects, and no correlation between mitochondrial respiration and HNE content was observed. Interestingly, T2D patients also had a 2-fold higher proportion of type 2X muscle fibers compared with obese subjects. In the whole study group type 2X fiber content correlated with several markers of insulin resistance, but not with state 3 respiration or respiration through ETC. These results provide

evidence for a qualitative defect in the functional capacity of muscle mitochondria in T2D patients, which together with higher type 2X fiber content may play a role in the muscle metabolic changes leading to type 2 diabetes in humans with obesity.

Non-hypothesis-driven approaches using DNA microarrays and proteomics have demonstrated a coordinated down regulation of nuclear-encoded genes encoding proteins of the respiratory complexes (I-V) (14-16) and a decreased protein content and phosphorylation of the ATP-synthase β -subunit in skeletal muscle of T2D patients (17). These studies have pointed to abnormalities in the respiratory complexes I-V in the inner mitochondrial membrane. Correspondingly, ETC activity measured as the activity of NADH:O₂ oxidoreductase (18) or succinate oxidase (19) has been reported to be decreased in muscle mitochondria of T2D patients and obese subjects compared with lean subjects. Furthermore, the activity of succinate oxidase was found to be 50% lower in subsarcolemmal mitochondria in T2D patients compared with obese subjects. However, the NADH:O₂ oxidoreductase activity corrected for mitochondrial content (CS activity) was not different between lean subjects, obese non-diabetic subjects and T2D patients (18). Moreover, although succinate oxidase activity related to mtDNA was reduced in T2D patients and obese subjects compared with lean subjects, no difference in succinate oxidase activity corrected for mitochondrial content was reported between obese subjects and T2D patients in either the total population or the subsarcolemmal population of mitochondria (19).

We report a decreased respiratory function per mitochondrion in obese T2D patients compared with obese non-diabetic subjects. This was evident in fully coupled isolated mitochondria using pyruvate+malate (state 3 respiration) as substrates, and in permeabilized uncoupled mitochondria using NADH+cytochrome c (maximal respiration through ETC) as substrates. A decreased respiration per mitochondrion suggests an intrinsic inhibition or damage to some or more of the involved mitochondrial pathways. When using pyruvate+malate as substrates the respiration in isolated mitochondria is a result of a complex interplay between TCA cycle activity, ETC activity, utilization of the membrane potential, ATP-synthase functionality and the activity of adenine nucleotide translocase (ANT). Which of these factors are responsible for the reduced state 3 respiration with pyruvate+malate in T2D patients compared with control subjects can not be

determined from the present data. However, the concomitant finding of a significant lower respiration through ETC in T2D patients cannot be explained by a defect in the conversion of pyruvate to acetyl-CoA. Moreover, respiration with pyruvate+malate is not increased further by addition of substrates having the potential to increase NADH production indicating that pyruvate dehydrogenase and TCA cycle activity are unlikely to be the limiting factors. Together with the finding of similar CS activity in the two groups, these results suggest the possibility of a defect within the respiratory complexes I-V in the inner mitochondrial membrane. The failure to observe a significant difference in respiration with palmitoyl-L-carnitine+malate between obese non-diabetic and type 2 diabetic subjects could at least in part be explained by a lower respiration with palmitoyl-L-carnitine+malate than with pyruvate+malate in isolated mitochondria. Thus, with the same variations in these measurements, a larger sample size is probably required to prove significance. In vivo, however, mitochondrial oxidative metabolism is challenged by a number of substrates simultaneously, and hence a qualitative defect within the respiratory complexes I-V could have a more pronounced negative effect on respiration with lipids. Since only men with manifest T2D were included, further studies are needed to determine to which extent these findings apply to females with T2D, and whether they are potential early defects in the pathogenesis of T2D.

In the present study, HbA1c was negatively associated with state 3 respiration and respiration through ETC in the whole study group. This is in accordance with a previous study showing a significant inverse relationship between the content and phosphorylation of ATP-synthase β -subunit and glucose levels (17). These results seem to support the hypothesis of a relationship between prolonged elevation in glucose levels and mitochondrial dysfunction caused by an increased production of reactive oxygen species (ROS) and oxidative stress (34). Mitochondria are the major source of ROS and oxidative stress (~90%), and also the primary target for oxidative damage, when ROS production exceeds the capacity of the endogenous ROS-scavenging system (35). Accordingly, several protein subunits in the mitochondrial respiratory complexes (I-V) are susceptible to oxidative damage (36). However, to our knowledge, no previous studies have reported measurements of oxidative stress in skeletal muscle of insulin resistant subjects. Surprisingly, we found no difference in muscle oxidative stress measured

as HNE between T2D patients and obese subjects. Although, we did not measure other oxidative modifications such as nitration and carbonylation or local oxidative damage within mitochondria, it is likely that these measures would parallel those of HNE. Our data do not rule out a role for oxidative stress in mitochondrial dysfunction in both obesity and T2D, but indicate that the observed difference in quality of mitochondrial respiration between T2D patients and obese subjects is not mediated by increased oxidative stress. It is therefore too early to conclude that the observed negative association between mitochondrial respiration and HbA1c is a causal relationship involving increased ROS production.

Numerous factors are known to influence the mitochondrial function. Some of these factors, including age, obesity, fiber type distribution, physical activity and UCP3 protein, have also been related to insulin resistance and T2D. Physical activity has been shown to have a positive influence on T2D by increasing insulin sensitivity and inducing weight loss. Muscular adaptations in relation to increased physical fitness include an increased muscle oxidative capacity and increased lipid oxidative capacity mainly caused by an increased mitochondrial volume (24). Therefore, it could be argued that a potential lower level of physical activity in T2D patients contributed to the decrease in mitochondrial respiration. However, previous studies have shown a strong correlation between CS activity and VO_{2max} (37), and in our study CS activity in T2D patients and control subjects was in fact similar. Furthermore, several studies have shown that physical training does not affect measures of respiration per mitochondrion when using pyruvate and palmitoyl-L-carnitine with malate as substrates (24,25,38). Finally, the values of state 3 respiration with pyruvate+malate observed in the sedentary, obese non-diabetic subjects in the present study were similar to those previously reported in young healthy trained and untrained subjects (38,39). For these reasons, we find it unlikely that the decreased respiration per mitochondrion in T2D patients was due to a potential minor difference in physical activity.

In the present study, T2D patients had a 2-fold higher content of type 2X fibers. This is in agreement with some (40,41), but not all studies (42,43). Furthermore, in accordance with earlier reports of a close correlation between fiber type composition and insulin sensitivity (41,44,45), we found that type 2X fiber content was significantly associated with several markers of insulin resistance including HOMA-IR and levels of FFA.

Recent studies indicate that mitochondria from different fiber types may have different metabolic characteristics. This includes a higher UCP3 content in type 2X fibers in human skeletal muscle (46), and a lower lipid oxidative capacity per mitochondria and higher hydrogenperoxide production in type 2 fibers from rodent muscle (27,47). Correspondingly, we observed a positive association between the proportion of type 1 fibers and mitochondrial respiration with palmitoyl-L-carnitine+malate and a correlation between type 2X fibers and non-coupled (state 4) respiration. In humans, muscle fiber type composition is more mixed than in rodents and generally type 2X fibers have a very low oxidative capacity. Although there was no relationship between mitochondrial respiration (state 3 or through ETC) and the proportion of type 2X fibers, we cannot rule out the possibility that decreased mitochondrial respiration in T2D patients was in part explained by an increased proportion of type 2X fibers compared with obese subjects. On the other hand, single-fiber analysis has demonstrated that reduced oxidative capacity and increased lipid content in skeletal muscle of T2D patients are independent of the effect of fiber type (42).

Previous studies have indicated that UCP3 may play an essential role in mitochondrial dysfunction in T2D patients (48). Despite extensive research the main function of UCP3 is still unknown. UCP3 has been hypothesized to function as a protective mechanism against ROS production by mild uncoupling of the inner mitochondrial membrane and as a transporting protein of non-esterified fatty acids from the matrix to the intermembrane space protecting the mitochondrion against peroxidized lipids (49). At present there is little evidence that UCP3 functions as a fatty acid transporter. However, there is some evidence supporting a role of UCP3 in the attenuation of mitochondrial ROS production (49). In the present study, we found no difference in UCP3 protein levels between T2D patients and obese subjects. This is in contrast to a recent study showing a

decreased UCP3 content in T2D patients (50) probably due to different matching of the study subjects with respect to age, body composition and physical activity. Our findings strongly indicate that the decrease in respiration per mitochondrion in T2D patients compared with obese subjects is not explained by changes in UCP3 protein. On the other hand, even normal levels of UCP3 levels may contribute to mitochondrial damage in T2D patients in face of an increased oxidative stress (34).

In summary, we provide evidence for a qualitative defect in the respiratory function per mitochondrion in skeletal muscle of T2D patients compared with obese non-diabetic subjects. This difference was not paralleled by increased oxidative stress in diabetic muscle. Moreover, the proportion of type 2X fibers was higher in T2D patients and correlated with several markers of insulin resistance, but not with mitochondrial respiration. These alterations may be important factors contributing to the progression from obesity to T2D in humans. We were not able to demonstrate a role for either UCP3 protein levels, oxidative stress or fiber type composition in the respiratory capacity of mitochondria, but a significant negative correlation between mitochondrial respiration and HbA1c indicates that elevated glucose levels may contribute to mitochondrial dysfunction in obese and T2D subjects perhaps by other mechanisms than oxidative stress.

Acknowledgments

The authors wish to express their gratitude to L. Hansen, C. B. Olsen and C. Christensen for excellent technical assistance. The work was supported by grants from the Danish Medical Research Council, the Novo Nordisk Foundation, the Danish Diabetes Association and the Institute of Clinical Research, University of Southern Denmark. M. M and K. H. was supported by fellowships from the Danish Medical Research Council.

References

1. Beck-Nielsen H, Vaag A, Poulsen P, Gaster M: Metabolic and genetic influence on glucose metabolism in type 2 diabetic subjects - experiences from relatives and twin studies. *Best Pract Res Clin Endocrinol Metab* 17:445-467, 2003
2. Petersen KF, Shulman GI: Etiology of insulin resistance. *Am J Med* 119:S10-S16, 2006
3. Hojlund K, Staehr P, Hansen BF, Green KA, Hardie DG, Richter EA, Beck-Nielsen H, Wojtaszewski JF: Increased phosphorylation of skeletal muscle glycogen synthase at NH₂-terminal sites during physiological hyperinsulinemia in type 2 diabetes. *Diabetes* 52:1393-1402, 2003
4. Kelley DE, Goodpaster BH, Storlien L: Muscle triglyceride and insulin resistance. *Annu Rev Nutr* 22:325-346, 2002
5. Levin K, Daa Schroeder H, Alford FP, Beck-Nielsen H: Morphometric documentation of abnormal intramyocellular fat storage and reduced glycogen in obese patients with Type II diabetes. *Diabetologia* 44:824-833, 2001
6. Brehm A, Krssak M, Schmid AI, Nowotny P, Waldhausl W, Roden M: Increased lipid availability impairs insulin-stimulated ATP synthesis in human skeletal muscle. *Diabetes* 55:136-140, 2006
7. Stump CS, Short KR, Bigelow ML, Schimke JM, Nair KS: Effect of insulin on human skeletal muscle mitochondrial ATP production, protein synthesis, and mRNA transcripts. *Proc Natl Acad Sci U S A* 100:7996-8001, 2003
8. Kelley DE, Mandarino LJ: Fuel selection in human skeletal muscle in insulin resistance: a reexamination. *Diabetes* 49:677-683, 2000
9. Kelley DE, Simoneau JA: Impaired free fatty acid utilization by skeletal muscle in non-insulin-dependent diabetes mellitus. *J Clin Invest* 94:2349-2356, 1994
10. Kelley DE, Goodpaster B Wing RR, Simoneau JA: Skeletal muscle fatty acid metabolism in association with insulin resistance, obesity, and weight loss. *Am J Physiol* 277:E1130-E1141, 1999
11. Simoneau JA, Kelley DE: Altered glycolytic and oxidative capacities of skeletal muscle contribute to insulin resistance in NIDDM. *J Appl Physiol* 83:166-171, 1997
12. Kim JY, Hickner RC, Cortright RL, Dohm GL, Houmard JA: Lipid oxidation is reduced in obese human skeletal muscle. *Am J Physiol Endocrinol Metab* 279:E1039-E1044, 2000
13. Simoneau JA, Veerkamp JH, Turcotte LP, Kelley DE: Markers of capacity to utilize fatty acids in human skeletal muscle: relation to insulin resistance and obesity and effects of weight loss. *FASEB J* 13:2051-2060, 1999
14. Patti ME, Butte AJ, Crunkhorn S, Cusi K, Berria R, Kashyap S, Miyazaki Y, Kohane I, Costello M, Saccone R, Landaker EJ, Goldfine AB, Mun E, DeFronzo R, Finlayson J, Kahn CR, Mandarino LJ: Coordinated reduction of genes of oxidative metabolism in humans with insulin resistance and diabetes: Potential role of PGC1 and NRF1. *Proc Natl Acad Sci U S A* 100:8466-8471, 2003
15. Mootha VK, Lindgren CM, Eriksson KF, Subramanian A, Sihag S, Lehar J, Puigserver P, Carlsson E, Ridderstrale M, Laurila E, Houstis N, Daly MJ, Patterson N, Mesirov JP, Golub TR, Tamayo P, Spiegelman B, Lander ES, Hirschhorn JN, Altshuler D, Groop LC: PGC-1 α -responsive genes involved in oxidative phosphorylation are coordinately downregulated in human diabetes. *Nat Genet* 34:267-273, 2003

16. Sreekumar R, Halvatsiotis P, Schimke JC, Nair KS: Gene expression profile in skeletal muscle of type 2 diabetes and the effect of insulin treatment. *Diabetes* 51:1913-1920, 2002
17. Hojlund K, Wrzesinski K, Larsen PM, Fey SJ, Roepstorff P, Handberg A, Dela F, Vinten J, McCormack JG, Reynet C, Beck-Nielsen H: Proteome analysis reveals phosphorylation of ATP synthase beta -subunit in human skeletal muscle and proteins with potential roles in type 2 diabetes. *J Biol Chem* 278:10436-10442, 2003
18. Kelley DE, He J, Menshikova EV, Ritov VB: Dysfunction of mitochondria in human skeletal muscle in type 2 diabetes. *Diabetes* 51:2944-2950, 2002
19. Ritov VB, Menshikova EV, He J, Ferrell RE, Goodpaster BH, Kelley DE: Deficiency of subsarcolemmal mitochondria in obesity and type 2 diabetes. *Diabetes* 54:8-14, 2005
20. Petersen KF, Dufour S, Befroy D, Garcia R, Shulman GI: Impaired mitochondrial activity in the insulin-resistant offspring of patients with type 2 diabetes. *N Engl J Med* 350:664-671, 2004
21. Iossa S, Mollica MP, Lionetti L, Crescenzo R, Tasso R, Liverini G: A possible link between skeletal muscle mitochondrial efficiency and age-induced insulin resistance. *Diabetes* 53:2861-2866, 2004
22. Madsen K, Ertbjerg P, Pedersen PK: Calcium content and respiratory control index of isolated skeletal muscle mitochondria: effects of different isolation media. *Anal Biochem* 237:37-41, 1996
23. Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RC: Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia* 28:412-419, 1985
24. Tonkonogi M, Sahlin K: Rate of oxidative phosphorylation in isolated mitochondria from human skeletal muscle: effect of training status. *Acta Physiol Scand* 161:345-353, 1997
25. Mogensen M, Bagger M, Pedersen PK, Fernstrom M, Sahlin K: Cycling efficiency in humans is related to low UCP3 content and to type I fibres but not to mitochondrial efficiency. *J Physiol* 2006
26. Wibom R, Hagenfeldt L, von DU: Measurement of ATP production and respiratory chain enzyme activities in mitochondria isolated from small muscle biopsy samples. *Anal Biochem* 311:139-151, 2002
27. Mogensen M, Sahlin K: Mitochondrial efficiency in rat skeletal muscle: influence of respiration rate, substrate and muscle type. *Acta Physiol Scand* 185:229-236, 2005
28. Passonneau JV, Lowry OH: *In Enzymatic Analysis. A practical guide.* Human Press, 1993,
29. Alp PR, Newsholme EA, Zammit VA: Activities of citrate synthase and NAD⁺-linked and NADP⁺-linked isocitrate dehydrogenase in muscle from vertebrates and invertebrates. *Biochem J* 154:689-700, 1976
30. Andersen JL, Aagaard P: Myosin heavy chain IIX overshoot in human skeletal muscle. *Muscle Nerve* 23:1095-1104, 2000
31. Hansen EA, Andersen JL, Nielsen JS, Sjogaard G: Muscle fibre type, efficiency, and mechanical optima affect freely chosen pedal rate during cycling. *Acta Physiol Scand* 176:185-194, 2002
32. Tonkonogi M, Fernstrom M, Walsh B, Ji LL, Rooyackers O, Hammarqvist F, Wernerman J, Sahlin K: Reduced oxidative power but unchanged antioxidative capacity in skeletal muscle from aged humans. *Pflugers Arch* 446:261-269, 2003

33. Shabalina IG, Petrovic N, Kramarova TV, Hoeks J, Cannon B, Nedergaard J: UCP1 and defense against oxidative stress. 4-Hydroxy-2-nonenal effects on brown fat mitochondria are uncoupling protein 1-independent. *J Biol Chem* 281:13882-13893, 2006
34. Evans JL, Goldfine ID, Maddux BA, Grodsky GM: Oxidative stress and stress-activated signaling pathways: a unifying hypothesis of type 2 diabetes. *Endocr Rev* 23:599-622, 2002
35. Fridlyand LE, Philipson LH: Reactive species and early manifestation of insulin resistance in type 2 diabetes. *Diabetes Obes Metab* 8:136-145, 2006
36. Choksi KB, Boylston WH, Rabek JP, Widger WR, Papaconstantinou J: Oxidatively damaged proteins of heart mitochondrial electron transport complexes. *Biochim Biophys Acta* 1688:95-101, 2004
37. Blomstrand E, Radegran G, Saltin B: Maximum rate of oxygen uptake by human skeletal muscle in relation to maximal activities of enzymes in the Krebs cycle. *J Physiol* 501 (Pt 2):455-460, 1997
38. Fernstrom M, Tonkonogi M, Sahlin K: Effects of acute and chronic endurance exercise on mitochondrial uncoupling in human skeletal muscle. *J Physiol* 554:755-763, 2004
39. Tonkonogi M, Krook A, Walsh B, Sahlin K: Endurance training increases stimulation of uncoupling of skeletal muscle mitochondria in humans by non-esterified fatty acids: an uncoupling-protein-mediated effect? *Biochem J* 351 Pt 3:805-810, 2000
40. Oberbach A, Bossenz Y, Lehmann S, Niebauer J, Adams V, Paschke R, Schon MR, Bluher M, Punkt K: Altered fiber distribution and fiber-specific glycolytic and oxidative enzyme activity in skeletal muscle of patients with type 2 diabetes. *Diabetes Care* 29:895-900, 2006
41. Hickey MS, Carey JO, Azevedo JL, Houmard JA, Pories WJ, Israel RG, Dohm GL: Skeletal muscle fiber composition is related to adiposity and in vitro glucose transport rate in humans. *Am J Physiol* 268:E453-E457, 1995
42. He J, Watkins S, Kelley DE: Skeletal muscle lipid content and oxidative enzyme activity in relation to muscle fiber type in type 2 diabetes and obesity. *Diabetes* 50:817-823, 2001
43. Zierath JR, He L, Guma A, Odegaard WE, Klip A, Wallberg-Henriksson H: Insulin action on glucose transport and plasma membrane GLUT4 content in skeletal muscle from patients with NIDDM. *Diabetologia* 39:1180-1189, 1996
44. Lillioja S, Young AA, Culter CL, Ivy JL, Abbott WG, Zawadzki JK, Yki-Jarvinen H, Christin L, Secomb TW, Bogardus C: Skeletal muscle capillary density and fiber type are possible determinants of in vivo insulin resistance in man. *J Clin Invest* 80:415-424, 1987
45. Marin P, Andersson B, Krotkiewski M, Bjorntorp P: Muscle fiber composition and capillary density in women and men with NIDDM. *Diabetes Care* 17:382-386, 1994
46. Russell AP, Wadley G, Hesselink MK, Schaart G, Lo S, Leger B, Garnham A, Kornips E, Cameron-Smith D, Giacobino JP, Muzzin P, Snow R, Schrauwen P: UCP3 protein expression is lower in type I, IIa and IIx muscle fiber types of endurance-trained compared to untrained subjects. *Pflugers Arch* 445:563-569, 2003
47. Anderson EJ, Neuffer PD: Type II skeletal myofibers possess unique properties that potentiate mitochondrial H₂O₂ generation. *Am J Physiol Cell Physiol* 290:C844-C851, 2006
48. Schrauwen P, Hesselink MK: Oxidative capacity, lipotoxicity, and mitochondrial damage in type 2 diabetes. *Diabetes* 53:1412-1417, 2004

49. Brand MD, Esteves TC: Physiological functions of the mitochondrial uncoupling proteins UCP2 and UCP3. *Cell Metab* 2:85-93, 2005
50. Schrauwen P, Mensink M, Schaart G, Moonen-Kornips E, Sels JP, Blaak EE, Russell AP, Hesselink MK: Reduced skeletal muscle UCP3 protein content in pre-diabetic subjects and type 2 diabetic patients: restoration by rosiglitazone treatment. *J Clin Endocrinol Metab* 2005

Table 1 Clinical, metabolic and muscle characteristics

	T2D	Control	P-value
Age (years)	55 ± 2	53 ± 2	0.36
Body mass index (kg/m ²)	31 ± 1	32 ± 1	0.77
Weight (kg)	99 ± 4	106 ± 6	0.36
Plasma glucose (mmol/l)	11.5 ± 1.6	5.8 ± 0.1	0.006
HbA1c (%)	7.4 ± 0.4	5.5 ± 0.1	0.001
C-peptide (pmol/l)	1312 ± 167	929 ± 97	0.08
Insulin (pmol/l)	100 ± 24	76 ± 12	0.41
FFA (mmol/l)	0.6 ± 0.1	0.6 ± 0.1	0.73
Total cholesterol (mmol/l)	5.7 ± 0.3	5.4 ± 0.3	0.60
Triglyceride (mmol/l)	1.4 ± 0.2	2.9 ± 1.1	0.28
HOMA-IR	8.0 ± 1.7	3.3 ± 0.51	0.03
CS activity	104 ± 8	110 ± 11	0.63
HAD activity	160 ± 13	161 ± 11	0.95
UCP3	1.4 ± 0.2	1.2 ± 0.3	0.69
UCP3 per mitochondrion	1.3 ± 0.2	1.1 ± 0.2	0.44

Values are mean ± SEM from 10 T2D patients and 8 obese control subjects. CS and HAD activity was measured as units · mg protein⁻¹ · min⁻¹. UCP3 was determined as arbitrary units (AU) per mg protein. UCP3 per mitochondrion was determined as mAU per CS activity.

Table 2 Mitochondrial characteristics

	T2D	Control	P-value
Pyruvate+malate			
RCI	11.5 ± 1.0	14.4 ± 0.8	0.04
P/O ratio	2.47 ± 0.03	2.63 ± 0.08	0.08
P/O ratio _{submax}	2.24 ± 0.02	2.25 ± 0.04	0.92
[ADP] _{½V_{max}}	29.7 ± 5.7	29.6 ± 8.0	0.99
Palmitoyl-L-carnitine+malate			
RCI	7.2 ± 0.8	7.9 ± 0.6	0.47
P/O ratio	2.30 ± 0.03	2.35 ± 0.03	0.32
P/O ratio _{submax}	1.67 ± 0.08	1.64 ± 0.04	0.77
[ADP] _{½V_{max}}	9.8 ± 1.9	9.5 ± 1.1	0.92

Values are mean ± SEM from 10 T2D patients and 8 obese control subjects. RCI is state 3 divided by state 4 respiration. P/O ratio is the amount of ADP used per oxygen atom during state 3 respiration. P/O ratio_{submax} is P/O ratio determined during submaximal respiration. [ADP]_{½V_{max}} is the concentration of ADP needed to induce 50% of state 3 respiration, a measure of mitochondrial ADP sensitivity.

Figure legends

Figure 1

Fiber type distribution and oxidative stress in human skeletal muscle from obese patients with type 2 diabetes (T2D) and obese control subjects. (A) Proportions (%) of type 1, type 2A and type 2X fibers in skeletal muscle from 10 diabetic (black bars) and 8 control (white bars) subjects. Data are mean \pm SEM. (B) Oxidative stress measured as 4-hydroxy-2-nonenal (HNE) in skeletal muscle of 10 diabetic (closed circles) and 8 control (open circles) subjects. The mean value for each group is indicated by a line. *P < 0.05 vs. controls.

Figure 2

Respiration in isolated mitochondria from human skeletal muscle of obese patients with type 2 diabetes (T2D) and obese control subjects. (A) Maximal respiration through the electron transport chain (ETC) in permeabilized isolated mitochondria using NADH and cytochrome c as substrates. Data are from 10 diabetic (closed circles) and 7 control (open circles) subjects. The mean value for each group is indicated by a line. (B) State 3 and state 4 respiration in isolated mitochondria with pyruvate+malate (Pyr) and (C) palmitoyl-L-carnitine+malate (Pal) as substrate combinations. Data represent mean \pm SEM from 10 diabetic (black bars) and 8 control (white bars) subjects. All measures of respiration were divided by CS activity (U CS) in the mitochondrial suspension to correct for mitochondrial content. *P < 0.05 vs. controls.

Figure 3

Correlation analysis in the whole study group (n =18). The levels of HbA1c were negatively associated with (A) state 3 respiration with pyruvate+malate (nmol/min/U CS) and (B) respiration through the electron transport chain (ETC), (nmol/min/U CS). The proportion of type 2X fibers was positively associated with (C) FFA levels and (D) HOMA-IR.

Figure 1




Figure 2




Figure 3

