

**Phosphorylation Barriers to Skeletal and Cardiac Muscle Glucose Uptakes
in High Fat Fed Mice:
*Studies in Mice with a 50% Reduction of Hexokinase II***

Patrick T. Fueger¹, Robert S. Lee-Young¹, Jane Shearer¹, Deanna P. Bracy^{1,3},
Sami Heikkinen⁴, Markku Laakso^{4,5}, Jeffrey N. Rottman^{2,3}, and David H. Wasserman^{1,3}

¹ Department of Molecular Physiology and Biophysics, ² Department of Internal Medicine, ³
Mouse Metabolic Phenotyping Center, Vanderbilt University School of Medicine, Nashville,
TN, 37232

⁴ A. I. Virtanen Institute and Department of Biochemistry and Biotechnology, ⁵ Department of
Medicine, University of Kuopio, Kuopio, Finland, 70211

Running Head: Muscle glucose uptake in high fat-fed HK^{+/-} mice

Current address for correspondence:
Patrick T. Fueger, PhD
Duke University Medical Center
Department of Pharmacology and Cancer Biology
4321 Medical Park Drive, Suite 200
Durham, NC 27704
e-mail: patrick.fueger@duke.edu

Received for publication 19 April 2007 and accepted in revised form 11 July 2007.

ABSTRACT

Objective. Muscle glucose uptake (MGU) is regulated by glucose delivery to, transport into, and phosphorylation within muscle. The aim of this study was to determine the role of limitations in glucose phosphorylation in the control of MGU during either physiological insulin stimulation ($4 \text{ mU}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$) or exercise with chow (CH) or high fat (HF) feeding.

Research Design and Methods. C57BL/6J mice with (HK^{+/-}) and without (WT) a 50% hexokinase II deletion were fed CH or HF diets and studied at 4 mo of age during a 120 min insulin clamp or 30 min of treadmill exercise (n = 8-10 mice/group). 2-deoxy[³H]glucose was used to measure R_g, an index of MGU.

Results. Body weight and fasting arterial glucose were increased by HF feeding but not HK^{+/-}. Both HF feeding and HK^{+/-} independently created fasting hyperinsulinemia, a response that was increased synergistically with combined HF feeding and HK^{+/-}. Whole-body insulin action was suppressed by ~25% with either HF feeding or HK^{+/-} alone, but by over 50% when the two were combined. Insulin-stimulated R_g was modestly impaired by HF feeding and HK^{+/-} independently (~15 to 20%), but markedly reduced by the two together (~40 to 50%). Exercise-stimulated R_g was reduced by ~50% with HF feeding and HK^{+/-} alone and was not attenuated further by combining the two.

Conclusions. In summary, impairments in whole body metabolism and MGU due to HF feeding and HK^{+/-} combined during insulin stimulation are additive. In contrast, combining HF feeding and HK^{+/-} during exercise causes no greater impairment in MGU than the two manipulations independently. This suggests that MGU is impaired during exercise by HF feeding due, in large part, to a limitation in glucose phosphorylation. Together, these studies show that the high fat fed mouse is characterized by defects at multiple steps of the MGU system that are precipitated by different physiological conditions.

Key words: hexokinase, glucose transport, 2-deoxyglucose, glycogen, clamp

The importance of limitations in muscle glucose phosphorylation capacity in insulin resistance is a point of contention. In rodent muscle, glucose phosphorylation to glucose 6-phosphate is catalyzed primarily by the hexokinase (HK) II isozyme. HK II content may be normal (1) or reduced (2) in insulin resistant states depending on the muscle fiber type composition. Regardless, glucose phosphorylation capacity is affected due to allosteric inhibition of HK II by acyl CoA and glucose 6-phosphate or compartmentation of HK II, its allosteric regulators, and its substrates (i.e. glucose and ATP). Insulin stimulation and exercise can be used to expose functional deficits in glucose phosphorylation capacity, that may otherwise be undetectable (3). Indeed, we previously employed such a strategy in conscious mice overexpressing HK II to provide evidence that glucose phosphorylation is, in fact, impaired by high fat feeding (4). This previous study showed that HK II overexpression does not cause a unique phenotype in the basal state, is unable to correct impairments in insulin-stimulated muscle glucose uptake (MGU) due to high fat feeding, but largely corrects impairments in exercise-stimulated MGU.

The aims of this study were to determine whether diminished glucose phosphorylation capacity induced by a genetic 50% reduction in HK II: a) impairs insulin-stimulated MGU in chow fed mice, b) worsens the insulin resistance of high fat fed mice; and c) further attenuates exercise-stimulated MGU observed with high fat feeding and accelerates the onset of fatigue. To this end, mice with (HK^{+/-}) and without (WT) a 50% HK II deletion fed either a standard chow or high fat diet were studied during insulin clamps and exercise. One premise underlying these studies was that if limitations to glucose phosphorylation were a cause of impaired MGU resulting from high fat feeding, then removal of 50% of HK II

protein should have little additional effect when combined with high fat feeding. Reciprocally, additivity would predict that the mechanism of impairment created by genetic and dietary factors are likely to be distinct. It was hypothesized that HK II reduction in high fat fed mice would further impair MGU during an insulin clamp, but not during exercise. Furthermore, it was hypothesized that a reduction in HK II would not affect exercise endurance, a functional consequence of metabolic dysregulation, in high fat fed mice.

EXPERIMENTAL PROCEDURES

Mouse maintenance and genotyping.

Procedures were approved by the Vanderbilt University Animal Care and Use Committee. Male Ukko1 (a mixture of BALB/c and DBA/2 strains) mice containing a partial deletion to the HK II gene (HK^{+/-}) that results in a 50% reduction in HK II activity in heart, skeletal muscle, and adipose tissue, but does not change HK I activity (5) were backcrossed onto the C57BL/6J background for at least 5 generations (6). At 3 wks of age, littermates were separated by gender and maintained in microisolator cages. Genotyping was performed as previously described (5) with the polymerase chain reaction on genomic DNA obtained from a tail biopsy and isolated by a DNeasy Tissue Kit (Qiagen, Valencia, CA). Upon weaning at 3 wks of age mice were fed standard chow or a high fat diet *ad libitum* and were studied at ~4 mo of age. The high fat (HF) diet contained 35.5% fat (Diet F3282, Bio-Serv, Inc., Frenchtown, NJ) whereas the chow diet contained 5.5 % fat (5001 Laboratory Rodent Diet, Purina, Richmond, IN).

Immunoblotting. Total GLUT4 and HKII protein were determined in homogenates of gastrocnemius and cardiac muscles. 20 µg of protein were resolved on 4-12% Bis-Tris SDS-PAGE gels, followed by

electrophoretic transfer to polyvinylidene fluoride membranes. Membranes were blocked with 1X milk buffer (Chemicon; Temecula, CA) for 15 minutes, probed with rabbit anti-GLUT4 (1:1000; Abcam) overnight at 4° C, and then incubated with anti-rabbit horseradish peroxidase (1:20000; Pierce, Rockford, IL, USA) for 1 h at 23° C. Membranes were exposed to chemiluminescent substrate and imaged using the VersaDoc imaging system (Bio-Rad; Hercules, CA). Membranes were stripped with a Re-Blot Western Blot Recycling Kit (Chemicon), probed with rabbit anti-HKII (1:1000; Chemicon), incubated with anti-rabbit horseradish peroxidase (1:20000), and developed as before. To confirm equal protein loading and transfer, membranes were stripped and re-probed with monoclonal anti-glyceraldehyde-3-phosphate dehydrogenase (GAPDH, 1:4000, Abcam) and then incubated with anti-mouse horseradish peroxidase (1:20000). All antibodies were diluted in 1% polyvinylpyrrolidone in TBS, and membranes were washed between antibody incubations three times with TBS for 10 min. Densitometry was performed using Quantity One Analysis Software (Bio-Rad Laboratories; Hercules, CA).

Echocardiography and blood pressure measurement. At 16 weeks of age transthoracic echocardiograms were performed on resting conscious mice using a 15-MHz transducer (Sonos 5500 system, Agilent) as previously described (7; 8). In addition, systolic blood pressure was measured in conscious mice using tail cuff plethysmography (9). Echocardiographic and blood pressure measurements were made twice and thrice, respectively, to account for acclimation.

Surgical procedures. Surgical procedures were similar to those described previously (10; 11) Briefly, mice of either sex were anesthetized with pentobarbital (70 mg/kg body weight) and the left common

carotid artery and the right jugular vein were catheterized for sampling and infusions, respectively. Free catheter ends were tunneled under the skin to the back of the neck, where they were attached via stainless steel connectors to lines made of Micro-Renathane (0.033 in OD), which were exteriorized and sealed with stainless steel plugs. Lines were kept patent by flushing daily with 10-40 µl saline containing 200 U/ml of heparin and 5 mg/ml of ampicillin. Animals were housed individually after surgery and body weight was recorded daily.

Hyperinsulinemic-euglycemic clamp and saline control experiments. Experiments were performed on 5 h fasted mice as previously described (4; 12-14). For hyperinsulinemic-euglycemic clamp and saline control experiments, 4 mU·kg⁻¹·min⁻¹ of insulin or an infusion of saline, respectively, was begun at a rate of 1.375 µL/min at t = -90 min. Both experiments were conducted in the same environment (e.g. dimensions of the experimental chamber and catheter connections), and thus data from the saline-infusion experiments serve as basal controls for clamp experiments. Euglycemia was maintained during clamps by frequent measurement of arterial blood glucose (~5 µl; HemoCue, Mission Viejo, CA) and infusing 50% dextrose as necessary. Mice received saline-washed red blood cells from a donor mouse as needed to prevent a fall in hematocrit of >5%. At t = 0 min, arterial blood (150 µL) was obtained and processed for baseline analyses. At t = 5 min, a 12 µCi bolus of 2-deoxy[³H]glucose ([2-³H]DG) was administered to determine R_g, an index of tissue-specific glucose uptake. At t = 7, 10, 15, 20 min, arterial blood (~50 µL) was sampled to determine glucose and [2-³H]DG. At t = 30 min, a final arterial sample (170 µL) was drawn and processed the same as the baseline blood sample, but with added blood removal for determination of plasma [2-³H]DG. Mice were anesthetized with a

pentobarbital infusion. The gastrocnemius and heart muscles were excised, frozen in liquid nitrogen, and stored at -70 C.

Exercise and sedentary control experiments. Experiments were performed on 5 h fasted mice as previously described (4; 13; 14). Approximately 1 h prior to an experiment, mice were placed on a treadmill to acclimate them to the changed environment. At $t = 0$ min, a baseline arterial blood sample (150 μ L) was drawn for the measurement of glucose, hematocrit, insulin and, non-esterified fatty acids (NEFAs). To prevent a fall in hematocrit, the remaining erythrocytes were washed once with 0.9% saline containing 10 U/mL of heparin and reinfused. The mice either remained sedentary or ran on a treadmill to 30 min at 16.7 m/min with a 0% grade ($n = 9-11$ for each experimental group and genotype). Both experiments were conducted in the same environment (e.g. dimensions of the experimental chamber on the treadmill and catheter connections), and thus data from sedentary experiments serve as basal controls for exercise experiments. At $t = 5$ min, a 12 μ Ci bolus of [2-³H]deoxyglucose ([2-³H]DG; New England Nuclear, Boston, MA) was administered to determine R_g . At $t = 10, 15,$ and 20 min, ~ 50 μ L of arterial blood was sampled to determine glucose and [2-³H]DG. At $t = 30$ min, a 150 μ L arterial blood was withdrawn in order to determine glucose, hematocrit, insulin, [2-³H]DG, and NEFAs, and mice were anesthetized with an arterial pentobarbital infusion. The gastrocnemius and heart muscles were excised, frozen in liquid nitrogen, and stored at -70 C.

Processing of plasma and muscle samples. Immunoreactive insulin was assayed with a double antibody method (15). NEFAs were measured spectrophotometrically by an enzymatic assay (Wako NEFA C kit, Wako Chemicals Inc., Richmond, VA). Following deproteinization with Ba(OH)₂ (0.3 N) and ZnSO₄ (0.3 N),

plasma [2-³H]DG radioactivity was determined by liquid scintillation counting.

Muscle was homogenized in 0.5% perchloric acid. Homogenates were centrifuged and neutralized with KOH. One aliquot was counted directly to determine [2-³H]DG and [2-³H]DGP radioactivity. A second aliquot was treated with Ba(OH)₂ and ZnSO₄ to remove [2-³H]DGP and any tracer incorporated into glycogen. Radioactivity in [2-³H]DG was then determined. [2-³H]DGP was the difference between the two aliquots. Muscle glycogen was measured according to the method of Chan and Exton (16).

Capillary density was determined in 5 μ m sections of paraffin-embedded gastrocnemius muscles following immunohistochemical detection of CD-31 (platelet endothelial cell adhesion molecule-1, Pecam-1) in endothelial cells. Endogenous peroxidase was quenched with 0.03% hydrogen peroxide and samples were treated with diluted rabbit serum prior to primary antibody addition. Slides were incubated with goat anti-CD-31/Pecam-1 (1: 400, Santa Cruz Biotechnology) for 45 min. The Vectastain ABC Elite (Vector Laboratories, Inc.) System and DAB+ (DakoCytomation) was used to produce visible staining. Slides were lightly counterstained with Mayer's haematoxylin, dehydrated and coverslipped. Capillaries in three visible fields were counted and averaged for each muscle.

Endurance capacity. In separate groups of animals, exercise endurance was measured as previously described (6). Briefly, mice were placed in an enclosed treadmill (Columbus Instruments) and allowed to acclimate to their surroundings for 45 min. Mice were then run at 20 m/min until they were no longer able to keep pace with the treadmill. Mice were encouraged to run as long as possible with the use of an electric grid placed at the end of the treadmill (1.5 mA, 200 ms pulses, 4 Hz). Mice were

defined as exhausted if they remained on the shock grid for five continuous seconds.

Calculations and statistical analysis.

The Insulin Sensitivity Index (ISI) was calculated by dividing the average glucose infusion rate during the last 30 min of an insulin clamp by the insulin concentration over the same time interval. R_g was calculated as previously described (4; 14; 17). To ascertain the effects of dietary and/or genetic manipulations, R_g data are expressed relative to chow-fed, WT mice. Gastrocnemius R_g for chow-fed, WT mice during saline, insulin clamp, sedentary, and exercise experiments were 1.1 ± 0.1 , 6.6 ± 0.8 , 2.6 ± 0.5 , and 24.4 ± 4.7 $\mu\text{mol}/100\text{g}/\text{min}$, respectively. Heart R_g for chow-fed, WT mice during saline, insulin clamp, sedentary, and exercise experiments were 29.9 ± 11.5 , 147.0 ± 12.6 , 23.3 ± 6.3 , and 39.8 ± 8.3 $\mu\text{mol}/100\text{g}/\text{min}$, respectively. Data are presented as means \pm SEM. Differences between groups were determined by ANOVA or repeated-measures ANOVA for time course data followed by a Tukey's post hoc test. The significance level was set at $P < 0.05$.

RESULTS

Baseline characteristics of mouse models. HK II protein was decreased in HK^{+/-} mice by 50% in the gastrocnemius and heart muscles compared to WT mice (**Figure 1**). While HF feeding did not alter gastrocnemius HK II content, it led to a reduction in heart HK II content albeit only in WT mice. Neither a partial HK II knockout nor high fat feeding altered total GLUT4 content.

Baseline characteristics in 5 h fasted WT and HK^{+/-} mice fed either chow or HF diet are shown in **Figure 2**. Partial HK II knockout did not alter body weight or blood glucose concentration in chow fed mice; it did, however, increase plasma insulin concentrations and decrease plasma NEFA concentrations.

HF feeding increased body weight and arterial glucose and insulin concentrations in both genotypes. HF feeding did not alter NEFAs in either genotype. While HK II partial knockout did not exacerbate the hyperglycemia created by HF feeding, HF feeding and partial HK II knockout together resulted in a synergistic increase in fasting insulin.

Partial HK II knockout increased systolic blood pressure and fractional shortening without alterations in heart rate or left ventricular mass (**Table 1**). HF feeding increased left ventricular mass, indicating cardiac hypertrophy, but did not alter heart rate, systolic blood pressure, or fractional shortening. Neither partial HK II knockout nor HF feeding altered capillary density in the gastrocnemius muscle (467 ± 72 , 418 ± 34 , 574 ± 125 , and 516 ± 70 capillaries/ mm^2 for WT-Chow, WT-HF, HK^{+/-}-Chow, and HK^{+/-}-HF, respectively).

Arterial blood glucose during the last 30 min of the saline infusion control for insulin clamps or 30 min sedentary control period for exercise studies are shown in **Figures 3A** and **3C**. There were no differences in blood glucose between groups.

Insulin-stimulated metabolism.

Arterial blood glucose was not different between groups during the saline-infusion control (basal) or hyperinsulinemic-euglycemic clamp experiments (**Figure 3A** and **3B**). The differences in insulin concentration observed in the basal condition remained during the insulin clamp (**Figure 4A**). During the insulin clamp, NEFAs were reduced by ~50% compared to saline-infusion in all groups (0.7 ± 0.1 , 0.8 ± 0.2 , 0.5 ± 0.1 , and 0.6 ± 0.1 mM for WT-Chow, WT-HF, HK^{+/-}-Chow, and HK^{+/-}-HF, respectively). The glucose infusion rate (GIR) was reduced in chow-fed, HK^{+/-} mice (**Figure 4B**). HF feeding in WT mice decreased GIR to the same extent as rates observed in chow-fed HK^{+/-} mice. Further, GIR was lower in HF-

fed, HK^{+/-} mice compared to their WT littermates also fed a HF diet. Because of the differences in the plasma insulin concentrations during the clamp, ISI was calculated. As with GIR, the ISI was lower in chow-fed, HK^{+/-} mice compared to their WT littermates, and HF feeding in WT mice decreased ISI to a comparable level observed in chow-fed HK^{+/-} mice (**Figure 4C**). The combination of HF feeding and partial HK II knockout dramatically decreased ISI compared to all other groups.

During the saline infusion, gastrocnemius R_g was not different between groups (**Figure 5A**). A partial HK II knockout decreased cardiac R_g during the saline infusion (**Figure 6A**). Neither a partial HK II knockout nor HF feeding significantly decreased insulin-stimulated R_g in gastrocnemius (**Figure 5B**). However, the combination of genetic and diet perturbations significantly decreased skeletal muscle insulin-stimulated R_g. In contrast, cardiac insulin-stimulated R_g was decreased by either partial HK II knockout or HF feeding alone, and this decrease was exacerbated by the combination of the two (**Figure 6B**). The attenuation of R_g was marked in HF fed mice with partial HK II knockout despite higher insulin levels.

Gastrocnemius glycogen was generally lower in mice with a partial HK II knockout, whereas cardiac glycogen was higher in mice with a partial HK II knockout (**Table 2**). Cardiac, but not gastrocnemius, glycogen increased dramatically during the insulin clamp.

Exercise metabolism and endurance capacity. Arterial blood glucose did not significantly change throughout the 30 min of exercise in chow fed WT (**Figure 3D**). HF feeding, a partial HK II knockout, and both together resulted in exercise-induced hyperglycemia such that by 15 min of exercise glucose was greater than chow fed WT mice. Despite the exercise-induced

hyperglycemia, insulin concentrations fell in chow fed HK^{+/-} and HF fed mice of both genotypes (data not shown). However, HF fed mice of either genotype still had higher insulin concentrations. While plasma NEFAs remained stable throughout exercise in chow fed WT mice, they fell in the other groups (data not shown).

Sedentary R_g was unaltered by diet or genotype in gastrocnemius but was elevated by partial HK II knockout in cardiac muscle (**Figure 5C** and **6C**). Exercise-stimulated R_g was blunted by HF feeding in gastrocnemius of WT mice (**Figures 5D**). A partial HK II knockout also severely impaired gastrocnemius R_g in chow fed exercising mice. In fact, it was reduced to the same rate as that seen in HF fed WT mice. HF feeding did not further impair gastrocnemius R_g in exercising HK^{+/-} mice. HF feeding dramatically increased cardiac exercise-stimulated R_g (**Figure 6D**). However, this diet-induced increase was severely dampened in mice with partial HK II knockout. HF feeding, partial HK II knockout, and both together decreased endurance capacity, as measured by time to exhaustion (**Figure 7**). Overall, the effects on endurance time closely paralleled those of exercise-stimulated R_g in gastrocnemius.

Gastrocnemius glycogen was lower in all mice following exercise compared to sedentary mice with the exception of WT, HF-fed mice (**Table 2**). As before, gastrocnemius glycogen was generally lower in mice with a partial HK II knockout but this observation did not hold true with cardiac glycogen.

DISCUSSION

We previously reported that mice with a 50% reduction in HK II content have a phenotype at rest that is largely indistinguishable from WT littermates, but exhibit an impaired exercise-induced increase in MGU in oxidative muscles of mice on a

mixed genetic background (14). In the present study we backcrossed mice with the disrupted HK II gene onto the C57BL/6J background to study the effect of this genetic manipulation in combination with HF feeding. The HF fed C57BL/6J mice are obese and insulin resistant compared to chow fed mice of the same strain (18; 19). Here we show that a quantitative genetic reduction of HK II predisposes mice to a greater insulin resistance due to HF diet. A 50% HK II knockout and HF feeding independently resulted in modest insulin resistances that were most evident by increased 5 h fasted insulin concentrations and a diminished rate of glucose disposal during the insulin clamp using a physiological insulin dose. However, when mice with half the normal complement of HK II were fed a HF diet, fasting insulin was fivefold greater than WT mice fed chow. Moreover, insulin resistance became marked as the combined effects of reduced HK II and HF feeding on whole body glucose disposal, insulin sensitivity, and heart and skeletal muscle R_g responses were additive. The additive nature of this interaction in conjunction with an effect of HF feeding on HK II protein level is consistent with the notion that muscle glucose delivery and/or transport, and not glucose phosphorylation, is/are the major causes of dietary insulin resistance in the mouse. There are, however, two alternate interpretations. One cannot strictly rule out that HF feeding may influence the compartmentation of HK during the insulin clamp. If this does occur, the additive effect of diet and genotype on glucose uptake could, in fact, result due to further impairment of glucose phosphorylation. It is also possible that HF diet influences the availability of allosteric regulators of HK II leading to further impairment of glucose phosphorylation.

The results presented here build upon our previous findings demonstrating that a genetic manipulation which resulted in

overexpression of HK II did not improve insulin action in HF fed mice (4). HF feeding reduced HK II protein content in the heart but not gastrocnemius. These studies provide further support that a functional deficit in glucose phosphorylation capacity is present with HF feeding even in the presence of normal HK II protein content. This may be due to the increased acyl-CoA (20) or HK II compartmentation (21; 22).

When the mice performed moderate exercise, plasma insulin and NEFA concentrations generally declined. In contrast, arterial blood glucose remained constant in chow-fed WT mice, but rose at the onset of exercise and reached a plateau by 15 min of the 30 min protocol in mice fed a HF diet with or without a partial HK II deletion such that blood glucose was greater in these groups. During exercise, the massive hyperemia (23; 24) and translocation of GLUT4 from the cytoplasm to sarcolemma (25-27) leads to a robust increase in MGU. The result of these two exercise-stimulated events is that glucose phosphorylation capacity becomes increasingly important in determining MGU (10; 13; 14; 28). Despite the relative hyperglycemia and hyperinsulinemia in HK^{+/-} mice fed chow and HF, the exercise induced increase in gastrocnemius R_g was reduced by 50% of the response seen in WT chow. In contrast to insulin stimulation, the combination of HF feeding and a 50% partial HK II knockout mice had no greater effect than either variable alone. This suggests that partial HK II knockout and HF feeding impair exercise-stimulated R_g through a common mechanism (i.e. glucose phosphorylation). This result is again consistent with a previous study where we showed that HK II overexpression in muscle virtually corrects the reduction in exercise-stimulated R_g seen with HF feeding (4).

HF feeding or a partial HK II knockout both independently impaired

endurance capacity. The combination of the two manipulations did not further reduce endurance. Interestingly, the premature fatigue was closely correlated with exercise-stimulated MGU, and thus endurance capacity and MGU may be linked in the mouse. We previously reported that skeletal muscle HK II content, exercise-stimulated MGU, and exercise endurance capacity are closely correlated (6). Here we show that HF feeding, which impairs exercise-stimulated MGU by 50%, is associated with a 40% reduction in endurance time in the mouse. Mice with a partial HK II knockout also have slightly less skeletal muscle glycogen than their WT littermates and this might contribute to their premature fatigue. In a similar manner, HF-fed WT mice are slightly resistant to skeletal muscle glycogenolysis and thus this inability to efficiently utilize glycogen as a fuel for contractions might promote fatigue. Therefore, the combination of impairments in MGU and limitations in glycogen metabolism created by a partial HK II knockout and/or HF feeding likely lead to accelerated fatigue.

We have previously shown that glucose phosphorylation is a critical determinant of insulin-stimulated MGU (4; 29; 30) just as it is with exercise-stimulated MGU (13; 31). Thus we hypothesized that decreasing HK II protein would create a monogenic form of insulin resistance. This was in fact the case as HK^{+/-} mice had fasting hyperinsulinemia and decreased GIR during a hyperinsulinemic-euglycemic clamp, independent of dietary conditions. Interestingly, the form of insulin resistance created by a partial HK II knockout was not manifested by decreases in skeletal muscle R_g during insulin stimulation. It was, however, associated with decreases in cardiac R_g, suggesting that the heart might be an important tissue to consider when dissecting out the cause of whole body insulin resistance. Strategies to increase muscle HK

activity, such as exercise training, may be an important component of combating dietary-induced insulin resistance as decreases in glucose phosphorylation capacity might worsen the prognosis of patients with metabolic syndrome.

HF feeding provoked a cardiovascular phenotype. Glucose uptake was increased both during rest and exercise in hypertrophic hearts of HF fed mice, despite a 50% reduction in cardiac HK II content. The rise in exercise-stimulated glucose uptake was, however, dampened in mice lacking one HK II allele. These results suggest that the control of glucose uptake exerted by glucose phosphorylation during exercise is different between the heart and skeletal muscle. In contrast to exercise, insulin-stimulated glucose uptake was lower in hearts of HF fed mice compared to WT littermates. Further, a loss of one HK II allele also led to a decrease in insulin-stimulated glucose uptake. These results closely match the qualitative decreases in cardiac HK II content created by HF feeding or a partial HK II knockout. Unlike exercise, insulin stimulation leads to a dramatic fall in the heart's preferred substrate, NEFAs, thereby forcing the heart to utilize glucose as a substrate. The end result is that the lower glucose phosphorylation capacity in HF fed or HK^{+/-} mice limits cardiac glucose uptake.

High fat feeding and commensurate weight gain leads to an increase in cardiac mass. It was of interest to determine if a reduction in the heart's ability to phosphorylate glucose resulting in decreased cardiac glucose uptake would limit cardiac remodeling following HF feeding. The HF feeding maneuver employed here increased left ventricular mass but did not alter heart rate or systolic blood pressure. Interestingly, HK^{+/-} mice had elevated systolic blood pressure compared to WT mice, but this degree of hypertension did not lead to compensatory hypertrophy, suggesting that a

reduction in glucose phosphorylation capacity might limit cardiac remodeling. Further, the reduction in HK II content altered cardiac function as evidenced by increased fractional shortening.

In summary, to ascertain the significance of the glucose phosphorylation barrier to diet-induced insulin resistance, mice absent one HK II allele were fed chow or HF diets. HF feeding in WT mice impaired the insulin- and exercise-stimulated increase in glucose kinetics. HF feeding in mice with a partial HK II deletion did not further impair exercise-stimulated MGU, but resulted in a greatly exaggerated state of insulin resistance. The latter was evident at both the muscle and whole body levels. These results demonstrate that glucose phosphorylation is an important barrier to MGU during exercise in HF fed mice. Consistent with earlier studies in HK II overexpressing mice, diet-induced insulin resistance does not, on the other hand, appear to be due to an increased glucose phosphorylation barrier since the effects of a reduction in HK II adds to the effects of HF

feeding. These data show the impact of a quantitative reduction (50%) in a protein involved in glucose phosphorylation contributes to the insulin resistant state resulting from an environmental factor (i.e. diet).

ACKNOWLEDGEMENTS

We thank Wanda Snead of the Vanderbilt Mouse Metabolic Phenotyping Center (MMPC) Hormone Assay Core for performing the insulin assays and Carlo Malabanan of the MMPC Metabolic Pathophysiology Core for excellent technical assistance. We thank Gemin Ni and ZhiZhang Wang of the MMPC Cardiovascular Pathophysiology Core for performance of echocardiography and blood pressure measurements. We greatly appreciate the assistance of Dr. Lillian Nanney and Kelly Parman of the MMPC Immunohistochemistry Core for measurement of CD31. This work was supported by National Institute of Diabetes and Digestive and Kidney Diseases Grants R01 DK-54902 and U24 DK-59637 (to DHW).

REFERENCES

1. Kim CH, Youn JH, Park JY, Hong SK, Park KS, Park SW, Suh KI, Lee KU: Effects of high-fat diet and exercise training on intracellular glucose metabolism in rats. *Am J Physiol Endocrinol Metab* 278:E977-984, 2000
2. Braithwaite SS, Palazuk B, Colca JR, Edwards CW, 3rd, Hofmann C: Reduced expression of hexokinase II in insulin-resistant diabetes. *Diabetes* 44:43-48, 1995
3. Fueger PT: Glucose phosphorylation as a barrier to muscle glucose uptake. *Clin Exp Pharmacol Physiol* 32:314-318, 2005
4. Fueger PT, Bracy DP, Malabanan CM, Pencek RR, Granner DK, Wasserman DH: Hexokinase II overexpression improves exercise-stimulated but not insulin-stimulated muscle glucose uptake in high-fat-fed C57BL/6J mice. *Diabetes* 53:306-314, 2004
5. Heikkinen S, Pietila M, Halmekyto M, Suppola S, Pirinen E, Deeb SS, Janne J, Laakso M: Hexokinase II-deficient mice. Prenatal death of homozygotes without disturbances in glucose tolerance in heterozygotes. *J Biol Chem* 274:22517-22523, 1999
6. Fueger PT, Shearer J, Krueger TM, Posey KA, Bracy DP, Heikkinen S, Laakso M, Rottman JN, Wasserman DH: Hexokinase II protein content is a determinant of exercise endurance capacity in the mouse. *J Physiol* 566:533-541, 2005
7. Rottman JN, Ni G, Khoo M, Wang Z, Zhang W, Anderson ME, Madu EC: Temporal changes in ventricular function assessed echocardiographically in conscious and anesthetized mice. *J Am Soc Echocardiogr* 16:1150-1157, 2003
8. Exil VJ, Roberts RL, Sims H, McLaughlin JE, Malkin RA, Gardner CD, Ni G, Rottman JN, Strauss AW: Very-long-chain acyl-coenzyme a dehydrogenase deficiency in mice. *Circ Res* 93:448-455, 2003
9. Weisberg AD, Albornoz F, Griffin JP, Crandall DL, Elokda H, Fogo AB, Vaughan DE, Brown NJ: Pharmacological inhibition and genetic deficiency of plasminogen activator inhibitor-1 attenuates angiotensin II/salt-induced aortic remodeling. *Arterioscler Thromb Vasc Biol* 25:365-371, 2005
10. Halseth AE, Bracy DP, Wasserman DH: Overexpression of hexokinase II increases insulin and exercise-stimulated muscle glucose uptake in vivo. *Am J Physiol* 276:E70-77, 1999
11. Niswender KD, Shiota M, Postic C, Cherrington AD, Magnuson MA: Effects of increased glucokinase gene copy number on glucose homeostasis and hepatic glucose metabolism. *J Biol Chem* 272:22570-22575, 1997
12. Ayala JE, Bracy DP, McGuinness OP, Wasserman DH: Considerations in the design of hyperinsulinemic-euglycemic clamps in the conscious mouse. *Diabetes* 55:390-397, 2006
13. Fueger PT, Bracy DP, Malabanan CM, Pencek RR, Wasserman DH: Distributed control of glucose uptake by working muscles of conscious mice: roles of transport and phosphorylation. *Am J Physiol Endocrinol Metab* 286:E77-84, 2004
14. Fueger PT, Heikkinen S, Bracy DP, Malabanan CM, Pencek RR, Laakso M, Wasserman DH: Hexokinase II partial knockout impairs exercise-stimulated glucose uptake in oxidative muscles of mice. *Am J Physiol Endocrinol Metab* 285:E958-963, 2003
15. Morgan CR, Lazarow A: Immunoassay of pancreatic and plasma insulin following alloxan injection of rats. *Diabetes* 14:669-671, 1965
16. Chan TM, Exton JH: A rapid method for the determination of glycogen content and radioactivity in small quantities of tissue or isolated hepatocytes. *Anal Biochem* 71:96-105, 1976

17. Kraegen EW, James DE, Jenkins AB, Chisholm DJ: Dose-response curves for in vivo insulin sensitivity in individual tissues in rats. *Am J Physiol* 248:E353-362, 1985
18. Burcelin R, Crivelli V, Dacosta A, Roy-Tirelli A, Thorens B: Heterogeneous metabolic adaptation of C57BL/6J mice to high-fat diet. *Am J Physiol Endocrinol Metab* 282:E834-842, 2002
19. Surwit RS, Kuhn CM, Cochrane C, McCubbin JA, Feinglos MN: Diet-induced type II diabetes in C57BL/6J mice. *Diabetes* 37:1163-1167, 1988
20. Thompson AL, Cooney GJ: Acyl-CoA inhibition of hexokinase in rat and human skeletal muscle is a potential mechanism of lipid-induced insulin resistance. *Diabetes* 49:1761-1765, 2000
21. Russell RR, 3rd, Mrus JM, Mommessin JI, Taegtmeier H: Compartmentation of hexokinase in rat heart. A critical factor for tracer kinetic analysis of myocardial glucose metabolism. *J Clin Invest* 90:1972-1977, 1992
22. Whitesell RR, Ardehali H, Printz RL, Beechem JM, Knobel SM, Piston DW, Granner DK, Van Der Meer W, Perriott LM, May JM: Control of glucose phosphorylation in L6 myotubes by compartmentalization, hexokinase, and glucose transport. *Biochem J* 370:47-56, 2003
23. Honig CR, Odoroff CL, Frierson JL: Capillary recruitment in exercise: rate, extent, uniformity, and relation to blood flow. *Am J Physiol* 238:H31-42, 1980
24. Palm T, Nielsen SL, Lassen NA: Vascular recruitment in forearm muscles during exercise. *Clin Physiol* 3:445-451, 1983
25. Ploug T, Galbo H, Ohkuwa T, Tranum-Jensen J, Vinten J: Kinetics of glucose transport in rat skeletal muscle membrane vesicles: effects of insulin and contractions. *Am J Physiol* 262:E700-711, 1992
26. Brozinick JT, Jr., Etgen GJ, Jr., Yaspelkis BB, 3rd, Ivy JL: The effects of muscle contraction and insulin on glucose-transporter translocation in rat skeletal muscle. *Biochem J* 297:539-545, 1994
27. Etgen GJ, Jr., Memon AR, Thompson GA, Jr., Ivy JL: Insulin- and contraction-stimulated translocation of GTP-binding proteins and GLUT4 protein in skeletal muscle. *J Biol Chem* 268:20164-20169, 1993
28. Halseth AE, Bracy DP, Wasserman DH: Limitations to exercise- and maximal insulin-stimulated muscle glucose uptake. *J Appl Physiol* 85:2305-2313, 1998
29. Fueger PT, Hess HS, Bracy DP, Pencek RR, Posey KA, Charron MJ, Wasserman DH: Regulation of insulin-stimulated muscle glucose uptake in the conscious mouse: role of glucose transport is dependent on glucose phosphorylation capacity. *Endocrinology* 145:4912-4916, 2004
30. Fueger PT, Shearer J, Bracy DP, Posey KA, Pencek RR, McGuinness OP, Wasserman DH: Control of muscle glucose uptake: test of the rate-limiting step paradigm in conscious, unrestrained mice. *J Physiol* 562:925-935, 2005
31. Fueger PT, Hess HS, Posey KA, Bracy DP, Pencek RR, Charron MJ, Wasserman DH: Control of exercise-stimulated muscle glucose uptake by GLUT4 is dependent on glucose phosphorylation capacity in the conscious mouse. *J Biol Chem* 279:50956-50961, 2004

Table 1. Cardiovascular parameters of conscious C57BL/6J mice.

	WT		HK ^{+/-}	
	Chow	High Fat	Chow	High Fat
HR, <i>bpm</i>	694 ± 12	711 ± 7	700 ± 5	695 ± 18
SBP, <i>mmHg</i>	109 ± 4	113 ± 5	120 ± 3*	115 ± 4
LVmass, <i>mg</i>	92 ± 5	109 ± 5 [†]	97 ± 4	115 ± 8 [†]
FS, %	54 ± 1	53 ± 2	57 ± 1*	53 ± 2
IVSd, <i>mm</i>	0.91 ± 0.01	0.98 ± 0.02 [†]	0.93 ± 0.01	0.97 ± 0.02 [†]
LVIDd, <i>mm</i>	3.12 ± 0.07	3.19 ± 0.06	3.02 ± 0.05	3.16 ± 0.10
LVPWd, <i>mm</i>	0.90 ± 0.02	1.04 ± 0.02 [†]	0.97 ± 0.03*	1.01 ± 0.03
IVSs, <i>mm</i>	1.69 ± 0.03	1.70 ± 0.04	1.78 ± 0.04	1.73 ± 0.05
LVIDs, <i>mm</i>	1.44 ± 0.06	1.52 ± 0.07	1.45 ± 0.04	1.57 ± 0.05
LVPWs, <i>mm</i>	1.34 ± 0.04	1.52 ± 0.03 [†]	1.45 ± 0.04*	1.57 ± 0.05 [†]

Data are means ± s.e.m. *n* = 8–10 per group. HR, heart rate; SBP, systolic blood pressure; LV_{mass}, left ventricular mass; FS, fractional shortening, IVSd, interventricular septal thickness in diastole; LVIDd, LV end-diastolic dimension; LVPWd, LV posterior wall thickness in diastole; IVSs, interventricular septal thickness in systole; LVIDs, LV end-systolic dimension; LVPWs, LV posterior wall thickness in systole. *, *p* < 0.05 vs. WT; [†], *p* < 0.05 vs. Chow.

Table 2. Muscle glycogen following physiological conditions in C57BL/6J mice.

	WT		HK ^{+/-}	
	Chow	High Fat	Chow	High Fat
Gastrocnemius glycogen, mg/g				
Saline infusion	3.3 ± 0.2	3.9 ± 0.3 [†]	2.1 ± 0.3 [*]	2.0 ± 0.3 [*]
Insulin clamp	3.9 ± 0.2 [‡]	3.8 ± 0.3	2.9 ± 0.4 [*]	2.5 ± 0.3 [*]
Sedentary	2.8 ± 0.2	3.0 ± 0.3	1.8 ± 0.2 [*]	2.6 ± 0.2
Exercise	1.9 ± 0.2 [‡]	2.6 ± 0.3 [†]	1.2 ± 0.2 ^{*,‡}	1.5 ± 0.2 ^{*,‡}
Heart glycogen, mg/g				
Saline infusion	0.6 ± 0.1	0.8 ± 0.2	1.1 ± 0.1 [*]	2.0 ± 0.4 ^{*,‡}
Insulin clamp	2.0 ± 0.4 [‡]	2.0 ± 0.3 [‡]	2.5 ± 0.4 [‡]	2.4 ± 0.3
Sedentary	0.9 ± 0.3	0.6 ± 0.1	0.8 ± 0.2	1.3 ± 0.2 ^{*,†}
Exercise	0.3 ± 0.1 [‡]	0.4 ± 0.1	0.7 ± 0.1 [*]	0.9 ± 0.2 ^{†,‡}

Data are means ± s.e.m. $n = 8-10$ per group. ^{*}, $p < 0.05$ vs. WT; [†], $p < 0.05$ vs. Chow; [‡], $p < 0.05$ vs. control (i.e., saline infusion for insulin clamp and sedentary for exercise).

FIGURE LEGENDS

Figure 1. Total muscle HK II and GLUT4 content. Immunoblotting was performed to measure total HK II (*B, D*) and GLUT4 (*C, E*) protein content in the gastrocnemius (*B, C*) and cardiac (*D, E*) muscles of wild type mice (*WT*) or partial HK II knockout ($HK^{+/-}$) mice fed either chow or a HF diet. Glyceraldehyde-3-phosphate dehydrogenase (GAPDH) serves as an internal control. Representative blots for the gastrocnemius are shown in *A*. Data are means \pm SEM for 4-8 mice/group. *, $p < 0.05$ vs. *WT-Chow*.

Figure 2. Metabolic parameters of chronically catheterized, 5 h-fasted C57/BL6J mice. Wild type (*WT*) or partial HK II knockout ($HK^{+/-}$) mice fed either chow or a HF diet had catheters surgically implanted in a carotid artery and jugular vein and were allowed to recover for at least 5 d. Body weight was measured (*A*) and mice were fasted for 5 h. Arterial blood was obtained and measured for blood glucose (*B*), insulin (*C*), and non-esterified fatty acids (NEFAs, *D*). Data are means \pm SEM for 9-12 mice/group. *, $p < 0.05$ vs. Chow; †, $p < 0.05$ vs. *WT*.

Figure 3. Arterial blood glucose concentration during experimental conditions. Wild type (*WT*) or partial HK II knockout ($HK^{+/-}$) mice fed either chow or a HF diet had catheters surgically implanted in a carotid artery and jugular vein and were allowed to recover for at least 5 d. Arterial blood glucose was measured during the last 30 min of a 120 min saline infusion (*A*) or hyperinsulinemic-euglycemic clamp (Insulin clamp, *B*) or during a 30 min sedentary (*C*) or exercise (*D*) period. Data are means \pm SEM for 9-12 mice/group. *, $p < 0.05$ vs. *WT HF*; †, $p < 0.05$ vs. all other groups; ‡, $p < 0.05$ vs. all other groups.

Figure 4. Insulin concentration, glucose infusion rate, and insulin sensitivity index during a hyperinsulinemic-euglycemic clamp. Wild type (*WT*) or partial HK II knockout ($HK^{+/-}$) mice fed either chow or a HF diet had catheters surgically implanted in a carotid artery and jugular vein and were allowed to recover for at least 5 d. After a 5 h fast, insulin was infused at 4 mU/kg/min and arterial blood glucose was maintained by infusing glucose into a jugular vein. Average arterial plasma insulin concentrations during the experiment are shown (*A*). The average glucose infusion rates over the final 30 min of the 120 min hyperinsulinemic-euglycemic clamp (*B*) and insulin sensitivity index (*C*) are reported. Data are means \pm SEM for 9-12 mice/group. *, $p < 0.05$ vs. *WT-Chow*; †, $p < 0.05$ vs. all other groups.

Figure 5. Glucose uptake into the gastrocnemius muscle during insulin-stimulation and exercise. Gastrocnemius R_g , an index of muscle glucose uptake, was measured in conscious, freely moving wild type (*WT*) or partial HK II knockout ($HK^{+/-}$) mice fed either chow or a HF diet during: (*A*) a control saline-infusion, (*B*) a hyperinsulinemic-euglycemic clamp (Insulin-stimulated), (*C*) sedentary, and (*D*) exercise. Data are expressed relative to the values obtained for *WT-Chow* mice and are means \pm SEM for 9-12 mice/group. Gastrocnemius R_g for chow-fed, *WT* mice during saline, insulin clamp, sedentary, and exercise experiments were 1.1 ± 0.1 , 6.6 ± 0.8 , 2.6 ± 0.5 , and 24.4 ± 4.7 $\mu\text{mol}/100\text{g}/\text{min}$, respectively. *, $p < 0.05$ vs. *WT-Chow*.

Figure 6. Glucose uptake into the heart during insulin-stimulation and exercise. Heart R_g , an index of muscle glucose uptake, was measured in conscious, freely moving wild type (*WT*) or partial HK II knockout ($HK^{+/-}$) mice fed either chow or a HF diet during: (*A*) a control saline-

infusion, (B) a hyperinsulinemic-euglycemic clamp (Insulin-stimulated), (C) Sedentary, and (D) exercise. Data are expressed relative to the values obtained for WT-Chow mice and are means \pm SEM for 9-12 mice/group. Heart R_g for chow-fed, WT mice during saline, insulin clamp, sedentary, and exercise experiments were 29.9 ± 11.5 , 147.0 ± 12.6 , 23.3 ± 6.3 , and 39.8 ± 8.3 $\mu\text{mol}/100\text{g}/\text{min}$, respectively. *, $p < 0.05$ vs. WT-Chow; †, $p < 0.05$ vs. all other groups.

Figure 7. Endurance capacity is impaired by partial HK II knockout or HF feeding. Wild type (WT) or HK II partial knockout ($HK^{+/-}$) mice fed either chow or a HF diet were run at 20 m/min on a motorized treadmill until exhaustion in order to determine exercise endurance capacity. Data shown are means \pm S.E.M. for 9-12 mice/group. *, $p < 0.05$ vs. WT-Chow.




Figure 1.




Figure 2.




Figure 3.




Figure 4.




Figure 5.




Figure 6.




Figure 7.