

Evidence that Processes Other than Gluconeogenesis may influence the Ratio of Deuterium on the Fifth and Third Carbons of Glucose:
Implications for the use of $^2\text{H}_2\text{O}$ to measure gluconeogenesis in humans

Gerlies Bock^{*}, MD, William C. Schumann⁺ PhD, Rita Basu^{*}, MD, Shawn C. Burgess[#] PhD,
Zheng Yan[#], Visvanathan Chandramouli⁺, PhD, Robert A. Rizza^{*}, MD and
Bernard R. Landau^{a +}, MD, PhD

^{*}Division of Endocrinology, Diabetes, Metabolism & Nutrition,
Mayo Clinic College of Medicine, Rochester, MN, Department of Medicine;

⁺Case Western Reserve University School of Medicine, Cleveland Ohio,
The Advanced Imaging Research Center; [#]University of Texas Southwestern Medical Center,
Dallas, TX

^a deceased

Running title: Transaldolase reaction

Corresponding Author:

Robert A. Rizza MD
Mayo Clinic
200 1st Street SW, Rm 5-194 Joseph
Rochester, MN 55905
E-mail: rizza.robert@mayo.edu

Received for publication 22 May 2007 and accepted in revised form 1 October 2007.

ABSTRACT

Objective: The deuterated water method uses the ratio of deuterium on carbons five and two (C5/C2) or three and two (C3/C2) to estimate the fraction of glucose derived from gluconeogenesis. The current studies determined if C3 and C5 glucose enrichment is influenced by processes other than gluconeogenesis.

Research Design: Six non-diabetic subjects were infused with [3,5-²H₂] glucose and insulin while glucose was clamped at ~5 mmol/L; the C5/C3 ratio was measured in the in UDP glucose pool using NMR and the acetaminophen glucuronide method.

Results: Whereas, the C5/C3 ratio of the infusate was 1.07, the ratio in UDP-glucose was less than 1.0 in all subjects both prior to (0.75±0.07) and during (0.67±0.05) the insulin infusion.

Conclusions: These data indicate that the deuterium on C5 of glucose is lost more rapidly relative to the deuterium on C3. The decrease in the C5/C3 ratio could result from exchange of the lower three carbons of fructose-6-phosphate with unlabeled three carbon precursors via the transaldolase reaction and/or selective retention of the C3 deuterium at the level of triosephosphate isomerase due to a kinetic isotope effect. Following ingestion of ²H₂O, these processes would increase the enrichment of C5 and decrease the enrichment of C3 respectively with the former causing an overestimation of gluconeogenesis using the C2/C5 ratio and the latter an underestimation using the C3/C2 ratio. Future studies will be required to determine whether the impact of these processes on the measurement of gluconeogenesis differs amongst the disease states being evaluated (e.g. diabetes or obesity).

Keywords: Transaldolase; triosephosphate isomerase, gluconeogenesis; endogenous glucose production; nuclear magnetic resonance

Abbreviations: GNG: Gluconeogenesis, GLG: Glycogenolysis, EGP: Endogenous glucose production; NMR: nuclear magnetic resonance; G3P: glyceraldehyde-3 phosphate; DHAP: dihydroxyacetone phosphate

Plasma glucose concentration is determined by the balance between the rate of glucose production and glucose uptake. Glucose released from the liver is derived from either the degradation of glycogen, referred to as glycogenolysis or the synthesis of new glucose from three carbon precursors, referred to as gluconeogenesis. Multiple methods have been used to estimate rates of glycogenolysis and gluconeogenesis in humans with the deuterated water method perhaps being the most widely accepted (1-7). This method is based on the assumption that following ingestion of $^2\text{H}_2\text{O}$, the fifth carbon of glucose (C5) is labeled with deuterium at the level of the triose phosphate isomerase reaction during gluconeogenesis whereas the second carbon of glucose (C2) is labeled with deuterium during both glycogenolysis and gluconeogenesis (1). Therefore the plasma C5/C2 glucose ratio is used to estimate the fraction of glucose derived from gluconeogenesis. Since the third carbon of glucose (C3) and C5 are concurrently labeled during the triose isomerase reaction, the C3/C2 ratio also has been used to assess gluconeogenesis (8). The absolute rate of gluconeogenesis is calculated by multiplying the fraction of glucose derived from gluconeogenesis times the tracer determined rate of endogenous glucose production (9-11). The rate of glycogenolysis is then determined by subtracting the rate of gluconeogenesis from glucose production.

In general, estimates derived with the deuterated water method are concordant with those derived using other methods of estimating gluconeogenesis (2,12). In addition, gluconeogenesis measured with the deuterated water method changes in a biologically plausible manner during perturbations such as fasting or diabetes (1,11,13-15). However, unfortunately there is no "gold standard" against which results derived with this method can be compared.

An alternative approach is to test the assumptions of the method. As noted above, a key assumption is that labeling of the fifth carbon of glucose with deuterium following ingestion of $^2\text{H}_2\text{O}$ only occurs during gluconeogenesis (1). The specific biochemical step in which the C5 hydrogen is exchanged for body water is during the equilibration between glyceraldehyde-3 phosphate and dihydroxyacetone phosphate at the triosephosphate isomerase reaction (1). The C3 hydrogen also is lost during this exchange. Therefore, whenever a triose molecule is converted to glucose, enrichment at either C5 or C3 can, in principle, be used to index gluconeogenesis. However, processes other than gluconeogenesis can influence enrichment of these carbons. For instance, C5 of glucose can become labeled when the lower three carbons of fructose-6-phosphate and/or sedoheptulose exchange with glyceraldehyde 3-phosphate by the transaldolase reaction (figure 1). Since this represents a simple exchange of the bottom 3 carbons of fructose-6-phosphate (C4-C6) with glyceraldehyde-3-phosphate rather than new hexose synthesis, generation of C5 enriched glucose via transaldolase will result in an overestimate of gluconeogenesis (16). Conversely, exchange of the C3 hydrogen with deuterium during the triosephosphate isomerase reaction may not be complete due to a primary kinetic isotope effect (17,18). If so, this would result in a lower than expected enrichment in C3 and therefore an underestimate of gluconeogenesis if the C3/C2 ratio is used. However, the extent (if at all) to which these processes occur in humans, is not known.

The present experiments addressed these questions by infusing $[3,5 \text{ } ^2\text{H}_2]$ glucose labeled in a 1/1 ratio (i.e. C5/C3 glucose ratio equaled 1.0) with deuterium on both the fifth and third carbons in non-diabetic humans prior to and during a hyperinsulinemic

euglycemic clamp. Enrichment in the UDP-glucose pool was measured with the acetaminophen glucuronide method. Since at steady state, UDP-glucose is in equilibrium with the glucose-6-phosphate pool and glucose-6-phosphate is derived from both plasma glucose and the fructose-6-phosphate pools, a C5/C3 ratio in the UDP-glucose pool that differs from that of the infusate indicates that processes in addition to gluconeogenesis influence this ratio.

RESEARCH DESIGN AND METHODS

After approval of the Mayo Institutional Review Board 6 subjects (4 women and 2 men, age 53 ± 5 , BMI 33.7 ± 1.6 , LBM 41.7 ± 3.0) gave written informed consent to participate in the study. Subjects were in good health, stable at weight, did not engage in vigorous exercise, were on no medication at the time of the study and did not have a history of diabetes in first degree relatives. All subjects were interviewed by a dietician and instructed to follow a weight maintaining diet consistent of 55% carbohydrate, 30% fat and 15% protein for a least one week prior to the study. Subjects were admitted to the Mayo General Clinical Research Center the evening before the study, ingested a standard 10 kcal/kg dinner at 1800 then remained fasting thereafter. At 0600 the following morning, an 18-gauge cannula was inserted in a forearm vein for tracer and hormone infusions. A second 18-gauge cannula was inserted in a retrograde fashion in a dorsal hand vein of the opposite arm, and the hand was placed in a heated box ($\sim 55^\circ\text{C}$) to enable sampling of arterialized venous blood. A urinary catheter was placed for urine collection. A primed (fasting glucose in mmol/l divided by 5.5 mmol/l times 12 μCi) continuous (0.12 $\mu\text{Ci}/\text{min}$) infusion of [$3\text{-}^3\text{H}$] glucose (New England Nuclear, Boston, MA) was started at 0700 and continued until the end of the study. In addition, two grams of liquid sugar free acetaminophen was given by mouth in split

doses at 0800 and 0930 to enable measurement of urinary acetaminophen glucuronide specific activity. At 0900, a primed (133 mg in the first four subjects; 62.5 mg in the final two subjects) -continuous (80 mg/hour in the first four subjects; 40 mg/hr in the final two subjects) infusion of [$3,5\text{-}^2\text{H}$] glucose (Omicron, Biochemicals Inc., Indiana) was started to enable the measurement of contribution of the transaldolase reaction to endogenous glucose production. A constant infusion containing somatostatin (60 ng/kg/min), glucagon (0.65 ng/kg/min), growth hormone (3 ng/kg/min) and insulin (0.50 mU/kg/min) was started at 1000 (i.e., time 0) and continued for 4 hours. Dextrose (D₅₀W) containing [$3\text{-}^3\text{H}$] glucose was infused as necessary to maintain plasma glucose concentrations at ~ 5.0 mmol/l as previously described (19). Arterialized venous blood samples were collected for measurements of glucose, hormones and tracer concentrations. Urine was collected from 0900 to 1000, 1000 to 1200 and 1200 to 1400 for analysis of urine acetaminophen glucuronide.

Analytical techniques. Plasma samples were placed on ice, centrifuged at 4°C , separated, and stored at -20°C until assay. Plasma glucose concentration was measured using a glucose oxidase method (Yellow Springs Instrument, Yellow Springs, OH). Plasma insulin concentrations were measured using a chemiluminescence assay with reagents obtained from Beckman (Access Assay; Beckman, Chaska, MN). Body composition was measured using dual-energy X-ray absorptiometry (DPX scanner; Lunar; Madison, WI). Plasma [$3\text{-}^3\text{H}$] glucose specific activity was measured using liquid scintillation counting as previously described (20).

Acetaminophen glucuronide was isolated as previously described (21). Briefly urine was defrosted and concentrated to a

small volume. The residue was dissolved in methanol and the methanol solution was filtered and evaporated to dryness. The residue was dissolved in water, pH raised to 9.0 with 10N NaOH and applied to an ion exchange column (spectragel 1x8 75-150um Spectrum) in its acetate form. The column was washed with water until urea test of fractions was negative and acetaminophen glucuronide was eluted with 5M acetic acid. The acetic acid fractions were evaporated off and the residue dissolved in 0.1M acetate buffer pH 5.0. The acetaminophen glucuronide was converted to monoacetoneglucose (MAG) by a slight modification of the method of Jones et al (22). Acetaminophen glucuronide in acetate buffer was added with glucuronidase (Helix Pomatia) and incubated for 48h at 37 °C. At the end of the incubation Ba(OH₂) and ZnSO₄ were added to the reaction mixture and the supernatant was passed down a column (Dowex 50W) in H⁺ form. The eluent was evaporated to dryness, then stored in a desiccator overnight to give glucuronolactone. The lactone was stirred overnight with anhydrous acetone containing 2% H₂SO₄. Water was added to quench the reaction and pH was raised to 5.0 with 1M Sodium Carbonate. This solution was evaporated off to dryness and extracted three times with tetrahydrofuran (THF) 10 ml each. The combined THF solution was reduced with 1M Lithium Borohydride in THF. The solution was quenched with water, made acidic with acetic acid, and evaporated to dryness. The residue was dissolved in 10 ml methanol and evaporated. This process was repeated twice more. The monoacetone glucose was dissolved in water and put through a 3ml Sep-Pak C-18 cartridge (Waters), washed with water and monoacetoneglucose eluted in 5.0% acetonitrile in water. This fraction evaporated off and dissolved in acetonitrile for NMR.

Analysis of C3 and C5 deuterium enrichments in MAG was performed by ²H NMR as previously described (23). The MAG samples were dissolved in 160 µl of HPLC grade acetonitrile with 5-10 ul of water and transferred to a 3 mm NMR tube. Deuterium NMR spectra of MAG were collected using a 14.1 T Varian INOVA spectrometer and 3 mm broadband probe, tuned to 92 MHz as previously described. A 90 degree pulse was applied and the signal was acquired over 1 s (sweep width = 1000 Hz) with no further delay. Deuterium spectra were collected in blocks of 256 acquisitions without lock followed by a single pulse on the proton channel. The proton position of the solvent peak was ascertained and Z₀ was automatically adjusted as a correction to any field drift since the collection of the last block. The end of acquisition all blocks were summed for the final spectrum. ²H NMR spectra were typically signal-averaged for 1-4 hours at 50 °C. NMR peak areas were fitted using the PC based NMR analysis software ACD (Advanced Chemistry Development Inc, Toronto Canada). The H3 and H5 represent the ²H peak area at the 3 and 5 positions of MAG, respectively. Baseline was chosen as the center of the noise across the span of the spectrum and corrected by the NMR software. Operator error was minimized by allowing the ADC software to line fit the NMR peaks automatically. Repeated measurements of NMR data by a single operator typically yield <5% SD, but is dependent on the signal/noise quality of the spectrum

Statistical analysis. Data in the text and figures are expressed as mean ± SEM. Rates are expressed as per kilogram of fat free mass. Values from -30 to 0 minutes were averaged and considered as basal and values from 210 to 240 minutes were averaged and considered as clamp. Student's paired t- test was used to determine if the C5/C3 ratio was less than

one. A $p < 0.05$ was considered as statistically significant. Values are means \pm SEM.

RESULTS

Plasma glucose, insulin, and C-peptide concentrations (figure 2). Plasma glucose concentrations did not differ prior to and during the clamp (5.1 ± 0.1 vs. 5.1 ± 0.1 mmol/l). Plasma insulin concentrations increased from 35 ± 5 pmol/l before the clamp to 130 ± 13 pmol/l during the clamp. Somatostatin resulted in essentially complete suppression of C-peptide.

Endogenous glucose production and glucose disappearance (figure 3). Endogenous glucose production averaged 17.3 ± 1.0 μ mol/kg/min before the clamp and decreased to 1.5 ± 1.4 μ mol/kg/min by the end of the clamp. On the other hand, glucose disappearance averaged 16.9 ± 0.8 μ mol/kg/min before the clamp and increased to 22.4 ± 2.3 μ mol/kg/min by the end of the clamp.

C5/C3 glucose ratio (figures 4 and 5). A typical spectra of MAG derived from UDP-glucose is shown in figure 4. Peak areas correspond to the relative deuterium enrichments of the two hydrogen positions of glucose. The C5/C3 glucose ratio in the infusate was 1.07 indicating that fifth and third carbons of glucose were equally labeled with deuterium. The C5/C3 ratio in the UDP glucose pool (figure 5) was lower than 1.0 before the insulin infusion in all four subjects in whom it was measured (0.75 ± 0.07) and during the insulin infusion in all six subjects (0.67 ± 0.05) indicating either selective loss of deuterium from the fifth carbon of glucose and/or selective retention of deuterium on the third carbon of glucose

DISCUSSION

The deuterated water method for the measurement of gluconeogenesis relies on the

assumption that there is extensive equilibration between the precursor hydrogens and body water (1,24). Therefore, enrichment of plasma with $^2\text{H}_2\text{O}$ results in the labeling of newly synthesized glucose with deuterium. The ratio of the deuterium on C5 (labeled during gluconeogenesis) to that on C2 (labeled during both gluconeogenesis and glycogenolysis due to equilibration between glucose-6-phosphate and fructose-6-phosphate pools) of glucose provides an estimate of the fraction of plasma glucose derived from gluconeogenesis (1). Since in the absence of an isotope effect, C5 and C3 are equally labeled during the triose phosphate isomerase reaction, in theory, the C3/C2 ratio also can be used to assess the fractional contribution of gluconeogenesis (8). The present study provides evidence that factors other than gluconeogenesis influence enrichment of C5 and/or C3 since the C5/C3 ratio of UDP-glucose was less than 1.0 both before and during infusion of insulin despite the fact the enrichment of the infused C5/C3 glucose equaled 1.0. This means that during equilibration of the fructose-6-phosphate, glucose-6-phosphate and UDP-glucose pools, the deuterium on the fifth carbon of glucose was selectively lost and/or the deuterium on the third carbon of glucose was selectively retained. Both scenarios have implications when the deuterated water method is used to measure gluconeogenesis.

Transaldolase exchange. The transaldolase reaction can exchange the bottom three carbons of C5 labeled fructose-6-phosphate with glyceraldehyde-3-phosphate. Selective loss of C5 deuterium occurs when the lower three carbons of fructose-6-phosphate exchange with unlabeled glyceraldehyde-3-phosphate via the transaldolase reaction (16,25). It is unlikely that the pentosephosphate pathway contributed significantly to our observations because deuterium on C3, but not on C5 is lost when

glucose-6-phosphate traverses the pentose phosphate pathway which would increase rather than decrease the C5/C3 ratio (16). On the other hand, in the absence of an isotope effect (see below) the deuteriums on C3 and C5 are equally lost during equilibration of glyceraldehyde-3-phosphate and dihydroxyacetone at the level of the triose isomerase reaction (16). It is conceivable that glyceraldehyde-3-phosphate can become labeled with deuterium during its equilibration with dihydroxyacetone at the level of the triose isomerase reaction. However, the probability would be very low under the present experimental conditions since $^2\text{H}_2\text{O}$ only was generated during metabolism of C5/C3 glucose which was infused in trace amounts.

We are unaware of prior studies in humans that have measured the effect of the transaldolase reaction on labeling of the fifth carbon of glucose. On the other hand, this reaction is well documented *in vitro* (16,25). Of note, in a parallel study using a similar approach, Jones et al (personal communication) observed a C5/C3 ratio of ~0.77 in 6 healthy subjects following an overnight fast and ingestion of [$\text{U}-^2\text{H}_7$] glucose. This ratio is entirely consistent with the ratios observed in the present experiment of 0.75 in the state and 0.67 during infusion of insulin. While exchange of the lower three carbons of fructose-6-phosphate with unlabeled glyceraldehyde-3-phosphate would result in a decrease in C5 enrichment in the present study, the opposite would have occurred if plasma had been enriched with $^2\text{H}_2\text{O}$. Under those conditions, glyceraldehyde-3-phosphate would be extensively labeled with deuterium during its equilibration with dihydroxyacetone and carbon exchange via the transaldolase reaction would increase C5 glucose enrichment in the absence of an increase in gluconeogenesis.

Triosephosphate isomerase kinetic isotope effect. Selective retention of deuterium on C3 relative to C5 due to a kinetic isotope effect at the level of the triosephosphate isomerase reaction also would reduce the C5/C3 ratio. This effect is well known *in vitro* (17,18,26-28), but has not yet been described *in vivo*. The origin of the kinetic isotope effect lies in the fact that ^2H and ^3H isotopes have 1 and 2 extra neutrons respectively, which doubles and triples their mass compared to the naturally abundant ^1H . If the location of the isotope label is in a position that participates in the transition state of a biochemical reaction, then the heavier mass isotope increases the activation energy for this step and a slower reaction rate is observed for isotopically labeled substrates compared to unlabeled substrates. This effect is particularly important in hydrogen isotope labeled substrates because they have highest degree of mass difference compared to their natural abundant isotope and their bonds are the most likely to be broken and formed in the transition state of biochemical reactions. This appears to be the case in the triosephosphate isomerase reaction. When [$3,5-^2\text{H}_2$]glucose is metabolized in the liver to the level of triosephosphates, [$1-^2\text{H}$]-DHAP and [$3-^2\text{H}$]-G3P are initially formed. In principle, the interconversion of these two metabolites by the triosephosphate isomerase reaction leads to an equivalent loss of ^2H label in both positions. In practice, Knowles and co-workers showed that 77% of label in [$3-^3\text{H}$]-G3P is exchanged with water (27), but only 13% of label in [$1-^3\text{H}$]-DHAP is exchanged with water during the triosephosphate isomerase reaction (17). For deuterium, the kinetic isotope effect at the C1 position of DHAP is 3-fold, but essentially negligible for the C2 position of G3P (26). This represents a substantial primary kinetic isotope effect during the exchange of the C1 hydrogen of DHAP (destined to become glucose C3 hydrogen) but not in the C2 hydrogen of G3P

(destined to become glucose C5 hydrogen). Under the present experimental conditions selective retention of deuterium enrichment in the 3 position compared to the 5 position also would result in a reduction of the C5/C3 ratio.

Implications for the use of $^2\text{H}_2\text{O}$ to measure gluconeogenesis in vivo. Deuterium enrichment of plasma glucose and urinary glucuronide has been measured after administration of deuterated water under a variety of conditions in both human (8,29,30) and animal studies by NMR (15,23,31). While the C3 enrichment typically has not been reported in these studies, it is easily visualized in the NMR spectra shown in the above references. We are unaware of any instance in which glucose C3 and C5 deuterium enrichment become equal to one another following administration of $^2\text{H}_2\text{O}$. In addition, measurement of gluconeogenesis in overnight fasted humans using the C3 enrichment yields a lower estimate of gluconeogenesis than does measurement of C5 enrichment (8). While, this observation is consistent with the reduced C5/C3 ratio observed in the present studies, it provides no insight into the origin of the discrepancy. However, studies in long term fasted rodents (24-hours) are helpful because under these conditions liver glycogen is depleted and which means essentially all of glucose production originates from gluconeogenesis (15,23). Under these conditions, measurement of C5 enrichment indicates that in excess of 95 % of glucose production is derived from gluconeogenesis. In contrast, measurement of C3 enrichment yielded a less likely value of ~75% gluconeogenesis (15,23). Thus, while the C5/C2 ratio approached 1.0 after an extended fast, the C3/C2 ratio did not which is consistent with a primary kinetic isotope effect. It should be noted that the fact that measurement of C5 enrichment during a prolonged fast yields plausible estimates of gluconeogenesis does

not rule out on going transaldolase exchange since under these conditions, C5 enrichment will approach plasma $^2\text{H}_2\text{O}$ enrichment and therefore cannot be further increased by exchange by labeled three carbon precursors. This situation differs from that which occurs under conditions of daily living where gluconeogenesis does not account for 100 % of glucose production. In this circumstance, C5 enrichment is substantially less than plasma $^2\text{H}_2\text{O}$ and therefore can be increased by exchange with labeled three carbon precursors.

Limitations. The present study is subject to limitations. Since enrichment in plasma was low, enrichment of C5/C3 in the UDP-glucose pool was measured by collecting urine at two hour intervals and extracting acetaminophen glucuronide. Since there presumably is a delay between the time glucuronide is formed within the liver and it is excreted in the urine, this method looks back at what occurred during the interval before sampling of the urine. Therefore the enrichment of glucuronide in the liver and urine at any given time may differ. However, C5/C3 ratio remained essentially constant over the five hours of study. This suggests that equilibrium between hepatic and urine glucuronide was reached during the first two hours of the study and remained unchanged throughout the remainder of the study. Perhaps more importantly from the perspective of the present experiments, our conclusions are based on the C5/C3 ratio rather than the degree to which steady state was achieved. This is of interest since endogenous glucose production gradually decreased over the same period of time to values that no longer differed from zero by the end of the study. Consistent with previous studies in animals (32), the progressive fall in endogenous glucose production in the absence of a change in the C5/C3 ratio suggests that insulin inhibited hepatic glucose release by

suppressing glucose-6-phosphatase activity over time without altering the equilibration amongst the fructose-6-phosphate, glucose-6-phosphate and UDP-glucose pools.

The C5/C3 ratio only was measured in the fasting state and during a euglycemic hyperinsulinemic clamp. We therefore do not know if the increase in hepatic glucose uptake that occurs in the presence of combined hyperglycemia and hyperinsulinemia would alter the rate of the transaldolase reaction and/or the rate of the isomerase reaction. Only six subjects were studied. Since the C5/C3 ratio in those subjects ranged from ~0.5 to ~0.9, we do not know if the same degree of variation would be observed in a larger population. However the ratio was less than one in *all* subjects strongly supporting our conclusion that processes other than gluconeogenesis influence this ratio. We only studied obese non-diabetic subjects. We therefore do not know if other metabolic conditions (e.g. obesity or diabetes) influences the C5/C3 ratio. On the other hand, if the C5/C3 ratio remains constant under the conditions in which such comparisons are made, then while enrichment of C5 glucose via the transaldolase reaction would make determination of the absolute rates of gluconeogenesis problematic, it would not preclude comparisons among groups.

CONCLUSIONS

In summary, the present data indicate that processes other than the rate of gluconeogenesis influence the C5/C3 deuterium ratio of urinary glucuronide after infusion of [3,5-²H]glucose in humans. The 25-30 percent reduction of the C5/C3 ratio in UDP-glucose relative to the ratio of 1.0 in the

infused C5/C3 glucose both in the fasting state and during a hyperinsulinemic euglycemic clamp is consistent with transaldolase exchange, a kinetic isotope effect at the level of triosephosphate isomerase or a combination of both. In any case, our results do not obviate the utility of the deuterated water method to estimate gluconeogenesis. While both the ²H₂O and isotopomer methods are affected by transaldolase exchange and/or an isotope effects at the level of the triose phosphate isomerase step, we believe, they remain the best available methods for assessing gluconeogenesis *in vivo*. However, it will be important in future studies to determine whether the processes other than gluconeogenesis that increase labeling C5 or reduce labeling of C3 following ingestion of deuterated water differ amongst the disease states being evaluated (e.g. diabetes or obesity).

ACKNOWLEDGMENTS

This study was supported by U.S. Public Health Service (DK29953, DK14507, UL1RR024150) and a Merck research infrastructure grant. GB was supported by an American Diabetes Association Mentor based fellowship. SCB is the recipient of an American Diabetes Association Junior Faculty Award 1-50-JF-05. NMR facilities are provided by a National Center for Research Resources, RR02584. We wish to thank B. Dicke, G. DeFoster, P. Reich, and B. Norby for technical assistance, M. Davis for assistance with preparation of the manuscript, and the staff of the Mayo Clinical Research Unit for assistance with the studies.

REFERENCES

1. Landau BR, Wahren J, Chandramouli V, Schumann WC, Ekberg K, Kalhan SC: Contributions of gluconeogenesis to glucose production in the fasted state. *Journal of Clinical Investigation* 98:378-385, 1996
2. Edgerton DS, Cardin S, Emshwiller M, Neal D, Chandramouli V, Schumann WC, Landau BR, Rossetti L, Cherrington AD: Small increases in insulin inhibit hepatic glucose production solely caused by an effect on glycogen metabolism. *Diabetes* 50:1872-1882, 2001
3. Kunert O, Stingl H, Rosian E, Krssák M, Bernroider E, Seebacher W, Zangger K, Staehr P, Chandramouli V, Landau BR, Nowotny P, Waldhäusl W, Haslinger E, Roden M: Measurement of fractional whole-body gluconeogenesis in humans from blood samples using ^2H nuclear magnetic resonance spectroscopy. *Diabetes* 52:2475-2482, 2003
4. Neese RA, Schwarz J-M, Faix D, Turner S, Letscher A, Vu D, Hellerstein MK: Gluconeogenesis and intrahepatic triose phosphate flux in response to fasting or substrate loads. *J.Biol.Chem.* 270:14452-14463, 1995
5. Rothman DL, Magnusson I, Katz LD, Shulman RG, Shulman GI: Quantitation of hepatic glycogenolysis and gluconeogenesis in fasting humans with ^{13}C NMR. *Science* 254:573-254, 1991
6. Peroni O, Large V, Beylot M: Measuring gluconeogenesis with [2- ^{13}C] glycerol and mass isotopomer distribution analysis of glucose. *Am.J.Physiol.* 269:E516-E523, 1995
7. Radziuk J, Lee WP: Measurement of gluconeogenesis and mass isotopomer analysis based on [U-(^{13}C)]glucose. *Am J Physiol Endocrinol Metab* 277:E199-E207, 1999
8. Jones JG, Carvalho RA, Sherry AD, Malloy CR: Quantitation of gluconeogenesis by ^2H nuclear magnetic resonance analysis of plasma glucose following ingestion of $^2\text{H}_2\text{O}$. *Anal Biochem* 277:121-126, 2000
9. Van Schaftingen E, Gerin I: The glucose-6-phosphatase system. *Biochem J* 362:513-532, 2002
10. Adkins A, Basu R, Persson M, Dicke B, Shah P, Vella A, Schwenk WF, Rizza R: Higher insulin concentrations are required to suppress gluconeogenesis than glycogenolysis in non-diabetic humans. *Diabetes* 52:2213-2220, 2003
11. Gastaldelli A, Baldi S, Pettiti M, Toschi E, Camastra S, Natali A, Landau BR, Ferrannini E: Influence of obesity and type 2 diabetes on gluconeogenesis and glucose output in humans: a quantitative study. *Diabetes* 49:1367-1373, 2000
12. Kunert O, Stingl H, Rosian E, Krssak M, Bernroider E, Seebacher W, Zangger K, Staehr P, Chandramouli V, Landau BR, Nowotny P, Waldhausl W, Haslinger E, Roden M: Measurement of fractional whole-body gluconeogenesis in humans from blood samples using ^2H nuclear magnetic resonance spectroscopy. *Diabetes* 52:2475-2482, 2003
13. Basu R, Chandramouli V, Dicke B, Landau B, Rizza R: Obesity and type 2 diabetes impair insulin-induced suppression of glycogenolysis as well as gluconeogenesis. *Diabetes* 54:1942-1948, 2005
14. Boden G, Chen X, Stein TP: Gluconeogenesis in moderately and severely hyperglycemic patients with type 2 diabetes mellitus. *Am J Physiol Endocrinol Metab* 280:E23-E30, 2001
15. Burgess SC, Jeffrey FMH, Storey C, Milde A, Hausler N, Merritt ME, Mulder H, Holm C, Sherry AD, Malloy CR: Effect of murine strain on metabolic pathways of glucose

- production after brief or prolonged fasting. *Am J Physiol Endocrinol Metab* 289:E53-E61, 2005
16. Landau B, Bartsch GE: Estimations of pathway contributions to glucose metabolism and the transaldolase reactions. *J Biol Chem* 241:741-749, 1966
 17. Fletcher SJ, Herlihy JM, Alberty WJ, Knowles JR: Energetics of triosephosphate isomerase: the appearance of solvent tritium in substrate glyceraldehyde 3-phosphate and in product. *Biochemistry* 15:5612-5617, 1976
 18. Rieder SV, Rose IA: The mechanism of the triosephosphate isomerase reaction. *J Biol Chem* 234:1007-1010, 1959
 19. Butler PC, Caumo A, Zerman A, O'Brien PC, Cobelli C, Rizza RA: Methods for assessment of the rate of onset and offset of insulin action during nonsteady state in humans. *Am.J.Physiol.* 264:E548-E569, 1993
 20. Rizza RA, Mandarino LJ, Gerich JE: Dose-response characteristics for effects of insulin on production and utilization of glucose in man. *Am.J.Physiol.* 240:E630-E639, 1981
 21. Stingl H, Chandramouli V, Schumann WC, Brehm A, Nowotny P, Waldhausl W, Landau BR, Roden M: Changes in hepatic glycogen cycling during a glucose load in healthy humans. *Diabetologia* 49:360-368, 2006
 22. Jones JG, Barosa C, Gomes F, Mendes AC, Delgado TC, Diogo L, Darcia P, Bastos M, Barros L, Fagulha A, Baptista C, Carvalheiro M, Caldeira MM: NMR derivatives for quantification of ^2H and ^{13}C -enrichment of human glucuronide from metabolic tracers. *Journal of Carbohydrate Chemistry* 25:203-217, 2006
 23. Burgess SC, Nuss M, Chandramouli V, Hardin DS, Rice M, Landau BR, Malloy CR, Sherry AD: Analysis of gluconeogenic pathways in vivo by distribution of ^2H in plasma glucose: comparison of nuclear magnetic resonance and mass spectrometry. *Anal Biochem* 318:321-324, 2003
 24. Landau BR, Wahren J, Chandramouli V, Schumann WC, Ekberg K: Use of $^2\text{H}_2\text{O}$ for estimating rates of gluconeogenesis: Application to the fasted state. *Journal of Clinical Investigation* 95:172-178, 1995
 25. Ljungdahl L, Wood HG, Racker E, Couri D: Formation of unequally labeled fructose 6-phosphate by an exchange reaction catalyzed by transaldolase. *J Biol Chem* 236:1622-1625, 1961
 26. Leadlay PF, Alberty WJ, Knowles JR: Energetics of triosephosphate isomerase: deuterium isotope effects in the enzyme-catalyzed reaction. *Biochemistry* 15:5617-5620, 1976
 27. Maister SG, Pett CP, Alberty WJ, Knowles JR: Energetics of triosephosphate isomerase: the appearance of solvent tritium in substrate dihydroxyacetone phosphate and in product. *Biochemistry* 15:5607-5612, 1976
 28. Rose IA, Kellermeyer R, Stjernholm R, Wood HG: The distribution of C^{14} in glycogen from deuterated glycerol- C^{14} as a measure of the effectiveness of triosephosphate isomerase *in vivo*. *J Biol Chem* 237:3325-3331, 1962
 29. Burgess SC, Weis B, Jones JG, Smith E, Merritt ME, Margolis D, Sherry AD, Malloy CR: Noninvasive evaluation of liver metabolism by ^2H and ^{13}C NMR isotopomer analysis of human urine. *Anal Biochem* 312:228-234, 2003
 30. Jones JG, Solomon MA, Cole SM, Sherry AD, Malloy CR: An integrated ^2H and ^{13}C NMR study of gluconeogenesis of TCA cycle flux in humans. *Am J Physiol Endocrinol Metab* 281:E848-E856, 2001

31. Burgess SC, Hausler N, Merritt M, Jeffrey FMH, Storey C, Milde A, Koshy S, Lindner J, Magnuson MA, Malloy CR, Sherry AD: Impaired tricarboxylic acid cycle activity in mouse livers lacking cytosolic phosphoenolpyruvate carboxykinase. *J Biol Chem* 47:48941-48949, 2004
32. Barzilai N, Rossetti L: Role of glucokinase and glucose-6-phosphatase in the acute and chronic regulation of hepatic glucose fluxes by insulin. *Journal of Biological Chemistry* 268:25019-25025, 1993

FIGURE LEGENDS

Figure 1

Reactions involved in the synthesis of glucose, glucose-6-phosphate and UDP-glucose

Figure 2

Plasma glucose, insulin and C-peptide concentrations observed before and during a hyperinsulinemic euglycemic clamp. An infusion of [3,5-²H₂] glucose was started at time -60 minutes and insulin at 0 minutes.

Figure 3

Rates of endogenous glucose production and glucose disappearance observed before and during a hyperinsulinemic euglycemic clamp. An infusion of [3,5-²H₂] glucose was started at time -60 minutes and insulin at 0 minutes.

Figure 4

Sample ²H nuclear magnetic resonance spectra of monoacetone glucose derived from UDP-glucose

Figure 5

The individual ratios of the deuterium enrichment on third and fifth carbons of UDP-glucose (C5/C3) measured in four subjects before and in six subjects during a hyperinsulinemic euglycemic clamp. An infusion of [3,5-²H₂] glucose was started at time -60 minutes and insulin at 0 minutes.

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5

