

Table appendix I. Haplotype association with plasma adiponectin levels in diabetic men without CVD

Haplotypes					Frequency	Mean difference, $\mu\text{g/mL}$			
-11365	-4034	-3964	+45	+276		Unadjusted	P	Adjusted*	P
C	A	A	T	G	0.20	Reference		Reference	
C	A	A	T	T	0.18	3.3	0.006	3.2	0.016
G	A	A	T	G	0.15	2.1	0.06	1.6	0.16
C	C	G	T	G	0.11	2.1	0.09	2.0	0.11
C	A	A	G	G	0.10	-0.4	0.39	-0.8	0.53
C	C	A	T	T	0.08	7.4	0.009	7.1	0.02
G	C	G	T	G	0.05	-2.5	0.91	-1.7	0.91

*: adjusting for age, BMI, smoking, alcohol consumption, physical activity, HbA1c, aspirin use, history of hypertension and history of hypercholesterolemia.

Table appendix II. Haplotype association with CVD risk in diabetic men

Haplotypes					Frequency		OR (95% CI)	P
-11365	-4034	-3964	+45	+276	Cases	Controls		
C	A	A	T	G	0.22	0.20	1.0	
C	A	A	T	T	0.19	0.18	1.09 (0.75-1.59)	0.65
G	A	A	T	G	0.13	0.14	0.91 (0.58-1.41)	0.67
C	C	G	T	G	0.15	0.11	1.30 (0.87-1.94)	0.20
C	A	A	G	G	0.10	0.10	1.02 (0.65-1.58)	0.93
C	C	A	T	T	0.08	0.08	0.97 (0.56-1.69)	0.92
G	C	G	T	G	0.04	0.05	0.86 (0.47-1.60)	0.64

Abbreviations: CVD, cardiovascular diseases; OR, odds ratio.