

# Effect of Oral Creatine Supplementation on Human Muscle GLUT4 Protein Content After Immobilization

B. Op 't Eijnde, B. Ursø, E.A. Richter, P.L. Greenhaff, and P. Hespel

The purpose of this study was to investigate the effect of oral creatine supplementation on muscle GLUT4 protein content and total creatine and glycogen content during muscle disuse and subsequent training. A double-blind placebo-controlled trial was performed with 22 young healthy volunteers. The right leg of each subject was immobilized using a cast for 2 weeks, after which subjects participated in a 10-week heavy resistance training program involving the knee-extensor muscles (three sessions per week). Half of the subjects received creatine monohydrate supplements (20 g daily during the immobilization period and 15 and 5 g daily during the first 3 and the last 7 weeks of rehabilitation training, respectively), whereas the other 11 subjects ingested placebo (maltodextrine). Muscle GLUT4 protein content and glycogen and total creatine concentrations were assayed in needle biopsy samples from the vastus lateralis muscle before and after immobilization and after 3 and 10 weeks of training. Immobilization decreased GLUT4 in the placebo group ( $-20\%$ ,  $P < 0.05$ ), but not in the creatine group ( $+9\%$  NS). Glycogen and total creatine were unchanged in both groups during the immobilization period. In the placebo group, during training, GLUT4 was normalized, and glycogen and total creatine were stable. Conversely, in the creatine group, GLUT4 increased by  $\sim 40\%$  ( $P < 0.05$ ) during rehabilitation. Muscle glycogen and total creatine levels were higher in the creatine group after 3 weeks of rehabilitation ( $P < 0.05$ ), but not after 10 weeks of rehabilitation. We concluded that 1) oral creatine supplementation offsets the decline in muscle GLUT4 protein content that occurs during immobilization, and 2) oral creatine supplementation increases GLUT4 protein content during subsequent rehabilitation training in healthy subjects. *Diabetes* 50:18–23, 2001

It is well established that high-dose (20–25 g per day) oral creatine intake can rapidly (3–5 days) raise muscle total creatine content. This elevation in muscle creatine storage is associated with increased muscle power output during short high-intensity exercise. In addition, it has been shown that long-term creatine intake can enhance the effects of weight training on muscle volume and strength (1,2). The use of creatine as an ergogenic supplement in sports has prompted interest in the potential of oral creatine supplementation to treat muscle atrophy and neuromuscular diseases. Thus, the recovery of muscle disuse atrophy due to immobilization was significantly enhanced by creatine supplementation (P.H., B.O.E., M. Van Leemputte, B.U., P.L.G., V. Labarque, S. Dymarkowski, P. Van Hecke, E.A.R, unpublished observations). Furthermore, creatine supplementation was found to have a beneficial impact on muscle functional capacity in various modes of mitochondrial cytopathies (3) and muscle dystrophies (4). At the same time, evidence is accumulating to suggest that creatine supplementation may be an effective neuroprotective agent to treat neurodegenerative diseases (5–7).

Interestingly, a number of recent observations also indicate that creatine supplementation might have a beneficial impact on glucose regulation. For instance, it has been shown that the ingestion of creatine in combination with carbohydrate supplements can stimulate postexercise muscle glycogen resynthesis (8), which is conceivably due to enhanced insulin-mediated muscle glucose uptake (9). Similarly, creatine intake in conjunction with carbohydrates was found to result in greater muscle creatine accumulation than creatine intake alone (10), which may be due to the fact that both glucose transport and creatine transport (11) in muscle cells are stimulated by insulin. On the other hand, a number of in vitro studies have found that high extracellular concentrations of guanidine compounds, including creatine, stimulate pancreatic insulin secretion (12,13). However, the extracellular creatine concentrations obtained by oral creatine intake in humans do not affect insulin secretion (14,15). Perhaps the most striking evidence to suggest that creatine supplementation might be an effective strategy to treat insulin resistance comes from a recent study on transgenic Huntington mice. The addition of creatine to the diet of the Huntington mice resulted in a marked neuroprotective effect and significantly reduced the hyperglycemia typical of these mice, while improving the glucose response to intravenous glucose injection (5).

Based on the above evidence, we speculate that creatine supplementation may enhance insulin-mediated muscle glucose uptake and glycogen synthesis, thereby beneficially impacting whole-body glucose homeostasis. This creatine

From the Faculty of Physical Education and Physiotherapy (B.O.E., P.H.), Exercise Physiology and Biomechanics Laboratory, Katholieke Universiteit Leuven, Leuven, Belgium; the Department of Human Physiology (B.U., E.A.R.), Copenhagen Muscle Research Center, University of Copenhagen, Copenhagen, Denmark; and the School of Biomedical Sciences (P.L.G.), Queens Medical Center, University of Nottingham, Nottingham, U.K.

Address correspondence and reprint requests to Peter Hespel, PhD, Faculty of Physical Education and Physiotherapy, Exercise Physiology and Biomechanics Laboratory, Tervuursevest 101, B-3001 Leuven, Belgium. E-mail: peter.hespel@flok.kuleuven.ac.be.

Received for publication 13 September 2000 and accepted 24 October 2000. Posted on the World Wide Web at [www.diabetes.org/diabetes](http://www.diabetes.org/diabetes) on 30 November 2000.

AMPK, AMP-activated protein kinase; CK, creatine kinase; DW, dry weight; MAPK, mitogen-activated protein kinase; RM, repetition maximum.

response might be particularly relevant to the prevention and/or treatment of disease states characterized by peripheral insulin resistance, such as type 2 diabetes, obesity, and inactivity (16). Furthermore, it is well established that muscle inactivity and training are effective stimuli to down- and upregulate muscle GLUT4 content and peripheral insulin sensitivity, respectively (17). Therefore, we investigated the effect of creatine supplementation on muscle GLUT4 protein content and total creatine and glycogen concentration in healthy volunteers during 2 weeks of leg immobilization and during 10 weeks of subsequent rehabilitation training. This report is part of a larger study (P.H., B.O.E., M. Van Leemputte, B.U., P.L.G., V. Labarque, S. Dymarkowski, P. Van Hecke, E.A.R.) that investigated the effects of creatine supplementation on muscle functional capacity during disuse atrophy in healthy subjects.

## RESEARCH DESIGN AND METHODS

**Subjects.** A total of 13 men and 9 women, aged 20–23 years, gave their informed written consent to participate in the study. They were instructed to abstain from taking any medication and to avoid making any changes in their usual physical activity level and other living habits during the period of the study. However, three of the women were taking oral contraceptives for the duration of the study. The local ethics committee approved the study protocol.

**Study protocol.** A double-blind study was performed over a 12-week period. At the start of the study, subjects were systematically assigned to either a creatine or a placebo group based on quadriceps muscle cross-sectional area and maximal isometric knee-extension torque to obtain two groups of similar distribution (P.H., B.O.E., M. Van Leemputte, B.U., P.L.G., V. Labarque, S. Dymarkowski, P. Van Hecke, E.A.R., unpublished observations). After baseline measurements had been taken, a light polyester cast, extending from groin to ankle, immobilized each subject's right leg at a knee angle of  $\sim 160^\circ$  for 2 weeks. Thereafter, the cast was removed, and the subjects underwent a standardized 10-week rehabilitation program. Each training session consisted of four series of 12 unilateral knee-extensions on a knee-extension apparatus, at a workload of 60% of maximal isometric knee-extension torque and at a rate of three sessions per week. Maximal knee-extension torque was measured at a  $90^\circ$  knee-angle at the start of each session using a calibrated force transducer. During the final 7 weeks of the training period, the series of four unilateral knee-extensions was increased to six. All training sessions were supervised by one of the investigators. During immobilization, the creatine group received 5 g creatine monohydrate four times per day, whereas the placebo group received placebo supplements (5 g maltodextrine, four times per day). During the training period, creatine and placebo supplementation was reduced to 5 g three times per day from week 1 to 3 and then to a single 5-g daily dose from week 4 to 10. The creatine supplements were flavored by the addition of citrate (60 mg/g creatine) and maltodextrine (940 mg/g creatine), whereas the placebo group ingested maltodextrine containing citrate (40 mg/g maltodextrine). Creatine and placebo powders were identical in taste and appearance. Before and after 2 weeks of immobilization, and after 3 and 10 weeks of rehabilitation, a percutaneous needle biopsy from the vastus lateralis muscle was obtained. The last training session preceded muscle sampling by at least 48 h. In addition, the subjects received a standardized dinner (855 kcal, 47% carbohydrate, 25% fat, and 28% protein) the evening before and a standardized breakfast (320 kcal, 65% carbohydrate, 15% fat, and 20% protein) the morning of muscle sampling. To collect each muscle biopsy, an incision was made through the skin and muscle fascia under local anesthesia (2–3 ml 1% lidocaine). During sessions 2–4, the incision was made either proximal or distal to the incision made at an earlier session. On removal from the limb, a piece of each muscle biopsy was immediately blotted and cleaned from visible connective tissue, rapidly frozen in liquid nitrogen, and stored at  $-80^\circ\text{C}$  for subsequent biochemical and immunochemical analyses.

**Biochemical and immunochemical analyses.** The biopsy samples were first freeze-dried, then washed twice in petroleum ether to remove fat, and finally dissected free of the remaining visible blood and connective tissue. A fraction of each sample was pulverized, and the powdered extracts were used for spectrophotometric determination of glycogen and free creatine and phosphocreatine concentrations (18). Another fraction was used for GLUT4 determination. An aliquot of the freeze-dried muscle was homogenized (Polytron) for 30 s on ice in a buffer with the following composition: 150 mmol/l NaCl, 1% NP<sub>4</sub>O, 0.5% deoxycholate, 0.1% SDS, and 50 mmol/l Tris, pH 8. The homogenate was incubated on ice for 1 h and spun for 15 min at 13,000g, and the supernatant (extract) was collected for analysis. Then, 100  $\mu\text{g}$  of the

extract were resolved by SDS-PAGE before electroblotting to polyvinylidene fluoride membranes. GLUT4 proteins were then detected by incubation in Tris-buffered saline with Tween (150 mmol/l NaCl, 50 mmol/l Tris, and 0.1% Tween 20) after blocking in 1% bovine serum albumin with a specific goat polyclonal antibody against the 13 COOH-terminal amino acids of GLUT4. Finally, GLUT4 was visualized by an alkaline phosphatase-labeled antibody and quantified on a phosphoimager (STORM; Molecular Dynamics, Sunnyvale, CA).

**Data analysis.** Data are means  $\pm$  SE. Muscle total creatine concentration was calculated as the sum of free creatine and phosphocreatine. Treatment effects (creatine versus placebo) were evaluated by a two-way analysis of variance, which was covariate adjusted for the baseline values (Statistica; Statsoft, Tulsa, OK). In addition to these primary analyses, we did a one-way analysis of variance to compare the values after immobilization and rehabilitation with the corresponding baseline values within each group. The statistical analyses of the GLUT4 data were performed on the raw data (densitometric counts).  $P < 0.05$  was considered statistically significant.

## RESULTS

**Muscle GLUT4 content.** Muscle GLUT4 concentrations were expressed relative to the corresponding baseline values that were set equal to 1 (Fig. 1). Muscle GLUT4 content at baseline was similar between the groups. In the placebo group, 2 weeks of immobilization decreased GLUT4 content on an average of 22% (range  $-10$  to  $-35\%$ ,  $P < 0.05$ ). Conversely, in the creatine group, muscle GLUT4 protein was stable ( $+9\%$  NS). In the placebo group, the rehabilitation training restored muscle GLUT4 content within 3 weeks to the baseline value, where it remained. However, in the creatine group, muscle GLUT4 content progressively increased during the 10-week rehabilitation period to a value that was  $\sim 40\%$  higher than in the placebo group at the end of the study ( $P < 0.05$ ).

**Muscle glycogen.** The initial muscle glycogen concentration was  $407 \pm 43$  mmol/kg dry weight (DW) in the placebo group versus  $379 \pm 19$  mmol/kg DW in the creatine group (NS) (Fig. 2). Immobilization did not change muscle glycogen concentration in either group. However, during the initial 3 weeks of rehabilitation training, muscle glycogen markedly increased in the creatine group ( $P < 0.05$ ), whereas it did not significantly change in the placebo group. Thus, after 3 weeks, muscle glycogen concentration was higher ( $P < 0.05$ ) in the creatine group ( $660 \pm 70$  mmol/kg DW) than in the placebo group ( $520 \pm 60$  mmol/kg DW). However, during the final 7 weeks of rehabilitation training, muscle glycogen reverted to baseline values in both groups.

**Muscle creatine content.** The muscle phosphocreatine and free creatine concentrations at baseline were similar between both groups (Table 1). During immobilization, phosphocreatine concentration decreased to  $\sim 15\%$  below the baseline value in the placebo group ( $P < 0.05$ ). This decrease was negated by creatine supplementation ( $P < 0.05$ ). In the placebo group, muscle phosphocreatine concentration returned to the preimmobilization baseline level within the initial 3 weeks of the rehabilitation period, after which it remained stable. On the other hand, in the creatine group, compared with the placebo group, the muscle phosphocreatine concentration increased to  $\sim 12\%$  above baseline value after 3 weeks of rehabilitation ( $P < 0.05$ ). However, this increase above baseline in phosphocreatine was reversed during the final stage of the rehabilitation period. Throughout the study, the muscle free creatine concentrations were not significantly different between the placebo and the creatine groups. In the placebo group, muscle total creatine concentration was not significantly changed compared with the baseline value during either immobilization or rehabilitation. Yet




FIG. 1. Effect of creatine supplementation on muscle GLUT4 protein content during immobilization and subsequent rehabilitation training. Data are means  $\pm$  SE ( $n = 8$ ) and are expressed relative to the baseline value that was set to be equal to 1. Muscle samples were taken from the vastus lateralis muscle before and after 2 weeks of immobilization and after 3 and 10 weeks of rehabilitation of the right leg. During immobilization and rehabilitation, subjects ingested creatine monohydrate (■) or placebo (□). See RESEARCH DESIGN AND METHODS for further details. \*Significant treatment effect compared with placebo,  $P < 0.05$ ; §significant time effect compared with the preimmobilization value.

in the creatine group, compared with the placebo group, the muscle total creatine concentration was higher at the end of the immobilization period, as well as after 3 weeks of rehabilitation ( $P < 0.05$ ). However, along with the declining muscle phosphocreatine levels, muscle total creatine returned to baseline by the end of the study.

## DISCUSSION

Our study investigated the impact of creatine supplementation on muscle GLUT4 content and glycogen and total creatine concentrations in healthy subjects during 2 weeks of voluntary leg immobilization followed by 10 weeks of reha-

bilitation training. Our data are the first to show that creatine supplementation prevents the loss of GLUT4 protein during muscle disuse and increases muscle GLUT4 content above normal levels during subsequent rehabilitation. Furthermore, muscle glycogen concentration was increased during the initial stages of the creatine supplementation.

Glucose transport across the plasma membrane is the rate-limiting step for glucose metabolism. Hence, muscle GLUT4 content is a primary determinant of insulin-stimulated muscle glucose uptake and metabolism (16). Thus, increasing muscle GLUT4 content by transgenic overexpression or by increased contractile activity enhances maximal insulin-stimulated muscle




FIG. 2. Effect of creatine supplementation on muscle glycogen concentration during immobilization and subsequent rehabilitation training. Data are means  $\pm$  SE ( $n = 8$ ). Muscle samples were taken from the vastus lateralis muscle before and after 2 weeks of immobilization and after 3 and 10 weeks of rehabilitation of the right leg. During immobilization and rehabilitation, subjects ingested creatine monohydrate (■) or placebo (□). See RESEARCH DESIGN AND METHODS for further details. \*Significant treatment effect compared with placebo,  $P < 0.05$ ; §significant time effect compared with the preimmobilization value.

TABLE 1  
Effect of creatine supplementation on muscle creatine content during immobilization and subsequent rehabilitation training

|                              | Before immobilization | After immobilization | 3 weeks of rehabilitation | 10 weeks of rehabilitation |
|------------------------------|-----------------------|----------------------|---------------------------|----------------------------|
| Free creatine (mmol/kg DW)   |                       |                      |                           |                            |
| Placebo                      | 31.3 ± 3.3            | 41.3 ± 3.6*          | 43.5 ± 5.4*               | 37.7 ± 2.9                 |
| Creatine                     | 30.6 ± 2.9            | 48.5 ± 4.5*          | 53.9 ± 5.4*               | 43.4 ± 4.0                 |
| Phosphocreatine (mmol/kg DW) |                       |                      |                           |                            |
| Placebo                      | 76.5 ± 1.8            | 64.9 ± 3.1*          | 73.8 ± 2.6                | 71.6 ± 2.2                 |
| Creatine                     | 82.4 ± 6.2            | 80.2 ± 5.8†          | 89.7 ± 6.8†               | 75.1 ± 6.3                 |
| Total creatine (mmol/kg DW)  |                       |                      |                           |                            |
| Placebo                      | 108.8 ± 2.8           | 106.2 ± 5.7          | 117.3 ± 5.1               | 109.3 ± 3.4                |
| Creatine                     | 113.9 ± 8.4           | 128.7 ± 9.9*†        | 143.6 ± 11.6*†            | 118.5 ± 8.0                |

Data are means ± SE of eight observations and represent concentrations measured in needle biopsy samples obtained from vastus lateralis muscle. Total creatine concentration was calculated as the sum of free creatine and phosphocreatine concentrations measured. Immobilization and rehabilitation procedures are described in RESEARCH DESIGN AND METHODS. \*Significant time-effect compared with the preimmobilization value,  $P < 0.05$ ; †significant treatment effect compared with placebo,  $P < 0.05$ .

glucose uptake. Conversely, reducing the content of GLUT4 by GLUT4 knockout, denervation, or aging impairs insulin-mediated muscle glucose uptake (19). Our data, therefore, suggest that creatine supplementation in humans may increase insulin sensitivity by increasing muscle GLUT4 content.

Over the last decade, substantial evidence has accumulated to show that endurance exercise training elevates muscle GLUT4 content and insulin-stimulated glucose uptake in both healthy (17,20–28) and insulin-resistant muscles (29,30). In this respect, the current study shows that in healthy individuals, a low volume (3 weekly sessions) of moderate resistance training (60% of 1 repetition maximum [RM]), in contrast with endurance training (23–26, 28) or daily maximal resistance training (31), is not a sufficient stimulus to increase muscle GLUT4 content. Ten weeks of rehabilitation training per se did not increase muscle GLUT4 content above the baseline level (Fig. 1). However, the same training regimen in conjunction with oral creatine supplementation resulted in a marked increase of muscle GLUT4 protein content. In fact, our observations indicate that oral creatine supplementation can probably increase GLUT4 protein content in skeletal musculature independent of exercise training. In keeping with earlier observations (17,20–22,31,32), muscle deconditioning by immobilization in the placebo subjects reduced GLUT4 protein content (~20%). Nevertheless, at the end of the immobilization period, GLUT4 content in the creatine group tended to increase by ~10%, which resulted in a 30% difference in muscle GLUT4 between placebo and creatine supplementation in the absence of a training sore, it is reasonable to conclude that creatine supplementation can increase GLUT4 protein content in human musculature during episodes of either reduced or increased physical activity.

Based on the current knowledge, it is difficult to reveal the molecular basis for the increase in muscle GLUT4 content that occurs during creatine supplementation. It has recently been observed in rats that short-term administration of aminoimidazole-4-carboximide riboside, an AMP-activated protein kinase (AMPK) agonist, increases muscle GLUT4 content (33). Creatine administration that increases AMPK activity by decreasing the phosphocreatine-to-creatine ratio (34) may, thus, explain the increase in GLUT4 protein content in the creatine group. And yet, in both groups the phosphocreatine-

to-creatine ratio decreased to the same degree during immobilization and remained below the baseline value during the subsequent rehabilitation period. Furthermore, it has recently been shown that the creatine kinase (CK) and AMPK enzymes colocalize in muscle cells (34). According to the prevailing opinion, in skeletal muscle, such coupling should serve to suppress muscle AMPK activity by maintaining high local ATP:AMP and phosphocreatine-to-creatine ratios in conditions of cellular stress, such as contractions (35). If anything, this inhibitory action is enhanced by the increased muscle phosphocreatine concentration established during the creatine supplementation (Table 1). Thus, evidence for a possible creatine-induced increase in AMPK activity has not been found. Alternatively, there is substantial evidence to suggest that cellular hydration status is an important factor controlling cellular protein turnover (36), which in muscle cells, excluding the contractile proteins, may involve other proteins important to energy homeostasis, such as GLUT4. Creatine is cotransported with Na ions across the sarcolemma, which initiates influx of Cl<sup>-</sup> and water to balance electroneutrality and osmolality (11). The resulting increase of cell volume may, in turn, act as an anabolic proliferative signal, which involves activation of the mitogen-activated protein kinase (MAPK) signaling cascade that plays a pivotal role in muscle protein synthesis regulation (37,38). It is warranted to further explore the possible role of intracellular creatine content in modulating the concerted actions of CK, AMPK, and MAPK in regulating GLUT4 synthesis and degradation in muscle cells.

The bulk of glucose in the human body is stored as muscle glycogen. The presence of a high muscle glycogen concentration, in general, indicates adequate insulin stimulation of muscle glucose uptake and glycogen synthesis. Furthermore, a high muscle glycogen concentration is a prerequisite for optimal endurance exercise performance (39). Robinson et al. (8) have recently demonstrated that carbohydrate intake in conjunction with creatine supplementation resulted in greater postexercise muscle glycogen resynthesis than carbohydrate intake alone. Accordingly, in the current study, during the initial 3 weeks of rehabilitation training, muscle glycogen concentration increased by ~30% in the placebo group, whereas a threefold greater increase occurred in the creatine group. This higher-than-average glycogen level,

established by creatine supplementation (>650 mmol/kg DW) (Fig. 2), corresponds with common glycogen levels in young healthy subjects after glycogen "supercompensation" (39). Given that no dietary instructions were administered to the subjects, our findings suggest that the addition of creatine supplementation to a standard diet may eventually result in a postexercise increment of muscle glycogen concentration similar to that found after a classical carbohydrate-enriched glycogen supercompensation dietary protocol (39). Interestingly, after 5 weeks of creatine supplementation, the increase of muscle glycogen content vanished, despite continued creatine supplementation. In fact, during both immobilization and rehabilitation, the pattern of muscle glycogen changes closely mimicked the fluctuations of muscle total creatine content (Table 1) (Fig. 2). In this respect, Low et al. (40) have provided clear evidence that osmotic swelling of muscle cells is a potent stimulus to muscle glycogen synthesis. The 30 mmol/kg DW increase of muscle total creatine, established after 3 weeks of training in the creatine group, was therefore probably sufficient to induce a degree of cell swelling necessary to enhance insulin-stimulated glycogen synthesis (40,36). If such an osmotic trigger mechanism indeed regulates insulin action on glycogen synthesis during creatine supplementation, then the decrease in muscle creatine content beyond 3 weeks of training might also explain the concurrent decrease in the muscle glycogen storage. The mechanism behind the decrease in muscle creatine content during the final stage of the study, despite continued creatine ingestion at a rate presumed to be sufficient for maintaining an elevated muscle creatine content (5 g/day), is unclear (2,41). Studies in rats have demonstrated that long-term high-dose creatine feeding induces a downregulation of muscle total Na-creatine cotransporter protein content (42). In addition, the low creatine transporter content in failing human myocardium has been found to be associated with a decrease in intracellular creatine storage (43).

In conclusion, the current findings provide strong evidence to suggest that 1) oral creatine supplementation can offset the decline of muscle GLUT4 protein content in skeletal musculature during disuse atrophy, and 2) oral creatine supplementation increases GLUT4 content during subsequent rehabilitation training. Based on the present findings, it is warranted to evaluate the potential of long-term creatine supplementation as a strategy to prevent or treat disease conditions characterized by peripheral insulin resistance.

#### ACKNOWLEDGMENTS

This study was supported by grant G.0331.98 from the Fonds voor Wetenschappelijk Onderzoek Vlaanderen, grant OT/94/31 from the Onderzoeksraad K.U.-Leuven, and grant 504-14 from the Danish National Research Foundation.

The authors thank Betina Bolmgreen, Irene Beck Nielsen, and Monique Ramaekers for providing skilled technical assistance.

#### REFERENCES

- American College of Sports Medicine roundtable: The physiological and health effects of oral creatine supplementation. *Med Sci Sports Exerc* 32: 706-717, 2000
- Vandenbergh K, Goris M, Van Hecke P, Van Leemputte M, Vangerven L, Hespel P: Long-term creatine intake is beneficial to muscle performance during resistance training. *J Appl Physiol* 83:2055-2063, 1997
- Tarnopolsky M, Martin J: Creatine monohydrate increases strength in patients with neuromuscular disease. *Neurology* 52:854-857, 1999
- Walter MC, Lochmüller H, Reilich P, Klopstock T, Huber R, Hartard M, Hennig M, Pongratz D, Müller-Felber W: Creatine monohydrate in muscular dystrophies: a double-blind, placebo-controlled clinical study. *Neurology* 54:1848-1850, 2000
- Ferrante RJ, Andreassen OA, Jenkins BG, Dedeoglu A, Kuemmerle S, Kubilus JK, Kaddurah-Daouk R, Hersch SM, Flint Beal M: Neuroprotective effects of creatine in a transgenic mouse model of Huntington's disease. *J Neurosci* 20:4389-4397, 2000
- Klivenyi P, Ferrante RJ, Matthews RT, Bogdanov MB, Klein AM, Andreassen OA, Mueller G, Wermer M, Kaddurah-Daouk R, Beal MF: Neuroprotective effects of creatine in a transgenic animal model of amyotrophic lateral sclerosis. *Nat Med* 5:347-350, 1999
- Matthews RT, Yang L, Jenkins BG, Ferrante RJ, Rosen BR, Kaddurah-Daouk R, Beal MF: Neuroprotective effect of creatine and cyclocreatine and cyclocreatine in animal models of Huntington's disease. *J Neurosci* 18:156-163, 1998
- Robinson TM, Sewell DA, Hultman E, Greenhaff PL: Role of submaximal exercise in promoting creatine and glycogen accumulation in human skeletal muscle. *J Appl Physiol* 87:598-604, 1999
- Richter EA, Garetto LP, Goodman MN, Ruderman NB: Enhanced muscle glucose metabolism after exercise: modulation by local factors. *Am J Physiol* 246:E476-E482, 1984
- Green AL, Hultman E, Macdonald IA, Sewell DA, Greenhaff PL: Carbohydrate ingestion augments skeletal muscle creatine accumulation during creatine supplementation in humans. *Am J Physiol* 271:E821-E826, 1996
- Odoom JE, Kemp GJ, Radda GK: The regulation of total creatine content in a myoblast cell line. *Mol Cell Biochem* 158:179-188, 1996
- Alsever RN, Georg RH, Sussman KE: Stimulation of insulin secretion by guanidinoacetic acid and other guanidine derivatives. *Endocrinology* 86:332-336, 1970
- Gerbitz K-D, Gempel K, Brdiczka D: Mitochondria and diabetes: genetic, biochemical, and clinical implications of the cellular energy circuit. *Diabetes* 45:113-126, 1996
- Green AL, Simpson EJ, Littlewood JJ, Macdonald IA, Greenhaff PL: Carbohydrate ingestion augments creatine retention during creatine feeding in humans. *Acta Physiol Scand* 158:195-202, 1996
- Steenge GR, Lambourne J, Casey A, Macdonald IA, Greenhaff PL: Stimulatory effect of insulin on creatine accumulation in human skeletal muscle. *Am J Physiol* 275:E974-E979, 1998
- Shepherd PR, Kahn BB: Glucose transporters and insulin action: implications for insulin resistance and diabetes mellitus. *N Engl J Med* 341:248-257, 1999
- Houmard JA, Tyndall GL, Midyette JB, Hickey MS, Dolan PL, Gavigan KE, Weidner ML, Dohm GL: Effect of reduced training and training cessation on insulin action and muscle GLUT4. *J Appl Physiol* 81:1162-1168, 1996
- Harris RC, Hultman E, Nordesjö L-O: Glycogen, glycolytic intermediates and high-energy phosphates determined in biopsy samples of musculus quadriceps femoris of man at rest: methods and variance of values. *Scand J Clin Lab Invest* 33:109-120, 1974
- Charron MJ, Katz EB, Olson AL: GLUT4 gene regulation and manipulation. *J Biol Chem* 274:3253-3256, 1999
- Host HH, Hansen PA, Nolte LA, Chen MM, Holloszy JO: Rapid reversal of adaptive increases in muscle GLUT4 and glucose transport capacity after training cessation. *J Appl Physiol* 84:79-802, 1998
- McCoy M, Proietto J, Hargreaves M: Effect of detraining on GLUT4 protein in human skeletal muscle. *J Appl Physiol* 77:1532-1536, 1994
- Vukovich MD, Arciero PJ, Kohrt WM, Racette SB, Hansen PA, Holloszy JO: Changes in insulin action and GLUT4 with 6 days of inactivity in endurance runners. *J Appl Physiol* 80:240-244, 1996
- Houmard JA, Shinebarger MH, Dolan PL, Leggett-Frazier N, Bruner RK, McCammon MR, Israel RG, Dohm GL: Exercise training increases GLUT4 protein concentration in previously sedentary middle-aged men. *Am J Physiol (Endocrinol Metab)* 264:E896-E901, 1993
- Houmard JA, Hickey MS, Tyndall GL, Gavigan KE, Dohm GL: Seven days of exercise increase GLUT4 protein content in human skeletal muscle. *J Appl Physiol* 79:1936-1938, 1995
- Ebeling P, Bourey R, Koranyi L, Tuominen JA, Groop LC, Henriksson J, Mueckler M, Sovijärvi A, Koivisto VA: Mechanism of enhanced insulin sensitivity in athletes: increased blood flow, muscle glucose transport protein (GLUT4) concentration, and glycogen synthase activity. *J Clin Invest* 92:1623-1631, 1993
- Houmard JA, Egan PC, Neuffer PD, Friedman JE, Wheeler WS, Israel RG, Dohm GL: Elevated skeletal muscle glucose transporter levels in exercise-trained middle-aged men. *Am J Physiol* 261:E437-E443, 1991
- Gulve EA, Spina RJ: Effect of 7-10 days of cycle ergometer exercise on skeletal muscle GLUT4 protein content. *J Appl Physiol* 79:1562-1566, 1995
- Phillips SM, Han XX, Green HJ, Bonen A: Increments in skeletal muscle GLUT1 and GLUT4 after endurance training in humans. *Am J Physiol* 270: E456-E462, 1996
- Banks EA, Brozinick JT Jr, Yaspelkis III BB, Kang HY, Ivy JL: Muscle glucose transport, GLUT4 content, and degree of exercise training in obese Zucker rats. *Am J Physiol* 263:E1010-E1015, 1992

30. Brozinick JT Jr, Etgen GJ Jr, Yaspelkis III BB, Kang HY, Ivy JL: Effects of exercise training on muscle GLUT-4 protein content and translocation in obese Zucker rats. *Am J Physiol* 265:E419–E427, 1993
31. Tabata I, Suzuki Y, Fukunaga T, Yokozeki T, Akima H, Funato K: Resistance training affects GLUT4 content in skeletal muscle of humans after 19 days of head-down bed rest. *J Appl Physiol* 86:909–914, 1999
32. Henriksen EJ, Rodnick KJ, Mondon CE, James DE, Holloszy JO: Effect of denervation or unweighting on GLUT4 protein in rat soleus muscle. *J Appl Physiol* 70:2322–2327, 1991
33. Holmes BF, Kurth-Kraczek EJ, Winder WW: Chronic activation of 5'-AMP-activated protein kinase increases GLUT4, hexokinase, and glycogen in muscle. *J Appl Physiol* 87:1990–1995, 1999
34. Hardie DG, Carling D, Carlson M: The AMP-activated/SNF1 protein kinase subfamily: metabolic sensors of the eukaryotic cell? *Annu Rev Biochem* 67:821–855, 1998
35. Wallimann T, Wyss M, Brdiczka D, Nicolay K, Eppenberger HM: Intracellular compartmentation, structure and function of creatine kinase isoenzymes in tissues with high and fluctuating energy demands: the 'phosphocreatine circuit' for cellular energy homeostasis. *Biochem J* 281:21–40, 1992
36. Häussinger D, Roth E, Lang F, Gerok W: Cellular hydration status: an important determinant of protein catabolism in health and disease. *Lancet* 341:1330–1332, 1993
37. Niisato N, Post M, Van Driessche W, Marunaka Y: Cell swelling activates stress-activated protein kinases, p38 MAP kinase and JNK, in renal epithelial A6 cells. *Biochem Biophys Res Commun* 266:547–550, 1999
38. Tilly BC, Gaestel M, Engel K, Edixhoven MJ, de Jonge HR: Hypo-osmotic cell swelling activates the p38 MAP kinase signaling cascade. *FEBS Lett* 395:133–136, 1996
39. Sherman WM, Costill DL, Fink WJ, Miller JM: Effect of exercise-diet manipulation on muscle glycogen and its subsequent utilization during performance. *Int J Sports Med* 2:114–118, 1981
40. Low SY, Rennie MJ, Taylor PM: Modulation of glycogen synthesis in rat skeletal muscle by changes in cell volume. *J Physiol (Lond)* 495:299–303, 1996
41. Hultman E, Söderlund K, Timmons JA, Cederblad G, Greenhaff PL: Muscle creatine loading in men. *J Appl Physiol* 81:232–237, 1996
42. Guerrero-Ontiveros ML, Wallimann T: Creatine supplementation in health and disease: effects of chronic creatine ingestion in vivo: downregulation of the expression of creatine transporter isoforms in skeletal muscle. *Mol Cell Biochem* 184:427–437, 1998
43. Neubauer S, Remkes H, Spindler M, Horn M, Wiesmann F, Prestle J, Walzel B, Ertl G, Hasenfuss G, Wallimann T: Downregulation of the Na<sup>+</sup>-creatine cotransporter in failing human myocardium and in experimental heart failure. *Circulation* 100:1847–1850, 1999