

# The Common Single Nucleotide Polymorphism E23K in $K_{IR}6.2$ Sensitizes Pancreatic $\beta$ -Cell ATP-Sensitive Potassium Channels Toward Activation Through Nucleoside Diphosphates

Christina Schwanstecher, Bettina Neugebauer, Miriam Schulz, and Mathias Schwanstecher

**E23K, a common polymorphism in the pore-forming subunit  $K_{IR}6.2$  of pancreatic  $\beta$ -cell ATP-sensitive  $K^+$  ( $K_{ATP}$ ) channels, is functionally relevant and thus might play a major role in the pathophysiology of common type 2 diabetes. In this study, we show that in the simultaneous presence of activatory and inhibitory nucleotides, the polymorphism exerts opposite effects on the potencies of these modulators: channel opening through nucleoside diphosphates is facilitated, whereas sensitivity toward inhibition through ATP is slightly decreased. The results support the conclusion that E23K predisposes to type 2 diabetes by changing the channel's response to physiological variation of cytosolic nucleotides, resulting in  $K_{ATP}$  overactivity and discrete inhibition of insulin release. *Diabetes* 51 (Suppl. 3): S363–S367, 2002**

**T**ype 2 diabetes is generally perceived as a polygenic disorder, with disease development being influenced by both hereditary and environmental factors (1). Genes encoding for key components of insulin secretion and glucose metabolism pathways have been widely considered as targets for defects in type 2 diabetes (2). One of these key proteins is the ATP-sensitive  $K^+$  channel ( $K_{ATP}$  channel) in pancreatic  $\beta$ -cells. This channel critically controls insulin secretion by coupling metabolism to electrical activity (3). The  $\beta$ -cell channel is assembled with a tetradimeric stoichiometry from two structurally distinct subunits: the inwardly rectifying potassium channel subunit ( $K_{IR}6.2$ ), which forms the pore, and the regulatory sulfonylurea receptor subunit 1 (SUR1) (4,5). While hypoglycemic sulfonylureas (e.g., glibenclamide) exert their effects on channel activity by

interacting with SUR1, there is strong evidence that the receptor site for inhibitory ATP is located on  $K_{IR}6.2$ .

E23K (substitution of a lysine [K] for a glutamic acid [E] in position 23) is one of three common missense single nucleotide polymorphisms (SNPs) that have been observed in  $K_{IR}6.2$  (E23K, L270V, I337V) (6–10). Recently, we presented evidence that this polymorphism predisposes to type 2 diabetes by inducing overactivity of  $K_{ATP}$  channels in the pancreatic  $\beta$ -cell (11). In particular, E23K markedly affects channel gating, significantly reducing the time spent in long interburst closed states and thereby producing an evident increase of spontaneous open probability ( $P_O$ ). Consistent with the idea that nucleotide-induced channel inhibition results from interaction with the interburst closed states (12,13), sensitivity toward inhibitory  $ATP^{4-}$  was found to be decreased (11).

In intact cells, however, other factors besides inhibitory ATP contribute to regulation of channel activity, with ADP presumably representing the most important of these additional parameters (14). ADP has at least three distinct effects on channel activity: 1) Free  $ADP^{3-}$  inhibits channel activity in isolated patches with a potency slightly lower than that of  $ATP^{4-}$  (half-maximal inhibitory concentration value [ $IC_{50}$ ] 49 vs. 8.9  $\mu$ mol/l, respectively) (11,15). 2) The  $Mg^{2+}$  complex of ADP (MgADP) per se potently activates the channels by interaction with a separate nucleotide site residing on SUR1 (16). 3) MgADP strongly reduces sensitivity toward inhibitory nucleotides (17).

In this study, we analyzed the impact of the E23K polymorphism of  $K_{IR}6.2$  on the regulation of channel activity by activating nucleoside diphosphates. The results support the conclusion that the polymorphism predisposes to type 2 diabetes by inducing overactivity of pancreatic  $\beta$ -cell  $K_{ATP}$  channels.

From the Institute of Pharmacology and Toxicology, University of Braunschweig, Braunschweig, Germany.

Address correspondence and reprint requests to Dr. Christina Schwanstecher, Institut für Pharmakologie und Toxikologie, Universität Braunschweig, Mendelssohnstraße 1, 38106 Braunschweig, Germany. E-mail: c.schwanstecher@tu-bs.de.

Received for publication 19 March 2002 and accepted in revised form 23 April 2002.

DMEM, Dulbecco's modified Eagle's medium;  $EC_{50}$ , half-maximal effective concentration;  $E_{max}$ , maximal effect; GDP, guanosine 5'-diphosphate; HG, high glucose;  $IC_{50}$ , half-maximal inhibitory concentration value;  $K_{ATP}$  channel, ATP-sensitive  $K^+$  channel; Kir channel, inwardly rectifying potassium channel; NDPK, nucleoside diphosphate kinase;  $P_O$ , open probability; SNP, single nucleotide polymorphism; SUR, sulfonylurea receptor.


The symposium and the publication of this article have been made possible by an unrestricted educational grant from Servier, Paris.

## RESEARCH DESIGN AND METHODS

**Materials.** All chemicals were obtained from the sources described elsewhere (18).

**Molecular biology.**  $K_{IR}6.2$  mutants were constructed as previously described (11). Briefly, point mutations were introduced into human (GenBank Q14654) or mouse (GenBank D50581)  $K_{IR}6.2$  by standard molecular biology techniques, and constructs were sequenced to verify PCR fidelity before transfection. For analysis of combinations of the polymorphisms (Figs. 1C and 2C), mutations were sequentially inferred into the same  $K_{IR}6.2$  cDNA.

**Electrophysiology.** Transfections were performed as previously described (11,18). COS-1 cells cultured in Dulbecco's modified Eagle's medium (DMEM)–high glucose (HG) (10 mmol/l glucose) supplemented with 10% FCS were plated at a density of  $8 \times 10^4$  cells per dish (35 mm) and allowed to attach overnight. Next, 20  $\mu$ g of pECE-human SUR1 (GenBank NP\_000343) comple-



**FIG. 1.** Effect of E23K on ATP potency in the presence of GDP. **A:** ATP sensitivity of channels reconstituted in COS-1 cells by co-expression of human SUR1 with human  $K_{IR}6.2_{wt}$  or  $K_{IR}6.2_{E23K}$ , as indicated. Representative currents recorded at  $-50$  mV from inside-out patches exposed to ATP and GDP (0.3 mmol/l), as shown by the lines above the records. Inward currents are depicted as downward deflections. Dashed lines indicate residual channel activity (percent) in the presence of 100  $\mu$ mol/l ATP (plus 0.3 mmol/l GDP). **B:** Potencies of ATP in the presence of 0.3 mmol/l GDP. Channel inhibition was recorded in inside-out patches as shown in **A**. Results are given as means  $\pm$  SE of 10–15 independent experiments.  $EC_{50}$  values and Hill coefficients are as follows: wild-type ( $\blacksquare$ ),  $71 \pm 4.5$   $\mu$ mol/l and 1.22; E23K ( $\square$ ),  $120 \pm 5.2$   $\mu$ mol/l and 1.34.  $P < 0.01$  for comparison of the  $EC_{50}$  values. **C:** The effect of E23K on the  $EC_{50}$  value for ATP was not altered by the additional introduction of L270V and/or I337V into human  $K_{IR}6.2$ . \* $P < 0.05$  vs. wild-type ( $n = 8-10$ ).

mentary DNA and 20  $\mu$ g of pECE-human  $K_{IR}6.2$  complementary DNA were mixed and used to transfect six 35-mm plates. In control experiments, we used pECE-hamster SUR1 (GenBank A56248) and pECE-mouse  $K_{IR}6.2$  instead of the human isoforms. For transfection, the cells were incubated for 4 h in a Tris-buffered salt solution containing DNA (5–10  $\mu$ g/ml) plus DEAE-dextran (1 mg/ml), 2 min in HEPES-buffered salt solution plus dimethyl sulfoxide (10%), and 4 h in DMEM-HG plus chloroquine (100  $\mu$ mol/l). Cells were then returned to DMEM-HG plus 10% FCS. Experiments in the inside-out configuration of the patch-clamp technique were performed 1–2 days after transfection at room temperature, as previously described (11,18). Membrane patches were clamped at  $-50$  mV. The intracellular bath solution contained (in mmol/l) 140 KCl, 2  $CaCl_2$  (free  $[Ca^{2+}] = 50$  nmol/l), 0.7 free  $Mg^{2+}$ , 10 EGTA, and 5 HEPES (pH 7.3), and the pipette solution contained (in mmol/l) 140 KCl, 2.6  $CaCl_2$ , 1.2  $MgCl_2$ , and 10 HEPES (pH 7.4). Test media were applied directly to the patch via a PF-8 rapid perfusion system (E.S.F. Electronic, Göttingen, Germany; diameter of the application pipette 200  $\mu$ m, flux rate 1 ml/min with the tip of the patch pipette positioned in the middle of the media stream).

The stability of a patch was analyzed in a 3-min test period before the start of the experiments (presence of 1  $\mu$ mol/l MgATP). Only patches with little "run-down" over this period ( $<20\%$  loss of channel activity) were chosen. E23K did not affect the percent of patches complying with this condition (wild-type 73%, 95% CI 62–84%; E23K 76%, 95% CI 65–87%;  $n = 60$  each). Patches were corrected for "run-down" over the measuring period by use of linear interpolation. Artifacts from incomplete wash-out of nucleotides or slow reversibility were excluded by ensuring that experiments with stepwise decrease of nucleotide concentrations yielded half-maximal effective concentration ( $EC_{50}$ ) values and slope factors identical to those from experiments with a stepwise increase of the concentration. The density of  $K_{ATP}$  channels per patch ranged from 100 to 700. Variations of channel densities did not affect the percent of sensitive channels,  $EC_{50}$  values, or Hill coefficients.

**Statistical analysis.** Data analysis and statistics were performed as previously described (11,18). Results are shown as records from representative single experiments or as means  $\pm$  SE ( $n = 8-15$ ).  $EC_{50}$  values and Hill coefficients ( $n$ ) were estimated by fitting the function  $\Delta P_O = 1/(1+(nucleotide)/EC_{50})^n$  to the data of each single experiment, where  $\Delta P_O$  is the maximal


nucleotide-induced change of the  $P_O$ .  $P$  values were calculated by the Mann-Whitney  $U$  test with correction for multiple comparisons.

## RESULTS

**Modulation of ATP action.** Both ADP and guanosine 5'-diphosphate (GDP) are thought to be involved in the regulation of  $K_{ATP}$  channel activity, with resting cytosolic concentrations of  $\sim 300$  and  $\sim 60$   $\mu$ mol/l, respectively (19,20). Each of these nucleotides is a potent activator of channel activity, but GDP (up to 2 mmol/l), in contrast to ADP, does not induce channel inhibition through interaction with the  $K_{IR}6.2$  site (21). Hence, to facilitate interpretation of results, GDP was used in the experiments.

Pure wild-type channels (SUR1/ $K_{IR}6.2$ ) transiently expressed in COS cells and analyzed in inside-out patches were strongly activated by the addition of 300  $\mu$ mol/l GDP (Fig. 1A). Exposure to increasing concentrations of ATP led to complete suppression of this activity, with an  $IC_{50}$  of  $71 \pm 4.5$   $\mu$ mol/l ( $n = 15$ ; Fig. 1A and B). E23K significantly weakened this effect, inducing a 1.7-fold rightwards shift of the concentration inhibition curve ( $IC_{50} = 120 \pm 5.2$   $\mu$ mol/l;  $n = 10$ ;  $P < 0.01$ ) (Fig. 1A and B). This effect was neither reproduced by L270V or I337V nor altered by additional introduction of these polymorphisms into the  $K_{IR}6.2_{E23K}$  cDNA (Fig. 1C).

**Modulation of GDP action.** Spontaneous channel activity of pure wild-type channels was partially suppressed by the addition of 10  $\mu$ mol/l ATP (Fig. 2A). Exposure to



**FIG. 2.** Effect of E23K on GDP potency in the presence of ATP. **A:** GDP sensitivity of channels reconstituted in COS-1 cells by co-expression of human SUR1 with human K<sub>IR</sub>6.2<sub>wt</sub> or K<sub>IR</sub>6.2<sub>E23K</sub>, as indicated. Representative currents recorded at  $-50$  mV from inside-out patches exposed to GDP and ATP (10  $\mu$ mol/l), as shown by the lines above the records. Dashed lines indicate channel activation (percent) by 30  $\mu$ mol/l GDP. **B:** Potencies of GDP in the simultaneous presence of various ATP concentrations (10 and 100  $\mu$ mol/l and 1 mmol/l). Channel activation was recorded in inside-out patches as shown in **A**. Results are given as means  $\pm$  SE of 10 independent experiments. EC<sub>50</sub> values and Hill coefficients are as follows. With 10  $\mu$ mol/l ATP: wild-type ( $\blacksquare$ ),  $89 \pm 4.1$   $\mu$ mol/l and 1.33, and E23K ( $\square$ ),  $45 \pm 2.2$   $\mu$ mol/l and 1.32; with 100  $\mu$ mol/l ATP: wild-type ( $\bullet$ ),  $362 \pm 18$   $\mu$ mol/l and 1.21, and E23K ( $\circ$ )  $219 \pm 7.2$   $\mu$ mol/l and 1.23; with 1 mmol/l ATP: wild-type ( $\blacklozenge$ )  $3.29 \pm 0.15$  mmol/l and 1.24, and E23K ( $\blacklozenge$ )  $1.95 \pm 0.08$  mmol/l and 1.23.  $P < 0.01$  for comparison of the EC<sub>50</sub> values for wild-type and E23K each. **C:** The effect of E23K on the EC<sub>50</sub> value for GDP was not altered by the additional introduction of L270V and/or I337V into human K<sub>IR</sub>6.2. \* $P < 0.05$  vs. wild-type ( $n = 8-10$ ).

increasing concentrations of GDP led to reactivation, with an EC<sub>50</sub> of  $89 \pm 4.1$   $\mu$ mol/l ( $n = 10$ ) (Fig. 2A and B). E23K significantly enhanced this effect, inducing a twofold leftwards shift of the concentration activation curve for GDP (IC<sub>50</sub> =  $45 \pm 2.2$   $\mu$ mol/l;  $n = 10$ ;  $P < 0.01$ ) (Fig. 2A and B). Equivalent effects were observed when the concentration of ATP was increased to 100  $\mu$ mol/l or 1 mmol/l ( $P < 0.01$  each) (Fig. 2B). Similar to the results presented above, the effect of E23K on GDP potency was neither reproduced by L270V or I337V nor altered by the additional introduction of these polymorphisms into the K<sub>IR</sub>6.2<sub>E23K</sub> cDNA (Fig. 2C). All results obtained with the human isoforms were confirmed in channels reconstituted from mouse K<sub>IR</sub>6.2 plus hamster SUR1 (results not shown).

## DISCUSSION

The results showed that besides reducing sensitivity toward inhibitory ATP, E23K in K<sub>IR</sub>6.2 increases sensitivity of pancreatic  $\beta$ -cell K<sub>ATP</sub> channels for activation through nucleoside diphosphates. Experiments were performed in the simultaneous presence of nucleoside di- and triphosphates, because in intact cells both types of nucleotides interact in a complex manner in channel control.

Which of the two effects is physiologically relevant in regulating insulin release—either inhibition of channel activity through interaction of nucleotides with their site on K<sub>IR</sub>6.2 or channel activation through binding to the SUR1 subunit, or both? Actually, in concurrence with the conclusion that insulin secretion might be controlled by

the cytosolic ATP/ADP ratio (22), there is evidence in favor of both sites. First, in resting pancreatic  $\beta$ -cells, estimates of the cytosolic concentration of ATP range from 0.6 to 3 mmol/l (19,22–24) and those of ADP, from 0.3 to 1.2 mmol/l (19,22). Consistent with previous reports (25), our experiments show modulation of channel activity if at 300  $\mu$ mol/l GDP the ATP concentration varies  $\sim$ 1 mmol/l (Fig. 1B) and, vice versa, at 1 mmol/l ATP the GDP concentration fluctuates  $\sim$ 300  $\mu$ mol/l (Fig. 2B). Thus, it seems plausible to assume that effects at both the inhibitory and the activatory nucleotide site are involved in physiological channel control. Second, in agreement with this conclusion, loss of the activatory effect of nucleoside diphosphates results in permanently closed channels and hypersecretion of insulin, despite severe hypoglycemia (as is seen in persistent hyperinsulinemic hypoglycemia of infancy) (4,5). Third, on the other hand, the importance of the inhibitory site is underlined by the observation that reduction of ATP sensitivity by no more than 3.6-fold has been shown to induce severe neonatal diabetes in transgenic mice (3). Hence, presumably both effects of E23K—reduced ATP sensitivity and increased nucleoside diphosphate potency—are physiologically relevant.

How does the initial observation of increased spontaneous P<sub>0</sub> relate to these effects (11)? In the absence of nucleotides, E23K reduces the time spent in interburst closed states by affecting as well the frequency and mean duration of these states (11). Thus, equilibrium between interburst and burst times seems shifted toward the latter

by both, stabilizing the intraburst states (open state plus closed state) and destabilizing interburst closure. Interestingly, similar changes have been observed in nucleoside diphosphate-induced channel activation (26), which suggests that E23K exerts its effect by facilitating the conformational change that in the physiological channel control results from SUR-K<sub>IR</sub> interaction. Hence, based on channel kinetics, we would expect E23K to increase the probability that nucleoside diphosphate binding results in channel opening and thus to induce a leftwards shift of the concentration activation curve.

What about ATP? In contrast to nucleoside diphosphates, ATP does not affect burst duration, but solely increases the average time spent in interburst closed states (27,28), suggesting exclusive binding to these states. E23K-induced reduction of the time spent in interburst closure should thus lower the amount of bound ATP. Higher ATP concentrations would be expected to compensate this effect by accelerated association and, accordingly, E23K should induce a rightwards shift of the concentration-inhibition curve. Thus, the effects of the polymorphism on spontaneous channel kinetics (11) appear consistent with the changes in nucleotide sensitivities observed in this study.

Considerable nucleoside diphosphate kinase (NDPK) activity has been demonstrated in pancreatic  $\beta$ -cells (29). Specifically, the effect of GDP on glibenclamide binding in particulate and solubilized  $\beta$ -cell membranes (30,31) suggests that this enzyme is closely linked to  $\beta$ -cell K<sub>ATP</sub> channels, and therefore NDPK activity in excised patches might have led to a distortion of the results obtained in this study. Yet there are clear arguments against this deduction. First, Hill coefficients of the concentration inhibition curves for ATP in the presence of GDP were 1.22 and 1.34 for wild-type and E23K channels, respectively (Fig. 1B), which is consistent with values expected in a medium void of ADP (18). Second, a rough calculation reveals that because of rapid and direct application of the test media (see RESEARCH DESIGN AND METHODS), enzymatically formed nucleotides (ADP or GTP) would at no time be expected to reach concentrations at the excised patch higher than 0.1% of educt (ATP or GDP) concentrations.

In conclusion, in the presence of physiological nucleotide concentrations, E23K in K<sub>IR</sub>6.2 sensitizes pancreatic K<sub>ATP</sub> channels toward activatory nucleoside diphosphates, whereas the inhibitory effect of ATP is reduced. Results support the inference that the polymorphism predisposes to type 2 diabetes through changing the channel's response to physiological variations of cytosolic nucleotides, resulting in K<sub>ATP</sub> overactivity and discrete inhibition of insulin release (11).

#### ACKNOWLEDGMENTS

This work was supported by grants from the Deutsche Forschungsgemeinschaft (M.Schw. and C.S.).

We thank Haide Fürstenberg, Claudia Ott, Beate Pieper, Carolin Rattunde, and Sabine Warmbold for excellent technical assistance.

#### REFERENCES

- Polonsky KS, Sturis J, Bell GI: Non-insulin-dependent diabetes mellitus: a genetically programmed failure of the beta cell to compensate for insulin resistance. *N Engl J Med* 334:777-783, 1996

- Permutt MA, Chiu K, Ferrer J, Glaser B, Inoue H, Nestorowicz A, Stanley CA, Tanizawa Y: Genetics of type II diabetes. *Recent Prog Horm Res* 53:201-216, 1998
- Koster JC, Marshall BA, Ensor N, Corbett JA, Nichols CG: Targeted overactivity of  $\beta$ -cell K<sub>ATP</sub> channels induces profound neonatal diabetes. *Cell* 100:645-654, 2000
- Aguilar-Bryan L, Clement JPIV, Gonzalez G, Kunjilwar K, Babenko A, Bryan J: Toward understanding the assembly and structure of K<sub>ATP</sub> channels. *Physiol Rev* 78:227-245, 1998
- Aguilar-Bryan L, Bryan J: Molecular biology of adenosine triphosphate-sensitive potassium channels. *Endocr Rev* 20:101-135, 1999
- Sakura H, Wat N, Horton V, Millns H, Turner RC, Ashcroft FM: Sequence variations in the human Kir6.2 gene, a subunit of the beta-cell ATP-sensitive K-channel: no association with NIDDM in white Caucasian subjects or evidence of abnormal function when expressed in vitro. *Diabetologia* 39:1233-1236, 1996
- Inoue H, Ferrer J, Warren-Perry M, Zhang Y, Millns H, Turner RC, Elbein SC, Hampe CL, Suarez BK, Inagaki N, Seino S, Permutt MA: Sequence variants in the pancreatic islet beta-cell inwardly rectifying K<sup>+</sup> channel Kir6.2 (Bir) gene: identification and lack of role in Caucasian patients with NIDDM. *Diabetes* 46:502-507, 1997
- Hansen L, Echwald SM, Hansen T, Urhammer SA, Clausen JO, Pedersen O: Amino acid polymorphisms in the ATP-regulatable inward rectifier Kir6.2 and their relationships to glucose- and tolbutamide-induced insulin secretion, the insulin sensitivity index, and NIDDM. *Diabetes* 46:508-512, 1997
- Hani EH, Boutin P, Durand E, Inoue H, Permutt MA, Velho G, Froguel P: Missense mutations in the pancreatic islet beta cell inwardly rectifying K<sup>+</sup> channel gene (KIR6.2/BIR): a meta-analysis suggests a role in the polygenic basis of type II diabetes mellitus in Caucasians. *Diabetologia* 41:1511-1515, 1998
- Gloyn AL, Hashim Y, Ashcroft SJ, Ashfield R, Wiltshire S, Turner RC, UK Prospective Diabetes Study (UKPDS 53): Association studies of variants in promoter and coding regions of beta-cell ATP-sensitive K-channel genes SUR1 and Kir6.2 with type 2 diabetes mellitus (UKPDS 53). *Diabet Med* 18:206-212, 2001
- Schwanstecher C, Meyer U, Schwanstecher M: K<sub>IR</sub> 6.2 polymorphism predisposes to type 2 diabetes by inducing overactivity of pancreatic  $\beta$ -cell ATP-sensitive potassium channels. *Diabetes* 51:875-879, 2002
- Koster JC, Sha Q, Shyng S-L, Nichols CG: ATP inhibition of K<sub>ATP</sub> channels: control of nucleotide sensitivity by the N-terminal domain of the Kir6.2 subunit. *J Physiol* 515:19-30, 1999
- Babenko AP, Gonzalez G, Bryan J: The N-terminus of K<sub>IR</sub> 6.2 limits spontaneous bursting and modulates the ATP-inhibition of K<sub>ATP</sub> channels. *Biochem Biophys Res Commun* 255:231-238, 1999
- Ashcroft FM, Gribble FM: ATP-sensitive K<sup>+</sup> channels and insulin secretion: their role in health and disease. *Diabetologia* 42:903-919, 1999
- Panten U, Heipel C, Rosenberger F, Scheffer K, Zunkler BJ, Schwanstecher C: Tolbutamide-sensitivity of the adenosine 5'-triphosphate-dependent K<sup>+</sup> channel in mouse pancreatic B-cells. *Naunyn Schmiedebergs Arch Pharmacol* 342:566-574, 1990
- Trapp S, Tucker SJ, Ashcroft FM: Activation and inhibition of K-ATP currents by guanine nucleotides is mediated by different channel subunits. *Proc Natl Acad Sci U S A* 94:8872-8877, 1997
- Kakei M, Kelly RP, Ashcroft SJ, Ashcroft FM: The ATP-sensitivity of K<sup>+</sup> channels in rat pancreatic B-cells is modulated by ADP. *FEBS Lett* 208:63-66, 1986
- Markworth E, Schwanstecher C, Schwanstecher M: ATP<sup>4-</sup> mediates closure of pancreatic beta-cell ATP-sensitive potassium channels by interaction with 1 of 4 identical sites. *Diabetes* 49:1413-1418, 2000
- Malaisse WJ, Sener A: Glucose-induced changes in cytosolic ATP content in pancreatic islets. *Biochim Biophys Acta* 927:190-195, 1987
- Detimary P, Van den Berghe G, Henquin JC: Concentration dependence and time course of the effects of glucose on adenine and guanine nucleotides in mouse pancreatic islets. *J Biol Chem* 271:20559-20565, 1996
- Schwanstecher C, Dickel C, Panten U: Interaction of tolbutamide and cytosolic nucleotides in controlling the ATP-sensitive K<sup>+</sup> channel in mouse beta-cells. *Br J Pharmacol* 111:302-310, 1994
- Detimary P, Dejonghe S, Ling Z, Pipeleers D, Schuit F, Henquin J-C: The changes in adenine nucleotides measured in glucose-stimulated rodent islets occur in  $\beta$  cells but not in  $\alpha$  cells and are observed in human islets. *J Biol Chem* 273:33905-33908, 1998
- Kennedy HJ, Pouli AE, Ainscow EK, Jouaville LS, Rizzuto R, Rutter GA: Glucose generates sub-plasma membrane ATP microdomains in single islet  $\beta$ -cells: potential role for strategically located mitochondria. *J Biol Chem* 274:13281-13291, 1999
- Gribble FM, Loussouarn G, Tucker SJ, Zhao C, Nichols CG, Ashcroft FM:

- A novel method for measurement of submembrane ATP concentration. *J Biol Chem* 275:30046–30049, 2000
25. Dunne MJ, West-Jordan JA, Abraham RJ, Edwards RH, Petersen OH: The gating of nucleotide-sensitive  $K^+$  channels in insulin-secreting cells can be modulated by changes in the ratio  $ATP^4^-/ADP^3^-$  and by nonhydrolyzable derivatives of both ATP and ADP. *J Membr Biol* 104:165–177, 1988
  26. Branstrom R, Corkey BE, Berggren PO, Larsson O: Evidence for a unique long chain acyl-CoA ester binding site on the ATP-regulated potassium channel in mouse pancreatic beta cells. *J Biol Chem* 272:17390–17394, 1997
  27. Enkvetchakul D, Loussouarn G, Makhina E, Shyng SL, Nichols CG: The kinetic and physical basis of  $K_{ATP}$  channel gating: toward a unified molecular understanding. *Biophys J* 78:2334–2348, 2000
  28. Enkvetchakul D, Loussouarn G, Makhina E, Nichols CG: ATP interaction with the open state of the  $K(ATP)$  channel. *Biophys J* 80:719–728, 2001
  29. Kowluru A, Metz SA: Characterization of nucleoside diphosphokinase activity in human and rodent pancreatic beta cells: evidence for its role in the formation of guanosine triphosphate, a permissive factor for nutrient-induced insulin secretion. *Biochemistry* 33:12495–12503, 1994
  30. Schwanstecher M, Löser S, Rietze I, Panten U: Phosphate and thiophosphate group donating adenine and guanine nucleotides inhibit glibenclamide binding to membranes from pancreatic islets. *Naunyn-Schmiedeberg's Arch Pharmacol* 343:83–89, 1991
  31. Schwanstecher M, Behrends S, Brandt C, Panten U: The binding properties of the solubilized sulfonylurea receptor from a pancreatic B-cell line are modulated by the  $Mg^{++}$ -complex of ATP. *J Pharmacol Exp Ther* 262:495–502, 1992