

PYY₃₋₃₆ Reinforces Insulin Action on Glucose Disposal in Mice Fed a High-Fat Diet

Anita M. van den Hoek,^{1,2} Annemieke C. Heijboer,^{1,3} Eleonora P.M. Corssmit,³ Peter J. Voshol,^{1,3} Johannes A. Romijn,³ Louis M. Havekes,^{1,2,4} and Hanno Pijl³

Peptide YY₃₋₃₆ (PYY₃₋₃₆) is released by the gut in response to nutrient ingestion. It modulates the activities of orexigenic neuropeptide Y (NPY) neurons and anorexigenic proopiomelanocortin (POMC) neurons in the hypothalamus to inhibit food intake. Because both NPY and POMC have also been shown to impact insulin action, we wondered whether PYY₃₋₃₆ could improve insulin sensitivity. To address this question, we examined the acute effect of intravenous PYY₃₋₃₆ on glucose and free fatty acid (FFA) flux during a hyperinsulinemic-euglycemic clamp in mice maintained on a high-fat diet for 2 weeks before the experiment. We also evaluated the effects of PYY₃₋₃₆ infusion on glucose uptake in muscle and adipose tissue in this experimental context. Under basal conditions, none of the metabolic parameters were affected by PYY₃₋₃₆. Under hyperinsulinemic conditions, glucose disposal was significantly increased in PYY₃₋₃₆-infused compared with vehicle-infused mice (103.8 ± 10.9 vs. $76.1 \pm 11.4 \mu\text{mol} \cdot \text{min}^{-1} \cdot \text{kg}^{-1}$, respectively; $P = 0.001$). Accordingly, glucose uptake in muscle and adipose tissue was greater in PYY₃₋₃₆-treated animals, although the difference with controls did not reach statistical significance in adipose tissue (muscle: 2.1 ± 0.5 vs. $1.5 \pm 0.5 \mu\text{mol/g tissue}$, $P = 0.049$; adipose tissue: 0.8 ± 0.4 vs. $0.4 \pm 0.3 \mu\text{mol/g tissue}$, $P = 0.08$). In contrast, PYY₃₋₃₆ did not impact insulin action on endogenous glucose production or FFA metabolism. These data indicate that PYY₃₋₃₆ reinforces insulin action on glucose disposal in mice fed a high-fat diet, through a mechanism that is independent of food intake and body weight. In contrast, it leaves glucose production and lipid flux largely unaffected in this experimental context. *Diabetes* 53:1949–1952, 2004

Peptide YY (PYY) belongs to a family of peptides that is critically involved in the regulation of appetite. It is synthesized in specialized cells (L-cells) that are found primarily in the distal gastrointestinal tract. Circulating PYY levels rise within 15 min after a meal in proportion to the amount of calories ingested and remain elevated for ~6 h (1). The two natural forms of this peptide, PYY₁₋₃₆ and PYY₃₋₃₆, have opposing effects on food intake (2): PYY₁₋₃₆ stimulates appetite, whereas PYY₃₋₃₆ (the major circulating form) inhibits feeding (3–5).

PYY₃₋₃₆ reduces food intake by acting on appetite regulation centers in the hypothalamus (3,6). Specifically, PYY₃₋₃₆ is an agonist of the presynaptic neuropeptide Y (NPY) Y2 receptor, which inhibits NPY neuronal activity in the arcuate nucleus and thereby activates adjacent proopiomelanocortin (POMC) neurons (3). In addition to their critical role in the control of feeding, both NPY and POMC affect insulin action. Intracerebroventricular infusion of NPY induces hyperinsulinemia and insulin resistance in rats (7,8). In contrast, central administration of α -melanocyte-stimulating hormone, a splice product of the POMC peptide, enhances insulin sensitivity (9). In view of the fact that PYY₃₋₃₆ inhibits NPY neuronal activity and activates that of POMC, we wondered whether it could improve insulin sensitivity “directly” (i.e., through a mechanism independent of food intake and body weight). To address this question, we infused PYY₃₋₃₆ intravenously and quantified its acute effects on glucose and fatty acid flux during a hyperinsulinemic-euglycemic clamp in mice without access to food during the experiment.

RESEARCH DESIGN AND METHODS

Male C57BL/6J mice were housed in a temperature-controlled room on a 12-h light/dark cycle and were fed a high-fat diet (43 energy% fat derived from bovine lard) with free access to water for 2 weeks to induce insulin resistance. All animal experiments were performed in accordance with the regulations of Dutch law on animal welfare, and the institutional ethics committee for animal procedures approved the protocol.

Hyperinsulinemic-euglycemic clamp. A total of 36 mice were fasted overnight with food withdrawn at 5:00 P.M. the day before the study. The next day, hyperinsulinemic-euglycemic clamps were performed as described earlier (10). PYY or vehicle was administered as a primed (0.15 μg)-continuous (0.25 $\mu\text{g/h}$) intravenous infusion during the whole experiment (basal and hyperinsulinemic period). In one series of experiments, glucose turnover was quantified, and in another series, free fatty acid (FFA) turnover was determined. First, basal rates of glucose or FFA turnover were measured by giving a primed-continuous infusion of ³H-glucose (prime: 0.7 μCi ; continuous: 1.2 $\mu\text{Ci/h}$; Amersham, Little Chalfont, U.K.) or ¹⁴C-palmitate (prime: 1.8 μCi ; continuous: 3 $\mu\text{Ci/h}$; Amersham), respectively, for 80 min. Subsequently, insulin was administered in a primed (4.1 mU)-continuous (6.8 mU/h) intra-

From ¹TNO Prevention and Health, Gaubius Laboratory, Leiden, the Netherlands; the ²Department of Internal Medicine, Leiden University Medical Center, Leiden, the Netherlands; the ³Department of Endocrinology and Metabolic Diseases, Leiden University Medical Center, Leiden, the Netherlands; and the ⁴Department of Cardiology, Leiden University Medical Center, Leiden, the Netherlands.

Address correspondence and reprint requests to Hanno Pijl, MD, Leiden University Medical Center, Department of Endocrinology and Metabolic Diseases, P.O. Box 9600, 2300 RC Leiden, Netherlands. E-mail: h.pijl@lumc.nl

Received for publication 31 March 2004 and accepted in revised form 12 May 2004.

A.M.V.D.H. and A.C.H. contributed equally to this work.

2-DG, 2-deoxy-D-[³H]glucose; EGP, endogenous glucose production; FFA, free fatty acid; NPY, neuropeptide Y; POMC, proopiomelanocortin; PYY, peptide YY.

© 2004 by the American Diabetes Association.

TABLE 1

Plasma parameters in overnight-fasted mice that received an intravenous infusion of PYY or vehicle under basal or hyperinsulinemic conditions

	Basal		Hyperinsulinemic	
	Vehicle	PYY	Vehicle	PYY
Glucose (mmol/l)	4.9 ± 0.6	5.0 ± 0.6	9.0 ± 0.9	9.2 ± 2.0
FFAs (mmol/l)	0.9 ± 0.2	1.0 ± 0.2	0.4 ± 0.1	0.5 ± 0.1
Insulin (ng/ml)	0.3 ± 0.1	0.3 ± 0.1	5.2 ± 2.9	6.7 ± 2.7
PYY ₃₋₃₆ (pg/μl)	<1	3.8 ± 0.7	<1	3.2 ± 0.7

Data are the means ± SD for at least seven mice per group.

venous infusion for 2–3 h to attain steady-state circulating insulin levels of ~6 ng/ml. A variable infusion of 12.5% D-glucose was used to maintain euglycemia, measured at 10-min intervals via tail bleeding (Freestyle; TheraSense, Diabetic Medical Systems, Vianen, the Netherlands). Blood samples (75 μl) were taken during the basal period (after 60 and 80 min) and during the clamp period (when glucose levels were stable and 20 and 40 min later) for determination of plasma glucose, FFA, and insulin concentrations and [³H]glucose and [¹⁴C]palmitate specific activities.

To assess insulin-mediated glucose uptake in individual tissues, 2-deoxy-D-[³H]glucose (2-DG; Amersham) was administered as a bolus (1 μCi) 40 min before the end of the clamp experiments in which FFA turnover was measured. At the end of the clamp, mice were killed, and muscle and adipose tissue were isolated and frozen in liquid nitrogen for subsequent analysis.

Analytical procedures. Plasma levels of glucose and FFAs were determined using commercially available kits (InstruChemie, Delfzijl, the Netherlands, and Wako, Neuss, Germany). Plasma insulin and PYY₃₋₃₆ concentrations were measured by a mouse insulin enzyme-linked immunosorbent assay and PYY₃₋₃₆ radioimmunoassay (Alpco, Windham, NH, and Phoenix Pharmaceuticals, Belmont, CA). Total plasma ³H-glucose was determined in 7.5 μl plasma and in supernatants after trichloroacetic acid (20%) precipitation and water evaporation to eliminate tritiated water. Total plasma ¹⁴C-palmitate was determined in 7.5 μl plasma after extraction of lipids by a modification of Bligh and Dyer's (11) method. Briefly, 7.5 μl plasma was dried and resuspended in 100 μl water. Then, 1.1 ml demi-water and 4.5 ml methanol:chloroform (2:1) were added and mixed thoroughly, after which 1.5 ml chloroform was added and mixed, and, finally, 1.5 ml demi-water was added and mixed. After centrifugation, the chloroform layer was collected, and the FFA fraction was separated from other lipid components by thin-layer chromatography on silica gel plates.

Tissue analysis. For determination of tissue 2-DG uptake, the homogenate of muscle and adipose tissue was boiled, and the supernatant was subjected to an ion-exchange column to separate 2-DG-6-phosphate from 2-DG as described previously (10,12,13).

Calculations. Turnover rates of glucose and FFAs (μmol · min⁻¹ · kg⁻¹) were calculated during the basal period and in steady-state clamp conditions as the rate of tracer infusion (dpm/min) divided by the plasma specific activity of ³H-glucose or ¹⁴C-palmitate (dpm/μmol). The ratio was corrected for body weight. Endogenous glucose production (EGP) was calculated as the difference between the tracer-derived rate of glucose appearance and the glucose infusion rate.

Tissue-specific glucose uptake in muscle and adipose tissue was calculated from tissue 2-DG content, corrected for plasma specific activity and expressed as micromoles per gram of tissue.


Statistical analysis. Differences between groups were determined by Mann-Whitney's nonparametric test for two independent samples. *P* < 0.05 was considered statistically significant. All values shown represent means ± SD.

RESULTS

Animals. Mice were 16–18 weeks old during the experiments. Body weight was 23.3 ± 1.2 g in the control group and 22.8 ± 1.4 g in the PYY group.

Plasma parameters. Plasma glucose, FFA, insulin, and PYY₃₋₃₆ concentrations in basal and hyperinsulinemic conditions are shown in Table 1. In the basal state, plasma parameters did not differ between PYY- and vehicle-infused animals, except for the plasma PYY₃₋₃₆ concentration. Under steady-state clamp conditions, plasma insulin and glucose concentrations were similar in both groups.

A



B




FIG. 1. Glucose disposal (A) and EGP (B) in overnight-fasted mice before (basal) and during (hyperinsulinemic) a hyperinsulinemic-euglycemic clamp. Before and during insulin infusion, the animals received an intravenous infusion of PYY or vehicle. Values represent the means ± SD for at least seven mice per group. **P* < 0.01 vs. vehicle. ■, vehicle; □, PYY.

Hyperinsulinemia suppressed FFA levels to a similar extent in PYY- and vehicle-infused mice.

Glucose turnover. Under basal conditions, glucose disposal was similar in PYY- and vehicle-infused mice (39.5 ± 10.9 vs. 45.9 ± 12.6 μmol · min⁻¹ · kg⁻¹, respectively; *P* = 0.19) (Fig. 1A). The rate of glucose infusion necessary to maintain euglycemia during insulin infusion was significantly higher in PYY-infused mice than in vehicle-infused animals (89.1 ± 7.1 vs. 50.9 ± 13.6 μmol · min⁻¹ · kg⁻¹, *P* = 0.001), indicating that intravenous PYY₃₋₃₆ administration acutely enhances insulin sensitivity. Hyperinsulinemia increased glucose disposal in both groups. However, the disposal rate was significantly higher in PYY-infused animals compared with vehicle-infused controls (103.8 ± 10.9 vs. 76.1 ± 11.4 μmol · min⁻¹ · kg⁻¹, respectively; *P* = 0.001) (Fig. 1A). In contrast, EGP, which was similar in PYY- and vehicle-treated mice under basal conditions, was suppressed by insulin to the same extent in both groups (by 62 ± 29 vs. 42 ± 18% from basal, respectively; *P* = 0.30) (Fig. 1B).




FIG. 2. Muscle-specific (A) and adipose tissue-specific (B) glucose uptake under hyperinsulinemic conditions in overnight-fasted mice that received an intravenous infusion of PYY or vehicle. Values represent the means \pm SD for at least seven mice per group. * $P < 0.05$ vs. vehicle. ■, vehicle; □, PYY.

Tissue-specific glucose uptake. Insulin-mediated 2-DG uptake in muscle and adipose tissue was greater in PYY-treated animals, but the difference with controls did not reach statistical significance in adipose tissue (muscle: 2.1 ± 0.5 vs. 1.5 ± 0.5 $\mu\text{mol/g}$ tissue, $P = 0.049$; adipose tissue: 0.8 ± 0.4 vs. 0.4 ± 0.3 $\mu\text{mol/g}$ tissue, $P = 0.08$) (Fig. 2).

FFA turnover. Basal FFA turnover (38.0 ± 14.2 vs. 42.3 ± 9.9 $\mu\text{mol} \cdot \text{min}^{-1} \cdot \text{kg}^{-1}$) was not different between PYY- and vehicle-infused animals and was suppressed to a similar extent by insulin in both groups (22.4 ± 12.3 vs. 21.3 ± 10.9 $\mu\text{mol} \cdot \text{min}^{-1} \cdot \text{kg}^{-1}$ in PYY- and vehicle-infused animals, respectively) (Fig. 3).

DISCUSSION

Here we show that intravenous PYY₃₋₃₆ administration acutely reinforces insulin action on glucose disposal in overnight-fasted mice maintained on a high-fat diet. In contrast, PYY₃₋₃₆ does not appear to impact the effect of insulin on EGP or FFA metabolism in this experimental context.

PYY₃₋₃₆ clearly enhanced insulin-induced glucose disposal as determined by tracer dilution methodology. Ac-




FIG. 3. FFA turnover in overnight-fasted mice before (basal) and during (hyperinsulinemic) a hyperinsulinemic-euglycemic clamp. Before and during insulin infusion, the animals received an intravenous infusion of PYY or vehicle. Values represent the means \pm SD for at least nine mice per group. ■, vehicle; □, PYY.

cordingly, 2-DG uptake in muscle and adipose tissue under hyperinsulinemic conditions was higher during PYY₃₋₃₆ infusion, albeit the difference with control attained statistical significance only for muscle. In contrast, PYY₃₋₃₆ did not significantly impact the capacity of insulin to inhibit EGP. Insulin action on FFA metabolism was not affected by PYY₃₋₃₆, either, as indicated by similar fatty acid turnover rates during hyperinsulinemia in PYY₃₋₃₆- and saline-infused animals. In light of the current experimental context, these data suggest that PYY₃₋₃₆ reinforces the impact of insulin on glucose disposal through a mechanism that is independent of food intake and body weight. In contrast, it appears to leave insulin action on glucose production and FFA turnover largely unaffected.

The plasma PYY₃₋₃₆ concentration rose to 3.2 ± 0.7 pg/ μl in response to PYY infusion. Relatively few studies have looked at the physiology of circulating PYY₃₋₃₆ in rodents. According to a recent article by Batterham et al. (3), postprandial PYY₃₋₃₆ concentrations amount to 112 pmol/l (~ 0.4 pg/ μl) in freely feeding rats. In a study by Lee et al. (14), plasma PYY levels in fasting mice were 0.18 pg/ml, which accords with our data (Table 1). We are not aware of any study measuring postprandial PYY₃₋₃₆ concentrations in mice. Thus, the dose of PYY₃₋₃₆ we administered may have induced supraphysiological PYY₃₋₃₆ levels. Therefore, our data do not allow a definitive inference as to whether the postprandial rise of circulating PYY₃₋₃₆ can reinforce insulin action.

Postprandial PYY₃₋₃₆ release is decreased in obese compared with lean subjects, and circulating levels correlate negatively with BMI. Conversely, intravenous PYY₃₋₃₆ infusion significantly reduces food intake in humans (15), and repeated administration of PYY₃₋₃₆ attenuates weight gain in rodents (3). These findings suggest that diminished PYY₃₋₃₆ release may contribute to the pathogenesis of obesity in animals and humans. The observations presented here allow us to hypothesize that low circulating PYY₃₋₃₆ levels may also compromise insulin action in obese subjects. Moreover, perhaps even more important, the data suggest that PYY₃₋₃₆ or synthetic analogs of this peptide may be useful tools in the clinical management of obesity and insulin resistance.

It remains to be established whether PYY₃₋₃₆ acts through hypothalamic neural circuits (by analogy with the mechanism guiding its effects on appetite) to enhance insulin-mediated glucose disposal. Because PYY₃₋₃₆ is a high-affinity agonist to the Y2 receptor (2) and the distribution of this receptor subtype is largely confined to the central nervous system (particularly the arcuate nucleus of the hypothalamus) (16), it is most likely that PYY₃₋₃₆ modulates insulin action by activation of Y2 receptors in the brain. As alluded to earlier, Y2 receptor signaling inhibits NPY neuronal activity in the arcuate nucleus of the hypothalamus and thereby activates adjacent POMC neurons (3). Hypothalamic overexpression of NPY induces obesity and insulin resistance in mice (7,8), whereas activation of melanocortin receptor subtypes 3 and 4 in the brain enhances insulin sensitivity (9). Thus, the present data are in keeping with the contention that PYY₃₋₃₆ modulates insulin action via hypothalamic Y2 receptor. The downstream mechanisms that actuate the effects of hypothalamic neuronal circuits on muscle and adipose tissue remain to be fully elucidated. At this point, vagotomy was shown to prevent the hyperinsulinemic effects of NPY, which suggests that the autonomic nervous system may be involved (17). Also, adrenalectomy prevents the obesity syndrome produced by chronic central NPY infusion and reverses the obese phenotype in leptin-deficient *ob/ob* mice (18,19), indicating that the pituitary-adrenal ensemble may also serve as a second messenger. Thus, neural and/or endocrine mechanistic routes may convey signals from hypothalamic nuclei to peripheral tissues to orchestrate energy balance and fuel flux. It is conceivable that similar routes partake in the control of insulin action by PYY₃₋₃₆.

In conclusion, PYY₃₋₃₆ reinforces insulin action in mice maintained on a high-fat diet through a mechanism that is independent of food intake and body weight. In this context, PYY₃₋₃₆ appears to predominantly impact insulin-mediated glucose disposal, whereas it leaves insulin action on glucose production largely unaffected. These novel data suggest that PYY₃₋₃₆ or synthetic analogs of this peptide may be valuable assets to our armamentarium of drugs designed to battle insulin resistance and type 2 diabetes.

ACKNOWLEDGMENTS

The research described in this article is supported by the Dutch Scientific Research Council/Netherlands Heart Foundation (Projects 980-10-017 and 907-00-002).

This study is conducted in the framework of the Leiden Center for Cardiovascular Research LUMC-TNO.

REFERENCES

1. Adrian TE, Ferri GL, Bacarese-Hamilton AJ, Fuessl HS, Polak JM, Bloom SR: Human distribution and release of a putative new gut hormone, peptide YY. *Gastroenterology* 89:1070-1077, 1985
2. Grandt D, Schimiczek M, Beglinger C, Layer P, Goebell H, Eysselein VE, Reeve JR Jr: Two molecular forms of peptide YY (PYY) are abundant in human blood: characterization of a radioimmunoassay recognizing PYY 1-36 and PYY 3-36. *Regul Pept* 51:151-159, 1994
3. Batterham RL, Cowley MA, Small CJ, Herzog H, Cohen MA, Dakin CL, Wren AM, Brynes AE, Low MJ, Ghatei MA, Cone RD, Bloom SR: Gut hormone PYY(3-36) physiologically inhibits food intake. *Nature* 418:650-654, 2002
4. Morley JE, Levine AS, Grace M, Kneip J: Peptide YY (PYY), a potent orexigenic agent. *Brain Res* 341:200-203, 1985
5. Hagan MM, Moss DE: Effect of peptide YY (PYY) on food-associated conflict. *Physiol Behav* 58:731-735, 1995
6. Challis BG, Pinnock SB, Coll AP, Carter RN, Dickson SL, O'Rahilly S: Acute effects of PYY3-36 on food intake and hypothalamic neuro peptide expression in the mouse. *Biochem Biophys Res Commun* 311:915-919, 2003
7. Marks JL, Waite K: Intracerebroventricular neuro peptide Y acutely influences glucose metabolism and insulin sensitivity in the rat. *J Neuroendocrinol* 9:99-103, 1997
8. Zarjevski N, Cusin I, Vettor R, Rohner-Jeanrenaud F, Jeanrenaud B: Chronic intracerebroventricular neuro peptide-Y administration to normal rats mimics hormonal and metabolic changes of obesity. *Endocrinology* 133:1753-1758, 1993
9. Obici S, Feng Z, Tan J, Liu L, Karkanas G, Rossetti L: Central melanocortin receptors regulate insulin action. *J Clin Invest* 108:1079-1085, 2001
10. Voshol PJ, Jong MC, Dahlmans VE, Kratky D, Levak-Frank S, Zechner R, Romijn JA, Havekes LM: In muscle-specific lipoprotein lipase-overexpressing mice, muscle triglyceride content is increased without inhibition of insulin-stimulated whole-body and muscle-specific glucose uptake. *Diabetes* 50:2585-2590, 2001
11. Bligh EG, Dyer WJ: A rapid method of total lipid extraction and purification. *Can J Biochem Biophys* 37:911-917, 1959
12. Rossetti L, Giaccari A: Relative contribution of glycogen synthesis and glycolysis to insulin-mediated glucose uptake: a dose-response euglycemic clamp study in normal and diabetic rats. *J Clin Invest* 85:1785-1792, 1990
13. Goudriaan JR, Dahlmans VE, Teusink B, Ouwens DM, Febbraio M, Maassen JA, Romijn JA, Havekes LM, Voshol PJ: CD36 deficiency increases insulin sensitivity in muscle, but induces insulin resistance in the liver in mice. *J Lipid Res* 44:2270-2277, 2003
14. Lee HM, Udipi V, Englander EW, Rajaraman S, Coffey RJ Jr, Greeley GH Jr: Stimulatory actions of insulin-like growth factor-I and transforming growth factor-alpha on intestinal neurotensin and peptide YY. *Endocrinology* 140:4065-4069, 1999
15. Batterham RL, Cohen MA, Ellis SM, Le Roux CW, Withers DJ, Frost GS, Ghatei MA, Bloom SR: Inhibition of food intake in obese subjects by peptide YY3-36. *N Engl J Med* 349:941-948, 2003
16. Parker E, Van Heek M, Stamford A: Neuropeptide Y receptors as targets for anti-obesity drug development: perspective and current status. *Eur J Pharmacol* 440:173-187, 2002
17. Sainsbury A, Rohner-Jeanrenaud F, Cusin I, Zakrzewska KE, Halban PA, Gaillard RC, Jeanrenaud B: Chronic central neuro peptide Y infusion in normal rats: status of the hypothalamo-pituitary-adrenal axis, and vagal mediation of hyperinsulinaemia. *Diabetologia* 40:1269-1277, 1997
18. Sainsbury A, Cusin I, Rohner-Jeanrenaud F, Jeanrenaud B: Adrenalectomy prevents the obesity syndrome produced by chronic central neuro peptide Y infusion in normal rats. *Diabetes* 46:209-214, 1997
19. Makimura H, Mizuno TM, Roberts J, Silverstein J, Beasley J, Mobbs CV: Adrenalectomy reverses obese phenotype and restores hypothalamic melanocortin tone in leptin-deficient *ob/ob* mice. *Diabetes* 49:1917-1923, 2000