

Subcutaneous Oxyntomodulin Reduces Body Weight in Overweight and Obese Subjects

A Double-Blind, Randomized, Controlled Trial

Katie Wynne,¹ Adrian J. Park,¹ Caroline J. Small,¹ Michael Patterson,¹ Sandra M. Ellis,² Kevin G. Murphy,¹ Alison M. Wren,¹ Gary S. Frost,² Karim Meeran,¹ Mohammad A. Ghatei,¹ and Stephen R. Bloom¹

This study investigated the effect of subcutaneously administered oxyntomodulin on body weight in healthy overweight and obese volunteers. Participants self-administered saline or oxyntomodulin subcutaneously in a randomized, double-blind, parallel-group protocol. Injections were self-administered for 4 weeks, three times daily, 30 min before each meal. The volunteers were asked to maintain their regular diet and level of physical exercise during the study period. Subjects' body weight, energy intake, and levels of adipose hormones were assessed at the start and end of the study. Body weight was reduced by 2.3 ± 0.4 kg in the treatment group over the study period compared with 0.5 ± 0.5 kg in the control group ($P = 0.0106$). On average, the treatment group had an additional 0.45-kg weight loss per week. The treatment group demonstrated a reduction in leptin and an increase in adiponectin. Energy intake by the treatment group was significantly reduced by 170 ± 37 kcal ($25 \pm 5\%$) at the initial study meal ($P = 0.0007$) and by 250 ± 63 kcal ($35 \pm 9\%$) at the final study meal ($P = 0.0023$), with no change in subjective food palatability. Oxyntomodulin treatment resulted in weight loss and a change in the levels of adipose hormones consistent with a loss of adipose tissue. The anorectic effect was maintained over the 4-week period. Oxyntomodulin represents a potential therapy for obesity. *Diabetes* 54:2390–2395, 2005

From the ¹Department of Metabolic Medicine, Imperial College Faculty of Medicine, Hammersmith Hospital, London, U.K.; and the ²Department of Dietetics, Imperial College Faculty of Medicine, Hammersmith Hospital, London, U.K.

Address correspondence and reprint requests to Prof. Stephen R. Bloom, Endocrine Unit, Imperial College Faculty of Medicine, Hammersmith Hospital, Du Cane Road, London W12 0NN, U.K. E-mail: s.bloom@imperial.ac.uk.

Received for publication 15 April 2005 and accepted in revised form 17 May 2005.

K.W. and A.J.P. contributed jointly to this study.

OLI, oxyntomodulin-like activity.

© 2005 by the American Diabetes Association.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

The prevalence of obesity is rapidly increasing worldwide; currently >65% of adults in the U.S. are overweight (1). Although even a modest weight loss can improve the health of obese individuals, efforts to treat the obesity pandemic have been unsuccessful. Novel therapeutic targets are urgently required.

Several gut hormones have been found to modulate appetite (2–4). Oxyntomodulin is a peptide product of the proglucagon gene released from the L-cells of the small intestine in response to food ingestion (5). Oxyntomodulin has been reported to reduce food intake by 19.3% during an intravenous infusion administered to normal-weight humans, an effect that continues for >12 h after infusion (6). Furthermore, in rodents, repeated intraperitoneal administration over 7 days has been associated with reduced white adipose tissue and a highly significant reduction in weight compared with controls (7). Thus, oxyntomodulin may offer a novel treatment for human obesity.

We hypothesized that self-administered subcutaneous oxyntomodulin would induce weight loss, reduce appetite, and alter the levels of adipose hormones in overweight and obese volunteers investigated in a 4-week community-based study.

RESEARCH DESIGN AND METHODS

Healthy male and female volunteers, ages 18–55 years, with a stable BMI of 25–40 kg/m² were recruited by advertisement and followed between January and August 2004. All potential subjects were nonsmokers with normal physical examination, routine blood tests, and electrocardiograms. Subjects were screened using the standard Dutch Eating Behavior Questionnaire (8) and SCOFF Questionnaire (9) and were excluded if they demonstrated abnormal eating behavior. Food preferences were assessed using a nine-point hedonistic scale to ensure that the study meal was acceptable. Subjects were given a subcutaneous injection of saline to assess whether they would find regular injections acceptable. A power calculation was used to estimate the number of subjects required to detect a significant effect of oxyntomodulin on a participant's body weight. Assuming a difference of 1 ± 0.2 kg to be clinically significant and using a significance level of $P < 0.05$ and 90% power, it was calculated that 12 subjects were required in each group; the final treatment group consisted of 14 subjects, and the final control group consisted of 12 subjects. Women of child-bearing age were advised to avoid pregnancy by using adequate contraception and were asked to provide regular urine tests to ensure they were not pregnant. The participants completed the study protocol

TABLE 1
Baseline parameters of the participants

	Control group	Treatment group	<i>P</i>
<i>n</i>	12	14	—
BMI (kg/m ²)	32.8 ± 0.9	33.7 ± 0.9	0.467
Age (years)	35.0 ± 2.0	32.5 ± 1.8	0.3573
Height (m)	1.64 ± 0.03	1.67 ± 0.02	0.3453
Weight (kg)	89.0 ± 4.7	94.6 ± 3.4	0.3300
Female/male	11/1	11/3	—
Leptin (ng/ml)	283 ± 24	325 ± 50	0.4561
Adiponectin (μg/ml)	111.4 ± 11.3	83.6 ± 9.5	0.0707
Insulin (pmol/l)	26.7 ± 8.5	34.5 ± 4.7	0.4151
Glucose (mmol/l)	4.61 ± 0.09	4.76 ± 0.11	0.3430
Cholesterol (mmol/l)	4.71 ± 0.26	4.66 ± 0.19	0.8820
HDL (mmol/l)	1.07 ± 0.06	1.12 ± 0.09	0.6747
Ghrelin (pmol/l)	493 ± 87	371 ± 65	0.2627
Peptide YY (pmol/l)	11.8 ± 1.6	12.2 ± 2.1	0.8881

Data are means ± SE. An unpaired *t* test was used to compare values between the two groups.

in the community and also attended the Charing Cross Hospital clinical trials unit at the start and end of the study for assessment of energy intake.

Ethical approval was obtained from the Riverside Medical Ethics Committee (reference no. 3397). All subjects gave informed written consent, and the study was performed in accordance with the Declaration of Helsinki.

The study followed a double-blind, placebo-controlled, parallel-group protocol designed to investigate the effect of oxyntomodulin on body weight, food intake, and adipose hormones. Oxyntomodulin was synthesized by Bachem U.K., sterile for culture and negative for pyrogen, as previously described (10). A subcutaneous dose of 400 nmol was chosen, which allowed us to achieve the same blood concentration as a previous human intravenous infusion study that demonstrated a reduction in calorie intake (6).

From recruitment, subjects were instructed to continue their current level of exercise and usual diet in a regimen of three meals per day. For 1 week before the study, all subjects self-administered saline injections three times daily, 30 min preprandially, into their abdominal subcutaneous tissue. All subjects prepared their injections by adding 0.25 ml sterile water to vials containing freeze-dried sodium chloride and self-administered injections using a 27-gauge needle. Subjects who failed to comply with the protocol during this week were removed from the study.

Subjects were randomly allocated to the control or treatment group by an independent investigator. Randomization was performed by stratification using the parameters BMI and age; subsequent randomization into the control or treatment group was performed by coin toss. The control and treatment groups were therefore matched for BMI and age and had similar mean height and weight measurements (Table 1). The estimated energy requirements for the control and treatment groups were not significantly different.

During the 4-week study period, subjects self-administered normal saline or oxyntomodulin (400 nmol), three times daily, 30 min preprandially. An identical protocol for injection preparation was used as in the previous week, and the saline and oxyntomodulin vials were indistinguishable. The subjects were reviewed weekly by blinded investigators to measure body weight, check injection technique, and monitor subjects for side effects and compliance. Adverse events during the injection period were recorded by diary. Protocol compliance was 94.4 ± 1.1% in the control group and 95.3 ± 1.2% in the treatment group, estimated by counting the number of returned empty vials. Compliance was also confirmed by checking for cutaneous sites of injection and inspecting the number of subjects' accumulated used needles. Subjects were weighed on each visit, and three fasting weights were recorded on three separate days at both the start and end of the study using a calibrated electronic scale (Marsden, London, U.K.) accurate to the nearest 100 g. Subjects wore the same light clothing and voided urine before each measurement. Preinjection blood samples were taken at the start and end of the study to measure leptin and adiponectin levels.

Energy intake was measured during three lunchtime study meals on days 1, 2, and 29. The day before each study meal, subjects refrained from alcohol and strenuous exercise and fasted overnight. Subjects first ingested a fixed 200-kcal breakfast and 210 min later self-administered their injection (designated as *t* = 0). A meal of known energy content was provided in excess quantity 30 min later, and subjects were requested to eat until they felt comfortably full. On day 1, all subjects administered a saline injection. On

days 2 and 29, the control group administered saline and the treatment group administered oxyntomodulin. Visual analog scales, 100 mm in length, were completed throughout the study meal day to evaluate subjective feelings of hunger, nausea, and meal palatability.

Blood samples were taken at *t* = -30, 0, 15, 30, 60, 90, 150, 210, and 240 min to screen for any effect of oxyntomodulin on other analytes (Table 3). All blood samples were collected from an antecubital fossa cannula into lithium/heparin tubes (LIP, Cambridge, U.K.) containing 2,000 kallikrein inhibitor units of aprotinin (Trasylol; Bayer) and stored on ice. After centrifugation, plasma was immediately separated and stored at -20°C until being analyzed. Blood pressure and pulse were measured before each blood sample on each of the study meal days.

Hormone measurements. Commercially available assays were used to measure plasma leptin and adiponectin (Linco Research, St. Charles, MO). Plasma oxyntomodulin-like activity (OLI) was measured using an established in-house radioimmunoassay (5). Plasma taken 30 and 90 min after injection was analyzed by gel permeation chromatography to assess oxyntomodulin degradation (6). Insulin, peptide YY, and ghrelin-like immunoreactivity were measured using in-house radioimmunoassays (11–13). All samples were assayed in duplicate and within one assay to eliminate interassay variation. Glucose and total and HDL cholesterol were measured by an Olympus Analyzer.

Statistical analyses. Combined data are expressed as means ± SE. Data were compared using paired *t* tests for within-group analyses and unpaired *t* tests for between-group analyses. The changes in body weight, food data, and adipose hormones were normally distributed and therefore analyzed using parametric analyses. Serial measurements of pulse and blood pressure over time were compared using two-way ANOVA. The integrated area under the curve for incremental plasma insulin was calculated using the trapezoid rule. Prism Version 3.0 was used for statistical calculations, and *P* < 0.05 was considered significant.

RESULTS

Participant flow. Eligibility for the study was assessed in 89 subjects; 33 healthy subjects were eligible according to study criteria, and 56 were excluded because of either abnormal eating behavior or coexisting health problems. Of the 33 eligible subjects, 31 agreed to enter the study. After 2 subjects were excluded in the week before randomization because of noncompliance with saline injections, 29 were randomized into the study phase: 16 into the treatment group and 13 into the control group. One female subject, randomized to the treatment group, withdrew at 2 weeks. Another female subject, randomized to the control group, withdrew at 3 weeks. Both subjects cited a dislike of self-administering injections as their reason for withdrawal. A male volunteer, subject X, was removed from the treatment group after day 2 due to nausea. Intention-to-treat analysis was not possible as these subjects were lost to follow-up. Thus 26 subjects completed the entire 4-week injection period, 14 within the treatment group and 12 within the control group.

Effect of oxyntomodulin on body weight. Repeated preprandial subcutaneous injection of oxyntomodulin resulted in a 2.4 ± 0.4% reduction in body weight, compared with a 0.5 ± 0.6% reduction in the control group (*P* = 0.0129) over the 4-week study period. This represented a significant weight loss of 2.3 ± 0.4 kg in the treatment group and 0.5 ± 0.5 kg weight loss in the control group (*P* = 0.0106) (Fig. 1). After 2 weeks, interim data demonstrated a significant weight loss of 1.1 ± 0.3 kg in the treatment group (*P* = 0.0343), which increased to 1.4 ± 0.4 kg after 3 weeks (*P* = 0.0331). Thus, on average, there was an additional 0.45-kg weight loss per week in the oxyntomodulin group.

Over the study period, there was a change in adipose hormones in the treatment group consistent with a reduction in adipose tissue. At the start of the study, the plasma




FIG. 1. Change in body weight after self-administration of subcutaneous saline ($n = 12$) or oxyntomodulin (OXM; $n = 14$). In the treatment group, a significant weight loss of 1.1 ± 0.3 kg was observed after 2 weeks ($*P = 0.0343$). This loss increased to 1.4 ± 0.4 kg after 3 weeks of injections ($*P = 0.0331$). A total weight loss of 2.3 ± 0.4 kg was observed by the end of the 4-week study period ($**P = 0.0106$). SE bars are shown.

leptin and adiponectin levels in the control and treatment groups were not significantly different. In the treatment group, the plasma leptin levels were reduced from baseline (325 ± 50 ng/ml) by 79 ± 35 ng/ml ($19.5 \pm 6.2\%$); this change was significant when compared with the control group ($P = 0.0185$) (Table 2). The plasma adiponectin levels were increased from baseline (83.6 ± 9.5 μ g/ml) by 8.3 ± 4.3 μ g/ml ($9.4 \pm 4.9\%$) in the treatment group; this change was also significant when compared with the control group ($P = 0.0068$) (Table 2).

Effect of oxyntomodulin on food intake and appetite.

A subcutaneous injection of oxyntomodulin reduced energy intake during the study meal. The energy intake of the control group was not significantly different on day 1 (716 ± 108 kcal), 2 (660 ± 99 kcal), or 29 (711 ± 87 kcal). On day 1, when the treatment group self-administered a saline injection, energy intake (678 ± 45 kcal) was not statistically different from that of the control group. However, subjects who self-administered oxyntomodulin had significantly lower energy intake on days 2 (508 ± 46 kcal; $P = 0.0007$) and 29 (428 ± 61 kcal; $P = 0.0023$) when compared with their energy intake on day 1 (Fig. 2).

There were no significant differences in the preprandial visual analog scales for hunger and nausea in the control and treatment groups 30 min after injection (data not shown). There also was no difference in the postprandial

TABLE 2
Changes in screened analytes over the 4-week study period

	Control group		Treatment group		P
	Initial value	Change in value at 4 weeks	Initial value	Change in value at 4 weeks	
<i>n</i>	12	12	14	14	—
Leptin (ng/ml)	283 ± 24	31 ± 25	325 ± 50	-79 ± 35	0.0185*
Adiponectin (μ g/ml)	111.4 ± 11.3	-7.6 ± 3.0	83.6 ± 9.5	8.3 ± 4.3	0.0068*
Insulin (pmol/l)	26.7 ± 8.5	-1.1 ± 5.4	34.5 ± 4.7	-7.8 ± 4.4	0.3394
Glucose (mmol/l)	4.61 ± 0.09	0.21 ± 0.09	4.76 ± 0.11	0.12 ± 0.08	0.4111
Cholesterol (mmol/l)	4.71 ± 0.26	-0.26 ± 0.15	4.66 ± 0.19	-0.15 ± 0.11	0.5154
HDL (mmol/l)	1.07 ± 0.06	-0.01 ± 0.03	1.12 ± 0.09	-0.02 ± 0.03	0.8459
Ghrelin (pmol/l)	493 ± 87	47 ± 58	371 ± 65	35 ± 44	0.8586
Peptide YY (pmol/l)	11.8 ± 1.6	-2.6 ± 1.5	12.2 ± 2.1	-0.6 ± 1.6	0.3651

Data are means \pm SE. An unpaired *t* test was used to compare changes in the control vs. the treatment group. *Significant difference.




FIG. 2. Energy intake at the study meal, 30 min after self-administration of subcutaneous saline ($n = 10$) or oxyntomodulin (OXM; $n = 12$). The data are expressed as the percent of each subject's energy intake on day 1, when all subjects self-administered a saline injection. On day 2, the control group consumed $92.6 \pm 3.8\%$ of its energy intake on day 1, whereas the treatment group consumed only $75.3 \pm 5.3\%$ ($P = 0.0194$). On day 29, the control group consumed $107.0 \pm 7.7\%$ of its day 1 intake and the treatment group consumed $65.3 \pm 9.2\%$ ($P = 0.0030$). There was no significant difference between the amount consumed on day 2 and day 29 in the treatment group ($P = 0.0938$) or the amount consumed on day 2 and day 29 in the control group ($P = 0.1480$). SE bars are shown.

assessment of subjective palatability of the food between the control (day 2, 44 ± 6 mm; day 29, 48 ± 8 mm) and treatment (day 2, 43 ± 5 mm; day 29, 44 ± 7 mm) groups.

Only two subjects in each group failed to complete the study meal protocol. These subjects did not consume the same food type at each study meal. In the interest of accuracy, these subjects were excluded from food intake and appetite analysis before unblinding. Thus, study meal data were analyzed with $n = 10$ for the control group and $n = 12$ for the treatment group.

Plasma levels of oxyntomodulin. After an injection of oxyntomodulin, plasma OLI increased to a peak of 972 ± 165 pmol/l at 30 min (Fig. 3). In comparison, on the days that subjects administered saline, the mean baseline level of OLI was 97.4 ± 5.5 pmol/l and the peak postprandial level of OLI was 116.5 ± 10.4 pmol/l at 150 min.

Gel permeation analysis (14) of plasma samples from the treatment group demonstrated a single immunoreactive peak eluting at the same position as synthetic oxyntomodulin, with no indication of any degradative fragments. There was also no antibody formation at the end of the study, as assessed by plasma binding of the oxyntomodulin label.

Hormone, lipid, and glucose levels. There was a statis-




FIG. 3. Oxyntomodulin-like immunoreactivity levels for the subjects within the treatment group ($n = 14$) after self-administration of subcutaneous saline or oxyntomodulin (OXM). The time of injection was designated as $t = 0$ min, and the study meal was consumed at $t = 30$ min. SE bars are shown.

tically significant increase in preprandial insulin levels 30 min after subcutaneous administration of oxyntomodulin, but the increase was small compared with the effect of meal ingestion and not sufficient to affect plasma glucose levels (Table 3). This effect on insulin was present on both day 2 and day 29 (Fig. 4). The area under the curve for the incremental insulin level was nonsignificantly lower on day 2 ($18.0 \pm 0.9 \text{ nmol} \cdot \text{min}^{-1} \cdot \text{l}^{-1}$; $P = 0.1378$) and significantly lower on day 29 ($14.9 \pm 0.7 \text{ nmol} \cdot \text{min}^{-1} \cdot \text{l}^{-1}$; $P = 0.0398$) over the 4-h postinjection period compared with the day saline was administered (day 1: $21.6 \pm 1.1 \text{ nmol} \cdot \text{min}^{-1} \cdot \text{l}^{-1}$).

The injection of oxyntomodulin did not acutely affect plasma levels of peptide YY or ghrelin 30 min after injection (Table 3). There were no other significant chronic preinjection changes observed over the 4-week study period in the analytes screened (e.g., in lipid parameters or insulin levels) (Table 2).

Adverse events. Short-lived minor discomfort at the injection site was reported by 8 of 14 of the subjects in the treatment group and 5 of 12 of the control group. In total, 6.9% of all oxyntomodulin injections and 6.6% of all saline injections were reported to cause minor discomfort; thus discomfort did not appear related to the substance injected.

Transient mild nausea was reported with 3% of oxyntomodulin injections compared with 0.2% of all saline injections ($P = 0.0389$). These data included subject Y, who experienced nausea associated with 13 of the first 47 injections. The dose subject Y administered was halved, and the incidence of nausea decreased to 2 out of the remaining 37 injections. On day 29, subject Y had a peak OLI level of 868 pmol/l, compared with the mean peak for the group of 972 ± 619 pmol/l, despite taking only half the




FIG. 4. Insulin levels for the treatment group ($n = 14$) after self-administration of saline (day 1) and oxyntomodulin (days 2 and 29). The time of injection was designated as $t = 0$ min, and the study meal was consumed at $t = 30$ min. After oxyntomodulin injection, there was a small but significant rise in preprandial insulin (***) day 2, $P = 0.0004$; day 29, $P = 0.0008$). On day 1, the integrated area under the curve for the incremental insulin level after the injection was $21.6 \pm 1.1 \text{ nmol} \cdot \text{min}^{-1} \cdot \text{l}^{-1}$. The area under the curve was nonsignificantly lower on day 2 ($18.0 \pm 0.9 \text{ nmol} \cdot \text{min}^{-1} \cdot \text{l}^{-1}$; $P = 0.1378$) and significantly lower on day 29 ($14.9 \pm 0.7 \text{ nmol} \cdot \text{min}^{-1} \cdot \text{l}^{-1}$; $P = 0.0398$). SE bars are shown.

dose of oxyntomodulin. A second participant, subject X, was removed from the treatment group after day 2 because of adverse effects and was not included in the analysis; after each of the initial three oxyntomodulin injections, subject X had significant nausea that interfered with daily activities. This subject's plasma OLI levels peaked at 2,140 pmol/l 15 min postinjection and were significantly higher than the rest of the treatment group over the entire 4-h study period ($P < 0.0001$ by two-way ANOVA). Another subject regularly experienced nausea for 2 days during her midmenstrual cycle and was included in the treatment group data in the interest of stringency.

The incidence of all other reported adverse events was $<2\%$ and not significantly different between the treatment and control groups. There were no effects of oxyntomodulin self-administration on pulse rate or systolic or diastolic blood pressure as measured during the study meal days.

Follow-up data. The treatment group regained 1.1 ± 0.4 kg 2 weeks after terminating the oxyntomodulin injections. The control group gained 0.5 ± 0.2 kg and were 0.0 ± 0.6 kg different from their initial weight. The difference in weight loss between the control and treatment group was no longer significant 2 weeks after the end of the study period ($P = 0.134$). One subject in the control group and one in the treatment group were lost to follow-up after the end of the study; therefore, the follow-up data were analyzed with $n = 11$ in the control group and $n = 13$ in the treatment group.

TABLE 3

Acute response of analytes to injection of saline or oxyntomodulin in the treatment group

	Day 1 (saline)		Day 2 (oxyntomodulin)			Day 29 (oxyntomodulin)		
	Initial value	Change at 30 min	Initial value	Change at 30 min	P	Initial value	Change at 30 min	P
Insulin (pmol/l)	34.5 ± 4.7	-6.1 ± 3.2	27.4 ± 3.6	32.6 ± 7.7	0.0004*	26.7 ± 4.3	40.5 ± 10.1	0.0008*
Glucose (mmol/l)	4.79 ± 0.09	-0.04 ± 0.06	4.73 ± 0.10	0.07 ± 0.08	0.300	4.88 ± 0.09	0.02 ± 0.07	0.6629
Ghrelin (pmol/l)	370 ± 65	-5 ± 23	375 ± 74	-42 ± 35	0.3429	405 ± 86	-49 ± 22	0.2327
Peptide YY (pmol/l)	12.2 ± 2.1	1.1 ± 2.2	15.2 ± 4.4	-3.5 ± 4.1	0.3377	12.1 ± 1.6	-2.6 ± 1.3	0.1637

Data are means \pm SE. A paired t test was used to compare the change from baseline 30 min after an injection of oxyntomodulin with the change on day 1 when the treatment group injected saline. *Significant difference.

DISCUSSION

Self-administering oxyntomodulin subcutaneously effectively reduced body weight by 2.4% over a 4-week period in healthy overweight and obese volunteers. This effect was substantial considering that all subjects were asked to continue their usual diet and level of exercise. The accompanying reduction in leptin and increase in adiponectin suggest that the weight loss demonstrated was at least partly due to a reduction in adipose tissue. A significant reduction in energy intake with preservation of subjective food palatability was also observed. The magnitude of this anorectic effect was sufficient to account for the weight loss in the treatment group (15).

The plasma OLI levels achieved by subcutaneous injection were similar to those achieved by an intravenous infusion of $3.0 \text{ pmol} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ in a study that reduced food intake by 19.3% in normal weight subjects (6). This anorectic effect was comparable with the effect observed in overweight subjects in the current study. This suggests that oxyntomodulin has an equipotent anorectic effect in normal weight and overweight subjects and that obesity does not confer resistance to the anorectic effect of oxyntomodulin.

A recent human intravenous infusion study demonstrated no effect of oxyntomodulin on insulin or glucose levels in normal-weight humans (6). In the current study, there was no evidence of a postprandial incretin effect. There was a small increase in preprandial plasma insulin after oxyntomodulin injection. However, this increase was not sufficient to alter plasma glucose levels. Other gut hormones, such as glucagon-like peptide, are known to have a powerful incretin effect (12). Indeed, glucagon-like peptide 1, while improving glucose tolerance when given to type 2 diabetic subjects (16), can cause hypoglycemia in nondiabetic subjects (17). There was no evidence of a change in baseline insulin, glucose, or lipid levels over the 4-week study period in this group of healthy obese subjects. Further studies are needed to assess to what degree long-term oxyntomodulin administration leads to an improvement in metabolic parameters in obese subjects with abnormal glucose and lipid metabolism.

The current study did not demonstrate an effect of oxyntomodulin on the circulating level of the orexigenic hormone ghrelin. It has been suggested that a proportion of the anorectic effect of intravenously administered oxyntomodulin is secondary to its suppressant effect on ghrelin (6). However, this does not seem a likely mechanism for the dose administered subcutaneously in the current study. The decrease in body weight was secondary to a significant reduction in energy intake. This reduction in energy intake was likely due to a reduction in appetite, as there was preservation of food satisfaction and therefore no evidence of taste aversion. Indeed, a previous study demonstrated that administration of intravenous oxyntomodulin reduces preprandial hunger scores in normal-weight humans (6). However, the current study did not demonstrate a change in subjective appetite scores before the meal after a subcutaneous injection of oxyntomodulin. This may have reflected a perception of early satiety rather than a reduced preprandial appetite or a failure of the visual analog scores as a tool of appetite measurement.

The self-administration of subcutaneous oxyntomodulin

was well-tolerated by the subjects, and good compliance rates were achieved. This was a fixed initial dose study; a dose of 400 nmol oxyntomodulin may not be appropriate for all subjects. Indeed, subject Y experienced a much lower incidence of nausea after the dose was reduced. Subject X demonstrated extremely high plasma OLI levels after oxyntomodulin injection and experienced significant nausea. These subjects may have absorbed oxyntomodulin more rapidly or metabolized the peptide more slowly. Larger groups studies are needed to assess if this was an idiosyncratic response.

These preliminary data suggest that the administration of oxyntomodulin could be an effective treatment for obesity. The reduction in food intake seen at the end of the study was statistically no different to the reduction on day 2, indicating that the efficacy of oxyntomodulin was maintained. In addition, the rate of weight loss was consistent over the 4-week study period. Oxyntomodulin levels are increased in patients who have had jejunoileal bypass surgery (18,19). This sustained elevation of plasma oxyntomodulin may be one of the factors leading to successful weight reduction in these patients. There was no evidence of oxyntomodulin antibody formation from the assay data. Together, this suggests that administration of oxyntomodulin beyond the 4-week study period may be expected to result in continued weight loss. However, these data are limited by our small sample size and the 4-week duration of the study. Longer-term clinical trials involving larger numbers of participants are required to demonstrate the long-term efficacy of oxyntomodulin as an antiobesity therapy.

In summary, subcutaneous self-administration of oxyntomodulin three times daily in the community reduced body weight, decreased food intake, and altered the levels of adipose hormones in overweight and obese nondiabetic human subjects. The anorectic effect was well maintained over the 4-week study period, and the weight loss compared favorably with that achieved through other drug therapies. Currently available pharmacological agents licensed for weight reduction therapy have limited efficacy (20). Several drugs now in development affect widely distributed central neurotransmitter systems and may therefore have a broad spectrum of side effects. Mimicking postprandial satiety by administering a natural postprandial hormone such as oxyntomodulin may provide a more specific treatment for obesity.

ACKNOWLEDGMENTS

K.W. and A.J.P. are Wellcome Trust Clinical Training Fellows. This study was funded with project grants from the Wellcome Trust and Medical Research Council. The funding sponsors had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; or preparation, review, and approval of the manuscript.

The study was conceived by K.W., A.J.P., and S.R.B. The study protocol and ethical application was formulated by K.W., A.J.P., C.J.S., M.P., S.M.E., K.G.M., A.M.W., G.S.F., K.M., M.A.G., and S.R.B. K.W. and A.J.P. enrolled patients and performed the study interventions, hormone assays, and data analysis. Dietetic support and data analysis were performed by S.M.E. in collaboration with G.S.F. K.W. and

A.J.P. drafted the manuscript, and all authors were involved in its critical appraisal and final approval.

The use of oxyntomodulin for the treatment of obesity is the subject of two pending patent applications (WO 2003/022304 and WO 2004/06285) in the name of Imperial College Innovations, which have been exclusively licensed to Thiakis Limited.

We thank the volunteers for their invaluable help with the study. We also thank Dr. David Stephens for statistical advice and the Department of Biochemistry, Charing Cross Hospital, for its assistance with the measurement of glucose and lipids.

REFERENCES

- Hedley AA, Ogden CL, Johnson CL, Carroll MD, Curtin LR, Flegal KM: Prevalence of overweight and obesity among US children, adolescents, and adults, 1999–2002. *JAMA* 291:2847–2850, 2004
- Batterham RL, Cowley MA, Small CJ, Herzog H, Cohen MA, Dakin CL, Wren AM, Brynes AE, Low MJ, Ghatei MA, Cone RD, Bloom SR: Gut hormone PYY(3–36) physiologically inhibits food intake. *Nature* 418:650–654, 2002
- Wren AM, Seal LJ, Cohen MA, Brynes AE, Frost GS, Murphy KG, Dhillo WS, Ghatei MA, Bloom SR: Ghrelin enhances appetite and increases food intake in humans. *J Clin Endocrinol Metab* 86:5992, 2001
- Flint A, Raben A, Astrup A, Holst JJ: Glucagon-like peptide 1 promotes satiety and suppresses energy intake in humans. *J Clin Invest* 101:515–520, 1998
- Ghatei MA, Uttenthal LO, Christofides ND, Bryant MG, Bloom SR: Molecular forms of human enteroglucagon in tissue and plasma: plasma responses to nutrient stimuli in health and in disorders of the upper gastrointestinal tract. *J Clin Endocrinol Metab* 57:488–495, 1983
- Cohen MA, Ellis SM, Le Roux CW, Batterham RL, Park A, Patterson M, Frost GS, Ghatei MA, Bloom SR: Oxyntomodulin suppresses appetite and reduces food intake in humans. *J Clin Endocrinol Metab* 88:4696–4701, 2003
- Dakin CL, Small CJ, Batterham RL, Neary NM, Cohen MA, Patterson M, Ghatei MA, Bloom SR: Peripheral oxyntomodulin reduces food intake and body weight gain in rats. *Endocrinology* 145:2687–2695, 2004
- Van Strien T, Rookus MA, Bergers GP, Frijters JE, Defares PB: Life events, emotional eating and change in body mass index. *Int J Obes* 10:29–35, 1986
- Morgan JF, Reid F, Lacey JH: The SCOFF questionnaire: assessment of a new screening tool for eating disorders. *BMJ* 319:1467–1468, 1999
- Batterham RL, Cohen MA, Ellis SM, Le Roux CW, Withers DJ, Frost GS, Ghatei MA, Bloom SR: Inhibition of food intake in obese subjects by peptide YY3–36. *N Engl J Med* 349:941–948, 2003
- English PJ, Ghatei MA, Malik IA, Bloom SR, Wilding JP: Food fails to suppress ghrelin levels in obese humans. *J Clin Endocrinol Metab* 87:2984, 2002
- Kreymann B, Williams G, Ghatei MA, Bloom SR: Glucagon-like peptide-1 7–36: a physiological incretin in man. *Lancet* 2:1300–1304, 1987
- Adrian TE, Savage AP, Sagor GR, Allen JM, Bacarese-Hamilton AJ, Tatemoto K, Polak JM, Bloom SR: Effect of peptide YY on gastric, pancreatic, and biliary function in humans. *Gastroenterology* 89:494–499, 1985
- Murphy KG, Abbott CR, Mahmoudi M, Hunter R, Gardiner JV, Rossi M, Stanley SA, Ghatei MA, Kuhar MJ, Bloom SR: Quantification and synthesis of cocaine- and amphetamine-regulated transcript peptide (79–102)-like immunoreactivity and mRNA in rat tissues. *J Endocrinol* 166:659–668, 2000
- Lean ME, James WP: Prescription of diabetic diets in the 1980s. *Lancet* 1:723–725, 1986
- Zander M, Madsbad S, Madsen JL, Holst JJ: Effect of 6-week course of glucagon-like peptide 1 on glycaemic control, insulin sensitivity, and beta-cell function in type 2 diabetes: a parallel-group study. *Lancet* 359:824–830, 2002
- Edwards CM, Todd JF, Ghatei MA, Bloom SR: Subcutaneous glucagon-like peptide-1 (7–36) amide is insulinotropic and can cause hypoglycaemia in fasted healthy subjects. *Clin Sci (Lond)* 95:719–724, 1998
- Sarson DL, Scopinaro N, Bloom SR: Gut hormone changes after jejunoileal (JIB) or biliopancreatic (BPP) bypass surgery for morbid obesity. *Int J Obes* 5:471–480, 1981
- Holst JJ, Sorensen TI, Andersen AN, Stadil F, Andersen B, Lauritsen KB, Klein HC: Plasma enteroglucagon after jejunoileal bypass with 3:1 or 1:3 jejunoileal ratio. *Scand J Gastroenterol* 14:205–207, 1979
- Finer N: Pharmacotherapy of obesity. *Best Pract Res Clin Endocrinol Metab* 16:717–742, 2002