

# HLA Genotyping Supports a Nonautoimmune Etiology in Patients Diagnosed With Diabetes Under the Age of 6 Months

Emma L. Edghill,<sup>1</sup> Rachel J. Dix,<sup>2</sup> Sarah E. Flanagan,<sup>1</sup> Polly J. Bingley,<sup>2</sup> Andrew T. Hattersley,<sup>1</sup> Sian Ellard,<sup>1</sup> and Kathleen M. Gillespie<sup>2</sup>

Children with permanent diabetes are usually assumed to have type 1 diabetes. It has recently been shown that there are genetic subgroups of diabetes that are often diagnosed during the neonatal period but may present later. A recent Italian study proposed that type 1 diabetes is rare before 6 months of age. We aimed to examine genetic susceptibility to type 1 diabetes in patients diagnosed with diabetes before the age of 2 years. We analyzed HLA class II genotypes, markers of autoimmune diabetes, in 187 children with permanent diabetes diagnosed at <2 years of age. Of the 79 subjects diagnosed at <6 months of age, 41% (95% CI 0.30–0.51) had type 1 diabetes–associated high-risk genotypes, a proportion similar to that in healthy population control subjects (44%,  $P = 0.56$ ). This group included 32 patients with mutations in the *KCNJ11* gene, which encodes Kir6.2 (44% high-risk HLA class II genotypes), and 47 in whom the etiology of diabetes was unknown (38% high-risk HLA class II genotypes). Of 108 patients diagnosed between 6 and 24 months of age, 93% (0.86–0.99) had high-risk HLA class II genotypes compared with 44% of the population control subjects ( $P < 0.0001$ ). We conclude that infants diagnosed with diabetes before 6 months of age are unlikely to have autoimmune type 1 diabetes and are most likely to have a monogenic etiology. *Diabetes* 55:1895–1898, 2006

Appropriate treatment regimens for diabetes require accurate characterization of disease (1). For patients diagnosed in infancy, this may be complicated; several different etiologies may cause diabetes, including type 1 diabetes and permanent neonatal diabetes (PNDM). Type 1 diabetes results from autoimmune destruction of insulin-producing  $\beta$ -cells and is characterized by the presence of multiple islet autoantibodies and high-risk HLA haplotypes for type 1 diabetes. HLA DRB1\*04-DQB1\*0302 and/or HLA DRB1\*03-

DQB1\*0201 are observed in >90% of affected children and in only 40% of the general population (2). The study of HLA genotypes may therefore help define which patients diagnosed in infancy are likely to have type 1 diabetes.

PNDM is typically defined as diabetes requiring permanent insulin treatment that is diagnosed in the first 3 months of life (3). Recent advances in determining the etiology of PNDM have demonstrated that heterozygous-activating mutations in the *KCNJ11* gene encoding the ATP-sensitive  $K^+$  channel subunit Kir6.2 account for ~40–64% of PNDM cases (4). To date, all activating *KCNJ11* mutations have been identified in subjects diagnosed under the age of 6 months (5–11). Some of these patients are diagnosed outside the defined neonatal period of 3 months and therefore highlight a need to investigate the etiologies of diabetes in infancy. The identification of patients with *KCNJ11* mutations is important, as most respond well to sulfonylureas and achieve better glycemic control following transfer from insulin to sulfonylurea tablets (6,8,12–14).

A recent Italian study suggested that children diagnosed with diabetes before 6 months of age do not have the genetic characteristics of autoimmune diabetes (15). Many patients from this cohort have since been diagnosed with *KCNJ11* mutations (10), but the etiology of diabetes in the remainder is uncertain. To determine the likelihood of type 1 diabetes in children diagnosed in infancy, we investigated the frequency of high-risk HLA class II genotypes in a cohort of 187 subjects diagnosed with permanent diabetes under the age of 2 years, 32 with *KCNJ11* mutations and 155 in whom mutations have been excluded.

## RESEARCH DESIGN AND METHODS

A total of 187 subjects were recruited worldwide. The inclusion criteria for this study were a diagnosis of diabetes before 24 months of age requiring continual insulin treatment from diagnosis. The physician's classification was type 1 diabetes in the majority of cases. This cohort included 104 U.K. samples, of whom 58 subjects were recruited from the 1972–1981 British Diabetic Association Under 2s cohort (16) and 19 from the Bart's Oxford study of childhood diabetes (17). Of the 187 samples, 79 were diagnosed at 0–6 months of age (22 of U.K. origin), 45 were diagnosed at 7–12 months (21 of U.K. origin), 32 were diagnosed at 13–18 months (30 of U.K. origin), and 31 were diagnosed at 19–24 months (31 of U.K. origin). Consent was obtained from all patients or their parents.

HLA genotypes on 621 adult U.K. Caucasian control subjects from the general population with no history of autoimmune disease have been described previously (18). All control subjects gave written informed consent. **Sequencing of *KCNJ11*.** All subjects were sequenced for *KCNJ11* as previously described (5–7,11 and personal communication with A.L. Gloyn, Diabetes Research Laboratories, University of Oxford, Oxford, U.K.).

From the <sup>1</sup>Institute of Biomedical and Clinical Science, Peninsula Medical School, Exeter, U.K.; and the <sup>2</sup>Department of Clinical Science at North Bristol, University of Bristol, Bristol, U.K.

Address correspondence and reprint requests to Dr. K.M. Gillespie, Medical School Unit, Southmead Hospital, Bristol BS10 5NB, U.K. E-mail: k.m.gillespie@bristol.ac.uk.

Received for publication 20 January 2006 and accepted in revised form 13 March 2006.

Additional information for this article can be found in an online appendix at <http://diabetes.diabetesjournals.org>.

PNDM, permanent neonatal diabetes.

DOI: 10.2337/db06-0094

© 2006 by the American Diabetes Association.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

TABLE 1

The frequency of HLA class II genotypes in 32 individuals with a defined mutation in *KCNJ11* and in 155 negative for mutations in *KCNJ11* 46 diagnosed before and 109 after the age of 6 months

	<i>KCNJ11</i> Mut +ve	<i>KCNJ11</i> Mut -ve (0–6 months)	Kir6.2 Mut -ve (>6 months)	Healthy control subjects
<i>n</i>	32	47	108	621
Genotype				
DR4-DQ8/DR3-DQ2	1 (3%)	4 (9%)	54 (50%)	17 (3%)
DR4-DQ8/X	6 (19%)	7 (15%)	18 (16%)	85 (14%)
DR3-DQ2/X	7 (22%)	7 (15%)	30 (28%)	167 (26%)
X/X	13 (40%)	21 (44%)	2 (2%)	249 (40%)
DR2-DQ6/ DR2-DQ6 or DR2-DQ6/X	5 (16%)	8 (17%)	4 (4%)	103 (17%)

Control data are provided for 621 healthy Caucasians. The genotypes HLA DRB1\*04-DQB1\*0302/ HLA DRB1\*03-DQB1\*0201, HLA DRB1\*04-DQB1\*0302/X, and HLA DRB1\*03-DQB1\*0201/X are described as DR4-DQ8/DR3-DQ2, DR4-DQ8/X, and DR3-DQ2/X, respectively, where X is not DRB1\*02-DQB1\*0602 (DR2-DQ6). +ve, mutation present; -ve, mutation not detected by sequencing.

**HLA class II genotyping.** HLA-DRB1, -DQA1, and -DQB1 genotyping was performed using a Dynal reverse SSOP method (Dynal Biotech, Warral, U.K.). Details of HLA class II genotypes in the 77 individuals from the Bart's Oxford study/British Diabetic Association Under 2s study and 621 U.K. control subjects were already available (7,18). In the absence of parental samples, haplotypes were identified on the basis of well-established patterns of linkage disequilibrium between HLA DRB1 and DQB1.

For the purposes of this study, the genotypes HLA DRB1\*04-DQB1\*0302/X or HLA DRB1\*03-DQB1\*0201/X (where X was not DRB1\*02-DQB1\*0602) were defined as high risk. This includes the highest risk genotype HLA DRB1\*04-DQB1\*0302/HLA DRB1\*03-DQB1\*0201. All other genotypes including those containing the protective DRB1\*02-DQB1\*0602 were defined as low risk.

**Statistical analysis.** To examine differences in the frequencies of HLA genotypes,  $\chi^2$  analysis was used. All data were stored according to existing data protection regulations.

## RESULTS

As previously reported, 32 of the 187 (17%) patients had heterozygous mutations in the *KCNJ11* gene. All subjects with a *KCNJ11* mutation were diagnosed with diabetes before 6 months of age (median 1.15 months [range 0.23–6]).

The overall results obtained are shown in Table 1. Of 32 individuals with a Kir6.2 mutation, 1 (3%) had the highest-risk HLA DRB1\*04-DQB1\*0302/HLA DRB1\*03-DQB1\*0201 genotype, whereas 16% had protective genotypes. These data are remarkably similar to those of the healthy control population, with frequencies of 3 and 17%, respectively. Of 47 children diagnosed under the age of 6 months with no mutation in the *KCNJ11* gene, 4 (9%) carried the highest-risk genotype for type 1 diabetes and 17% were positive for protective genotypes, also very similar to the healthy control population. This is in marked contrast to 108 children diagnosed at >6 months of age, where 50% had the highest-risk HLA genotype and only 4% had protective genotypes. These data, including mutation status, country of origin, HLA data, and age at diagnosis, are available in an online appendix (available at <http://diabetes.diabetesjournals.org>).

Overall, individuals diagnosed under 6 months of age, irrespective of *KCNJ11* mutation status, had very similar frequencies of type 1 diabetes-associated genotypes (41% [95% CI 0.3–0.51]) compared with the healthy control population (44%,  $P = 0.56$ ). In contrast, 93% (101 of 108) (95% CI 0.86–0.96) of children diagnosed with diabetes over the age of 6 months, who screened negative for a *KCNJ11* mutation, had at least one high-risk haplotype for type 1 diabetes with a similar distribution in the three 6-month intervals: 6–12 months (89%), 12–18 months (94%), and 18–24 months (97%), all different from those <6 months ( $P < 0.00001$  for all) (Fig. 1).

Since the role of HLA class II genotypes in susceptibility to type 1 diabetes is less well characterized in non-European populations, we analyzed European Caucasians with permanent diabetes separately and showed that the results were very similar to those observed in the international cohort. The European Caucasian subjects diagnosed with diabetes at <6 months of age ( $n = 50$ ) had the same frequency of high-risk HLA class II genotypes as normal population (44%,  $P = 0.99$ ), whereas those diagnosed at >6 months ( $n = 92$ ) had a higher prevalence (93%) of high susceptibility HLA compared with those diagnosed at <6 months ( $P < 0.00001$ ).

## DISCUSSION

We have shown that HLA class II high-risk genotypes for susceptibility to type 1 diabetes are much more common in subjects diagnosed with diabetes over the age of 6 months than in those diagnosed before 6 months. The frequency of high-risk HLA class II genotypes in patients diagnosed before 6 months, regardless of *KCNJ11* mutation status, was similar to that found in control subjects. Individuals diagnosed before 6 months of age are therefore unlikely to have type 1 diabetes, and even those who do not have mutations in the *KCNJ11* gene are likely to have other forms of monogenic diabetes.

Our data support a previous elegant study (15) of HLA class II genotypes in 111 Italian subjects diagnosed under 12 months of age, which suggested that individuals diagnosed with diabetes under the age of 6 months frequently have protective HLA genotypes for type 1 diabetes. They also showed that patients diagnosed before 6 months of age were less likely to have  $\beta$ -cell autoantibodies ( $n = 46$ ) and more likely to have low birth weight (small for gestational age) than patients diagnosed after 6 months. Low birth weight is an indicator of reduced insulin secretion in utero and is consistent with a genetic cause of reduced insulin secretion in utero. This has been described with known monogenic causes of neonatal diabetes (19) such as heterozygous-activating *KCNJ11* mutations (6) and homozygous glucokinase mutations (20). In contrast, there is evidence that HLA genotypes associated with type 1 diabetes are associated with high birth weight (21).

Although this is the largest study of HLA genotypes in patients aged between 0 and 2 years, it is difficult to be certain of an absolute cutoff at 6 months. The fact that the prevalence of high-risk HLA is slightly lower in those




FIG. 1. The prevalence of high-risk HLA class II genotypes (DR4-DQ8/DR3-DQ2, DR4-DQ8/X, and DR3-DQ2/X, where X is not DR2-DQ6) in all subjects from the international cohort ( $n = 187$ ) diagnosed with diabetes under 24 months of age and in the normal control population. The group diagnosed under 6 months was compared with the three age-groups over 6 months. \*\*\*\* $P < 0.00001$ .

diagnosed between 6 and 12 months of age compared with those between 12 and 24 months of age indicates that there may be a few patients with nonautoimmune diabetes in the 6- to 12-month age range.

Our data emphasize that, as expected, there is not a role for HLA class II genotyping in the classification of type 1 diabetes on an individual basis. While the absence of a high-risk HLA genotype makes type 1 diabetes unlikely, the presence of high-risk HLA does not exclude nonautoimmune diabetes. Approximately 40% of patients with *KCNJ11* mutations have high-risk HLA genotypes, similar to their prevalence in the general population.

The risk associated with type 1 diabetes HLA haplotypes differs between continents (22). In this study, we assumed that HLA-DRB1\*04-DQB1\*0302 and HLA-DRB1\*03-DQB1\*0201 are high risk in all populations. We addressed this potential source of error by analyzing subjects of European Caucasian origin separately and obtained results similar to those of the international cohort.

We have shown that the frequency of HLA class II high-risk genotypes is significantly different in subjects with insulin-treated permanent diabetes diagnosed over and under 6 months of age. Those diagnosed under 6 months of age have a distribution of high-risk genotypes similar to that of the normal population. Our study confirms that children diagnosed with diabetes under 6 months of age are unlikely to have type 1 diabetes and should be screened for *KCNJ11* gene mutations.

#### ACKNOWLEDGMENTS

This study was funded by Diabetes UK and the Wellcome Trust at the University of Bristol and at Peninsula Medical School. A.T.H. is a Wellcome Trust Research Fellow.

We thank Prof. Steve Gough for providing HLA data on healthy control subjects.

#### REFERENCES

- Pearson ER, Starkey BJ, Powell RJ, Gribble FM, Clark PM, Hattersley AT: Genetic aetiology of hyperglycaemia determines response to treatment in diabetes. *Lancet* 362:1275–1281, 2003
- Devendra D, Eisenbarth GS: 17. Immunologic endocrine disorders. *J Allergy Clin Immunol* 111 (Suppl. 2):S624–S636, 2003
- Shield JP: Neonatal diabetes: new insights into aetiology and implications. *Horm Res* 53 (Suppl. 1):7–11, 2000
- Hattersley AT, Ashcroft FM: Activating mutations in Kir6.2 and neonatal diabetes: new clinical syndromes, new scientific insights, and new therapy. *Diabetes* 54:2503–2513, 2005
- Gloyn AL, Cummings EA, Edghill EL, Harries LW, Scott R, Costa T, Temple IK, Hattersley AT, Ellard S: Permanent neonatal diabetes due to paternal germline mosaicism for an activating mutation of the *KCNJ11* gene encoding the Kir6.2 subunit of the beta-cell potassium adenosine triphosphate channel. *J Clin Endocrinol Metab* 89:3932–3935, 2004
- Gloyn AL, Pearson ER, Antcliff JF, Proks P, Bruining GJ, Slingerland AS, Howard N, Srinivasan S, Silva JM, Molnes J, Edghill EL, Frayling TM, Temple IK, Mackay D, Shield JP, Sunnik Z, van Rhijn A, Wales JK, Clark P, Gorman S, Aisenberg J, Ellard S, Njolstad PR, Ashcroft FM, Hattersley AT: Activating mutations in the gene encoding the ATP-sensitive potassium-channel subunit Kir6.2 and permanent neonatal diabetes. *N Engl J Med* 350:1838–1849, 2004
- Edghill EL, Gloyn AL, Gillespie KM, Lambert AP, Raymond NT, Swift PG, Ellard S, Gale EA, Hattersley AT: Activating mutations in the *KCNJ11* gene

- encoding the ATP-sensitive K<sup>+</sup> channel subunit Kir6.2 are rare in clinically defined type 1 diabetes diagnosed before 2 years. *Diabetes* 53:2998–3001, 2004
8. Sagen JV, Raeder H, Hathout E, Shehadeh N, Gudmundsson K, Baevre H, Abuelo D, Phornphutkul C, Molnes J, Bell GI, Gloyn AL, Hattersley AT, Molven A, Sovik O, Njolstad PR: Permanent neonatal diabetes due to mutations in KCNJ11 encoding Kir6.2: patient characteristics and initial response to sulfonylurea therapy. *Diabetes* 53:2713–2718, 2004
  9. Vaxillaire M, Populaire C, Busiah K, Cave H, Gloyn AL, Hattersley AT, Czernichow P, Froguel P, Polak M: Kir6.2 mutations are a common cause of permanent neonatal diabetes in a large cohort of French patients. *Diabetes* 53:2719–2722, 2004
  10. Massa O, Iafusco D, D'Amato E, Gloyn AL, Hattersley AT, Pasquino B, Tonini G, Dammacco F, Zanette G, Meschi F, Porzio O, Bottazzo G, Crino A, Lorini R, Cerutti F, Vanelli M, Barbetti F: KCNJ11 activating mutations in Italian patients with permanent neonatal diabetes. *Hum Mutat* 25:22–27, 2005
  11. Flanagan SE, Edghill E, Gloyn AL, Stals K, Ellard S, Hattersley AT: Kir6.2 gene mutations are a common cause of diabetes before 6 months and the presence of neurological features varies with the site of the mutation. *Diabet Med* 22: 10, 2005
  12. Klupa T, Edghill EL, Nazim J, Sieradzki J, Ellard S, Hattersley AT, Malecki MT: The identification of a R201H mutation in KCNJ11, which encodes Kir6.2, and successful transfer to sustained-release sulphonylurea therapy in a subject with neonatal diabetes: evidence for heterogeneity of beta cell function among carriers of the R201H mutation. *Diabetologia* 48:1029–1031, 2005
  13. Zung A, Glaser B, Nimri R, Zadik Z: Glibenclamide treatment in permanent neonatal diabetes mellitus due to an activating mutation in Kir6.2. *J Clin Endocrinol Metab* 89:5504–5507, 2004
  14. Codner E, Flanagan S, Ellard S, Garcia H, Hattersley AT: High-dose glibenclamide can replace insulin therapy despite transitory diarrhea in early-onset diabetes caused by a novel R201L Kir6.2 mutation. *Diabetes Care* 28:758–759, 2005
  15. Iafusco D, Stazi MA, Cotichini R, Cotellessa M, Martinucci ME, Mazzella M, Cherubini V, Barbetti F, Martinetti M, Cerutti F, Prisco F: Permanent diabetes mellitus in the first year of life. *Diabetologia* 45:798–804, 2002
  16. Botha JL, Parker H, Raymond NT, Swift PG: Diabetes diagnosed before the age of 2 years: mortality in a British cohort 8–17 years after onset. *Int J Epidemiol* 21:1132–1137, 1992
  17. Bingley PJ, Bonifacio E, Williams AJ, Genovese S, Bottazzo GF, Gale EA: Prediction of IDDM in the general population: strategies based on combinations of autoantibody markers. *Diabetes* 46:1701–1710, 1997
  18. Simmonds MJ, Howson JM, Heward JM, Cordell HJ, Foxall H, Carr-Smith J, Gibson SM, Walker N, Tomer Y, Franklyn JA, Todd JA, Gough SC: Regression mapping of association between the human leukocyte antigen region and Graves disease. *Am J Hum Genet* 76:157–163, 2005
  19. Hattersley AT, Tooke JE: The fetal insulin hypothesis: an alternative explanation of the association of low birth weight with diabetes and vascular disease. *Lancet* 353:1789–1792 1999
  20. Njolstad P, Sovik O, Cuesta-Munoz A, Bjorkhaug L, Massa O, Barbetti F, Undlien D, Shiota C, Magnuson M, Molven A, Matschinsky F, Bell G: Neonatal diabetes mellitus due to complete glucokinase deficiency. *N Engl J Med* 344:1588–1592, 2001
  21. Larsson HE, Lynch K, Lernmark B, Nilsson A, Hansson G, Almgren P, Lernmark A, Ivarsson SA: Diabetes-associated HLA genotypes affect birthweight in the general population. *Diabetologia* 48:1484–1491, 2005
  22. Ronningen KS, Keiding N, Green A: Correlations between the incidence of childhood-onset type I diabetes in Europe and HLA genotypes. *Diabetologia* 44 (Suppl. 3):B51–B59, 2001