

Genetic Influences of Adiponectin on Insulin Resistance, Type 2 Diabetes, and Cardiovascular Disease

Claudia Menzaghi,¹ Vincenzo Trischitta,^{1,2} and Alessandro Doria³

Recent evidence points to molecules secreted by the adipose tissue, or adipokines, as possible links between increased adipose mass and metabolic abnormalities. Among these molecules, adiponectin has drawn much attention because of its insulin-sensitizing and antiatherogenic actions, suggesting that genetic deficits in its production or action may contribute to insulin resistance and coronary artery disease (CAD). A meta-analysis of the data published to date supports this hypothesis. Two independent effects, corresponding to the two linkage disequilibrium blocks that can be identified at the adiponectin locus, appear to be present. In the 5' block, the g.-11391G→A variant has a modest but significant effect on adiponectinemia, with a mean difference between genotypes of 1.64 ng/ml (95% CI 0.88–2.41). In the 3' block, the g.+276G→T variant is a strong determinant of insulin resistance and CAD, with minor allele homozygotes having a lower homeostasis model assessment of insulin resistance (HOMA_{IR}) index (–0.36 units, 95% CI 0.24–0.47) and a lower cardiovascular risk (odds ratio 0.55, 95% CI 0.38–0.80) than carriers of other genotypes. No consistent effect on BMI or risk of type 2 diabetes is evident. Polymorphisms in the genes coding for the adiponectin receptors may also influence the risk of insulin resistance and CAD, but data on these genes are still too sparse to draw firm conclusions. In summary, the studies published to date indicate that polymorphisms at the adiponectin locus are indeed predictors of circulating adiponectin levels, insulin sensitivity, and atherosclerosis, highlighting the pivotal role of this adipokine in the modulation of metabolism and atherogenesis. *Diabetes* 56:1198–1209, 2007

From the ¹Research Unit of Diabetology and Endocrinology, Scientific Institute "Casa Sollievo della Sofferenza," San Giovanni Rotondo, Italy; the ²Department of Clinical Sciences, University La Sapienza, Rome, Italy; and the ³Research Division, Joslin Diabetes Center, and the Department of Medicine, Harvard Medical School, Boston, Massachusetts.

Address correspondence and reprint requests to Alessandro Doria, MD, PhD, MPH, Section on Genetics & Epidemiology, Joslin Diabetes Center, One Joslin Place, Boston, MA 02215. E-mail: alessandro.doria@joslin.harvard.edu; or to Claudia Menzaghi, PhD, Research Unit of Diabetology and Endocrinology, Scientific Institute "Casa Sollievo della Sofferenza," Viale Padre Pio, 71013 San Giovanni Rotondo (FG), Italy. E-mail: c.menzaghi@operapadrepio.it.

Received for publication 13 April 2006 and accepted in revised form 2 February 2007.

Published ahead of print at <http://diabetes.diabetesjournals.org> on 15 February 2007. DOI: 10.2337/db06-0506.

AdipoR1, adiponectin receptor 1; AdipoR2, adiponectin receptor 2; CAD, coronary artery disease; HOMA_{IR}, homeostasis model assessment of insulin resistance; LD, linkage disequilibrium; SNP, single nucleotide polymorphism; TZD, thiazolidinedione; UTR, untranslated region; WMD, weighted mean difference.

© 2007 by the American Diabetes Association.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Our view of adipose tissue has undergone dramatic changes over the past 15 years. Previously believed to be a mere energy depot, adipose tissue is now considered a major endocrine organ regulating whole-body metabolism as well as inflammatory and immune responses (1,2). These actions are mediated by a number of molecules—collectively known as adipokines—that are secreted by adipocytes and act in an autocrine, paracrine, or endocrine fashion, adapting metabolic fluxes to the amount of stored energy (1,2). The discovery of such endocrine function of the adipose tissue has prompted the hypothesis that a genetic dysregulation of the adipokine network may contribute to the pathogenesis of insulin resistance and related disorders such as type 2 diabetes and cardiovascular disease. Of all the molecules that have been shown to be produced by the adipose tissue, adiponectin has drawn special attention, largely due to its effects on both insulin sensitivity and inflammation, and the fact that its expression and serum levels can be modulated by peroxisome proliferator-activated receptor (PPAR)- γ agonists drugs (Fig. 1). In this article, we will review the evidence that has been thus far gathered on the role of genetic variants in the adiponectin and adiponectin receptors genes as modulators of adiponectin-circulating levels and susceptibility to insulin resistance traits. We will also discuss the directions in which research on this topic is heading.

ADIPONECTIN: A SALUTARY ADIPOKINE

Adiponectin, also known as adipocyte complement-related protein 30 (Acrp30), gelatin-binding protein 28 (Gbp28), adipose most abundant transcript 1 (apM1), or AdipoQ, is exclusively produced by adipocytes (3–6). It is abundantly present in serum, where it circulates in two higher-order forms: a low-molecular weight dimer of trimers and a larger high-molecular weight complex of 12–18 subunits (7). Serum levels are 15% higher in women than in men (8). Data from both animal and human studies indicate that adiponectin has insulin-enhancing as well as anti-inflammatory actions (rev. in 9). Adiponectin levels are markedly reduced in obese/diabetic mice, and injection of the adiponectin globular domain to these animals ameliorates insulin resistance, an effect that can be ascribed to an enhancement of fatty acid- β oxidation in skeletal muscle and a decrease of hepatic gluconeogenesis mediated by AMP-activated protein kinase (10–13). Furthermore, adiponectin knockout mice show increased susceptibility to diet-induced insulin resistance as well as injury-induced arterial stenosis and neointimal formation (14,15). Consistent with the results from animal models,




FIG. 1. Proposed salutary effects of adiponectin.

several cross-sectional studies have shown an association between low serum adiponectin levels and insulin resistance, type 2 diabetes, and cardiovascular disease in humans (16–19). A similar inverse relationship between adiponectin levels and incidence of insulin resistance, type 2 diabetes, and atherosclerosis has been demonstrated in follow-up studies (20–25). While it cannot be excluded that also in these studies low adiponectin levels are due to underlying disease processes, such prospective data suggest that hypoadiponectinemia is a determinant rather than a consequence of these conditions.

The salutary effects of adiponectin on metabolic traits and vascular functions have raised the hypothesis that genetic variants decreasing the production or affecting the function of this molecule or of its receptors may contribute to the etiology of insulin resistance and the chronic disorders that are frequently associated with this metabolic abnormality. The gene coding for adiponectin, officially named *ADIPOQ*, is placed on chromosome 3q27. This genomic region has been shown to segregate with insulin resistance traits and type 2 diabetes in several family studies (that is, these traits were more similar in relatives who shared this chromosomal region than those who did not), providing a further rationale for studying this gene as a candidate for these conditions (26–29). The *ADIPOQ* gene includes three exons, spanning a total of 16 kb of genomic sequence (Fig. 2). A recent, systematic analysis of this locus in Europeans suggests that this gene is organized in two linkage disequilibrium (LD) blocks separated by a region of looser LD placed in the middle of the first intron (30) (Fig. 2). A similar two-block structure has been observed in Chinese (31) and Hispanics, although

the between-block boundaries are shifted by a few kilobasepairs on the 5' side in the latter population (32).

During the past 5 years, several polymorphisms at this locus have been repeatedly tested, individually or in combination as haplotypes, for association with low adiponectin levels, measures of adiposity, features of the insulin resistance syndrome, and type 2 diabetes, in both Caucasians and Japanese individuals. Studies have mostly concerned four single nucleotide polymorphisms (SNPs), which were among the first to be discovered by targeted resequencing efforts (33,34). Two of these (g. –11391G→A and g. –11377C→G) are placed in the first LD block, in the immediate 5' flanking region of the gene, and the other two (g. +45T→G and g. +276G→T) are placed in the second LD block, in exon 2 and intron 2, respectively. In general, the results of these studies have been contradictory with regard to whether variability at this locus has an impact on metabolic phenotypes and which polymorphisms are responsible for such an effect (30,32,34–57). Some of these discrepancies may relate to the small size of several of these studies, providing limited power to study moderate genetic effects, whereas others may reflect genuine differences between populations in genetic background or environmental exposures. However, despite these problems, some consistent patterns of association can be identified through meta-analyses as described below for different phenotypes.

IMPACT OF *ADIPOQ* GENE VARIANTS ON ADIPONECTIN CIRCULATING LEVELS

It is estimated that a substantial proportion (between 30 and 70%) of the variability in plasma adiponectin levels is




FIG. 2. Schematic representation of the adiponectin (*ADIPOQ*) gene. The exon-intron organization of the gene is indicated by boxes and lines; closed and open boxes represent the exons for coding and noncoding regions, respectively. The arrows show the positions of polymorphic variants that have been commonly studied. The pairwise LD between SNPs (r^2) is indicated in the triangle at the bottom of the figure (red = 1.0, yellow = 0.7–0.9, green = 0.4–0.6, blue = 0.1–0.3, black = 0.0). Adapted from Heid et al. (30).

accounted for by genetic factors (54,56,58–63). A recent study of 1,700 Caucasian subjects has shown that polymorphisms in both blocks of the *ADIPOQ* gene contribute to such genetic modulation (30). Other studies, however, have failed to detect such effects or have provided contradictory results with regard to the polymorphic sites that are involved. To summarize the results obtained to date and draw some conclusions, albeit provisional, on the role of adiponectin variants in the control of adiponectin levels, we conducted a systematic meta-analysis of all published data concerning SNPs for which a total of >2,000 individuals have been studied. Relevant articles were identified by means of a PubMed search using the phrase “[*ADIPOQ* or adiponectin polymorphisms] and [adiponectin].” Additional studies were collected by searching the reference lists from the publications identified by the PubMed search. Reports on children and on adults undergoing treatments that could possibly interfere with adiponectin levels were excluded. Three SNPs (g.-11391G→A, g.+45T→G, and g.+276G→T) fulfilled these criteria (33,34,36,41,43,44,46,49,50,55,62–66). For each of these variants, a multivariate, random-effect meta-analysis was performed to estimate the weighted mean differences (WMDs) in adiponectin levels between genotypes along with their 95% CIs. Between-study heterogene-

ity and the possible presence of publication bias were assessed by the Cochran’s *Q* test and Macaskill’s inverse pooled variance weighting method, respectively (67,68).

A significant association with adiponectin levels was observed for SNPs g.-11391G→A and g.+276G→T (Table 1). These are probably two independent genetic effects, since the two SNPs belong to different LD blocks (Fig. 2). No association was observed for SNP g.+45T→G (Table 1). SNP g.-11391G→A was associated with adiponectin levels according to a dominant model with allele A carriers (i.e., GA+AA) having higher adiponectin levels compared with GG carriers. Although these data must be taken with caution because of a possible publication bias ($P = 0.04$) (Table 1), they are compatible with recent *in vitro* data supporting a biological function of this SNP, with the allele A enhancing the *ADIPOQ* promoter activity, probably through the displacement of an as yet unidentified transcription factor with inhibitory properties (69).

Results are not as clear cut for SNP g.+276T→G. Adiponectin levels seem to progressively increase from GG homozygotes to heterozygotes and TT homozygotes, but the statistical significance of this association is marginal and is present only for heterozygotes, probably because of the small number of TT subjects. The lack of significance for TT homozygotes may also be related to the

TABLE 1
Meta-analysis of the association between SNPs *g.*-11391G>A, *g.*+45T→G, and *g.*+276G→T and circulating adiponectin levels

Marker	Study (ref.)	Subjects (<i>n</i>)	WMD (95% CI)	WMD (95% CI)	<i>P</i>	<i>P</i> (hetero- geneity)	<i>P</i> (Publica- tion bias)
<i>g.</i> -11391G→A	Fumeron (36)*	453	A/A+A/G vs. G/G 2.39 (0.43–4.34)				
	Poitou (66)*	65	4.60 (–0.31 to 9.51)				
	Schwarz (50)*	550	1.69 (0.73–2.65)				
	Tanko (81)*	283	6.60 (1.08–12.12)				
	Vasseur (55)*	703	1.12 (0.64–1.60)				
A/A +A/G vs. G/G		2,054	1.64 (0.88–2.41)		0.014	0.11	0.04
<i>g.</i> +45T→G			T/G vs. T/T	G/G vs. T/T			
	Fumeron (36)*	457	1.68 (–0.85 to 4.21)	2.78 (–4.05 to 9.61)			
	Jang (41)†	902	0.01 (–0.55 to 0.57)	0.49 (–0.35 to 1.33)			
	Lee (43)†	427	2.35 (1.58–3.12)	1.33 (0.14–2.52)			
	Mackevics (44)*	1,733	0.55 (0.14 to –0.96)	2.53 (1.17–3.89)			
	Menzaghi (62)*	399	2.80 (–3.63 to 9.23)	–3.00 (–16.09 to 10.09)			
	Pollin (63)‡	548	1.50 (–0.79 to 3.79)	3.30 (–2.16 to 8.76)			
	Takahashi (33)†	142	–0.50 (–2.16 to 1.16)	–2.10 (–4.31 to 0.11)			
	Takahashi (33)†	77	–0.20 (–1.83 to 1.43)	–1.10 (–3.27 to 1.07)			
	Tanko (81)*	281	–1.10 (–6.05 to 3.85)	–12.00 (–16.67 to –7.33)			
Vasseur (55)*	703	–0.03 (–0.52 to 0.46)	–0.13 (–1.25 to 0.99)				
T/G vs. T/T		5,388	0.35 (–0.53 to 1.23)		0.44	<0.00001	0.48
G/G vs. T/T				–0.38 (–2.85 to 2.09)	0.76	<0.00001	0.28
<i>g.</i> +276G→T			G/T vs. G/G	T/T vs. G/G			
	Fumeron (36)*	452	–1.36 (–3.43 to 0.71)	–1.66 (–5.60 to 2.28)			
	Hara (34)†	480	1.30 (–0.32 to 2.92)	3.10 (0.24–5.96)			
	Iwashima (64)†	312	0.40 (–0.15 to 0.95)	–0.50 (–1.38 to 0.38)			
	Jang (41)†	902	0.21 (–0.34 to 0.76)	1.45 (0.58–2.32)			
	Lee (43)†	427	–0.70 (–1.52 to 0.12)	–6.65 (–7.23 to –6.07)			
	Mackevics (44)*	1,733	0.45 (0.11–0.79)	1.17 (0.56–1.78)			
	Menzaghi (62)*	403	1.80 (–4.23 to 7.83)	12.70 (2.71–22.69)			
	Mousavinasab (65)*	244	1.60 (0.69–2.51)	4.60 (2.28–6.92)			
	Pollin (63)‡	544	0.80 (–1.39 to 2.99)	3.00 (–0.06 to 6.06)			
	Salmenniemi (49)	156	–0.15 (–1.52 to 1.22)	–0.16 (–1.84 to 1.52)			
	Tanko (81)*	282	1.80 (–1.92 to 5.52)	7.50 (–0.42 to 15.42)			
	Vasseur (55)*	635	0.59 (0.07–1.11)	0.62 (–0.11 to 1.35)			
G/T vs. G/G		6,570	0.39 (0.01–0.79)		0.05	0.043	0.95
T/T vs. G/G				1.10 (–1.00 to 3.19)	0.31	0.00001	0.12

Race/ethnicity: *Caucasian, †Asian, ‡Old Order Amish. Ref. 33 appears twice for the two different study populations included in that report.

highly significant between-study heterogeneity in the data concerning the comparison between TT and GG carriers ($P = 0.00001$) (Table 1), which prompts caution in the interpretation of these findings. The cause of such heterogeneity (observed also for SNP *g.*+45) is not known, but it may relate to the different levels of circulating adiponectin reported in different studies, reflecting disparities in the criteria used to recruit the study populations and/or in the assays used to measure adiponectin concentrations.

If we assume that SNP *g.*+276G→T is indeed associated with adiponectin levels, it seems unlikely that this polymorphism is per se responsible for an effect on adiponectin expression, given the results of a recent in vitro study showing no difference between alleles in the binding with adipocyte nuclear factors (69). Thus, SNP *g.*+276T→G may be a mere marker in LD with an as yet unidentified functional variant. Of note, SNP *g.*+276G→T is in moderate to strong LD ($r^2 > 0.50$) with several polymorphisms placed in the 3' untranslated region (UTR)—a region known for playing a pivotal role in the control of gene expression by binding proteins that regulate mRNA processing, translation, or degradation (70). In particular, an insertion/deletion polymorphism at position +2019 in the

3'UTR (having an r^2 of 0.58 with *g.*+276G→T) is a strong candidate for a functional role, given that most of the linkage between *ADIPOQ* locus and adiponectin circulating levels observed in family study appears to be accounted for by this variant (63). Functional studies in transfected cells examining allelic difference in mRNA stability are needed to investigate the role of 3'UTR SNPs on the modulation of adiponectin levels.

In addition to common SNPs, also several rare variants including G84R, G90S, I164T, and R112C have been reported as modulators of adiponectin levels (54,64,71). The R112C and I164T substitutions, which are associated with low adiponectin levels, appear to disrupt the assembly of adiponectin trimers, which may, in turn, impair adiponectin secretion from adipocytes (72). By contrast, both the G84R and G90S variants have been shown to hamper the formation of high-molecular weight multimers, which are deemed to be essential for adiponectin action (72). The functional link between these sequence differences and adiponectin levels, if any, must be further investigated by functional studies in transfected adipose cells.

Although the results of the meta-analysis indicate a role for variants in the *ADIPOQ* gene in the modulation of

TABLE 2
Meta-analysis of the association between SNPs g.+45T→G and g.+276G→T and HOMA_{IR}

Marker	Study (ref.)	Subjects (n)	WMD (95% CI)	WMD (95% CI)	P	P (heterogeneity)	P (publication bias)
<i>g. +45T→G</i>							
			T/G vs. T/T	G/G vs. T/T			
	Hara (34)†	480	-0.04 (-0.19 to 0.11)	0.04 (-0.20 to 0.28)			
	Jang (41)†	902	0.14 (-0.07 to 0.35)	-0.15 (-0.39 to 0.09)			
	Lee (43)†	427	0.08 (-0.14 to 0.30)	-0.07 (-0.34 to 0.48)			
	Menzaghi (45)*	399	-0.10 (-0.30 to 0.10)	-0.18 (-0.57 to 0.21)			
	Nakatani (46)†	194	0.08 (-0.12 to 0.28)	0.35 (0.00-0.70)			
	Tanko (81)*	281	-0.41 (-1.05 to 0.23)	-0.45 (-2.54 to 1.64)			
	Yang (82)†	245	-0.40 (-0.97 to 0.17)	-0.83 (-1.44 to -0.22)			
T/G vs. T/T		2,928	-0.02 (-0.12 to 0.08)		0.71	0.27	0.10
G/G vs. T/T				-0.09 (-0.36 to 0.17)	0.48	0.036	0.93
<i>g. +276G→T</i>							
			G/T vs. G/G	T/T vs. G/G			
	Hara (34)†	480	-0.16 (-0.30 to -0.02)	-0.42 (-0.66 to -0.18)			
	Jang (41)†	902	-0.03 (-0.23 to 0.17)	-0.36 (-0.59 to -0.13)			
	Lee (43)†	427	0.04 (-0.18 to 0.26)	0.09 (-0.34 to 0.52)			
	Menzaghi (45)*	403	-0.23 (-0.44 to -0.02)	-0.35 (-0.65 to -0.05)			
	Nakatani (46)†	194	-0.15 (-0.36 to 0.06)	-0.39 (-0.62 to -0.16)			
	Tanko (81)†	282	-0.20 (-0.92 to 0.52)	-0.81 (-1.71 to 0.09)			
G/T vs. G/G		2,688	-0.12 (-0.20 to -0.04)		0.005	0.49	0.26
T/T vs. G/G				-0.36 (-0.47 to -0.24)	<0.0001	0.37	0.85

Race/ethnicity: *Caucasian, †Asian.

adiponectin secretion, it must be emphasized that variability at this locus explains only a small proportion (~2–8%) of serum adiponectin variance (30,54,56,60,62,63). If one considers that between 30 and 70% of this variance is accounted for by genetic factors, other loci are likely to play a bigger role. Consistent with such a scenario is the recent observation of a major locus for adiponectin levels on 3q27 that appears to be distinct from the *ADIPOQ* gene (60). The existence of additional genetic modulators is also supported by the observation of linkage with adiponectin levels at locations other than 3q27 (58,59,61–63). Among these, the region of 14q13 is especially interesting since linkage has been detected at this location in two independent studies (59,62). Of note, the linkage at these loci is not attenuated by adjustment for BMI, suggesting that the effect of these genes on adiponectin levels is not secondary to an effect on adiposity (60–62). By contrast, the linkage signal observed on chromosome 15 in Chinese decreases significantly after BMI adjustment, indicating that the gene(s) at this locus may modulate adiponectin levels through an effect on adipose mass (58). Genes with an effect on adiponectin levels may also include known modulators of insulin sensitivity such as PPAR- γ 2 and tumor necrosis factor (TNF)- α , although conflicting results have been obtained in different populations (73–77).

IMPACT OF *ADIPOQ* GENE VARIANTS ON BMI, INSULIN RESISTANCE, AND TYPE 2 DIABETES

Low plasma adiponectin levels have been consistently associated with body weight, insulin resistance, and increased risk of type 2 diabetes in cross-sectional as well as prospective studies (16–18,20–22). Furthermore, adiponectin has been shown in animal models to decrease body weight and to independently increase insulin sensitivity (11,13). Thus, one would expect that the same polymorphisms that are associated with adiponectin levels would also display an association with metabolic traits. Such prediction has not been unequivocally confirmed. In

the first LD block, a haplotype characterized by a G at positions g.-11391 and g.-11377 was found to be associated with type 2 diabetes in a case-control study of >2,000 French Caucasians (54,55). This association has been replicated with regard to SNP g.-11377C→G in studies of populations from Japan and Sweden (39,47), but not in other studies (37,56). In the second block, positive findings have been reported for SNPs at positions g.+45, g.+276, g.+712, g.+3,639, and g.+4,545, which are in strong LD with each other ($r^2 > 0.8$ for SNPs g.+45, + g.+712, and g.+2019; $r^2 > 0.8$ for g.+276 and g.+3,696) (Fig. 2), as well as for less frequent variants such as G84R, G90S, I164T, and R112C. These sequence differences have been variably associated with increased BMI, insulin resistance-related traits, and type 2 diabetes in cross-sectional studies of populations of different origin (34,35,38,40–43,45,46, 51,53). Some of these variants have also been reported to influence weight gain, impaired glucose tolerance, and type 2 diabetes risk in prospective studies (31,36,50,78). Also in this case, however, results have been far from uniform, being characterized by the same lack of replication displayed by genetic studies of other complex traits.

To better understand whether *ADIPOQ* variants are genuine determinants of metabolic traits, we conducted a meta-analysis using an approach similar to that described above for adiponectin levels. Applying the same criterion of having access to at least 2,000 subjects per SNP-phenotype combination, sufficient data were available in the literature for SNPs g.-11391G→A and g.-11377C→G with regard to type 2 diabetes (34,36,37,39,40,47,54–56,79), and for SNPs g.+45T→G and g.+276G→T with regard to BMI, insulin resistance, and type 2 diabetes (34,36–41,43,45–47,51–57,65,79–82). The only evidence for a significant association was observed between SNP g.+276G→T and insulin resistance, as defined by the surrogate marker HOMA_{IR} (homeostasis model assessment of insulin resistance) (Table 2). This index was higher in +276 GG carriers as compared with TG and TT subjects, indicating higher insulin sensitivity

in carriers of allele T—the same allele that showed an association, albeit marginal, with increased adiponectin levels (Table 2). All the studies were homogeneous, and no publication bias was present (Table 2). No significant global effects were observed between any of the SNPs and risk of type 2 diabetes (Table 3) or higher BMI (Table 4), although these data must be taken with caution since evidence of between-study heterogeneity and/or publication bias was detected for some SNP–phenotype combinations (Tables 3 and 4). Such negative results are in apparent contrast with those reported by recent prospective studies on the role of SNP *g.*+45T→G on weight gain (36) and SNPs *g.*-11391G→A, *g.*-11377G→C, and *g.*+45T→G on type 2 diabetes (50,78). The reasons for such discrepancies are unknown, but in addition to the obvious possibility of false positives due to chance or bias, one should also consider the role of differences among populations in environmental exposures that may be critical for genetic effects to become manifest. Differences in study design (i.e., cross-sectional versus prospective) must also be taken into account, given the possibility of survival bias obscuring genuine genetic effects in cross-sectional studies.

IMPACT OF *ADIPOQ* GENE VARIANTS ON THE RISK OF CORONARY ARTERY DISEASE

Another notable area of investigation is the relationship between genetic variants in the *ADIPOQ* gene and cardiovascular risk. In addition to modulating atherogenesis through its insulin-sensitizing actions, adiponectin has direct anti-atherogenic effects on the arterial wall by inhibiting monocyte adhesion to the endothelium, smooth muscle cell proliferation, and foam cell formation (9,14,15). Administration of recombinant adenovirus expressing human adiponectin to apoE-deficient animals causes a 30% reduction in the formation of atherosclerotic lesions in the absence of any effects on metabolic traits (83,84). Consistent with these findings, men with adiponectin levels in the highest quartile have been shown to have a significantly lower incidence of cardiovascular disease than men with adiponectin levels in the lowest quartile, independently of other cardiovascular risk factors or systemic inflammation markers (85,86).

Studies of the association between adiponectin polymorphisms and cardiovascular outcomes have not been as numerous as those concerning adiponectin levels or metabolic traits, but the overall number of study subjects is such that some preliminary conclusions can be drawn. Four populations, all consisting of subjects with type 2 diabetes, have been examined. The first one, recruited by Lacquemant et al. (87) in Switzerland and France, is a cross-sectional sample of 162 cases with coronary artery disease (CAD) and 315 control subjects without significant CAD. The second one is a similar set of 142 CAD-positive cases and 234 CAD-negative control subjects recruited in Southern Italy (88). The last two consist of incident CAD case and control subjects from the Nurses Health Study (989 women) and the Health Professional Follow-up Study (879 men), both of which are being conducted in the U.S. (89,90). In the French-Swiss population, SNP *g.*+45 was found to be a significant predictor of CAD, with an odds ratio (OR) of 1.9 (95% CI 1.2–2.9) (87). This effect, however, was not confirmed in the other three populations, which instead showed a significant association with SNP *g.*+276 (88–90). When the data from the four populations are considered together in a meta-analysis (including a total of

827 CAD-positive case and 1,887 CAD-negative control subjects), individuals homozygous for allele T at *g.*+276 have about half the cardiovascular risk of allele G carriers (OR 0.55, 95% CI 0.4–0.8) (89). While this result will have to be confirmed in additional populations, it is noteworthy that the pattern of association is consistent with that that is found for this SNP with regard to insulin resistance and adiponectin levels.

ADIPOQ GENE VARIANTS AS POSSIBLE PREDICTORS OF THIAZOLIDINEDIONES EFFICACY

The PPAR- γ agonists thiazolidinediones (TZDs) are potent insulin sensitizers that are increasingly used for the treatment of type 2 diabetes. Several studies have shown that treatment with TZDs increases serum adiponectin, raising the hypothesis that some of the antidiabetic actions of these agents are due to their effects on adiponectin levels (91–94). The hypoglycemic response to TZDs varies in the population, with 30–40% of individuals who have type 2 diabetes or are at risk for type 2 diabetes failing to respond to this therapy (95–99). Given the possible links between the antidiabetic action of TZDs and the effect of these drugs on adiponectin levels, genetic variation in the *ADIPOQ* gene might account for such differences in TZD responsiveness. Studies exploring this hypothesis are in their infancy. In the only published report to date, Kang et al. (80) have found that patients with type 2 diabetes carrying the G allele at either position *g.*+45 or position *g.*+276 were less responsive to the TZD rosiglitazone in terms of both a reduction of fasting plasma glucose and A1C and an increase of circulating adiponectin levels. This result, however, has been obtained in a very small cohort of type 2 diabetic patients. If confirmed in larger populations, these findings may translate into clinical tools, allowing the identification of the best candidates for TZD treatment, with obvious implications for devising more cost-effective approaches to diabetes care.

IMPACT OF GENETIC VARIABILITY IN THE ADIPONECTIN RECEPTORS ON INSULIN RESISTANCE, TYPE 2 DIABETES, AND CORONARY ARTERY DISEASE

The molecules mediating the biological functions of adiponectin were not known until 2003, when adiponectin receptors 1 and 2 (AdipoR1 and AdipoR2) were cloned by Yamauchi et al. (100). In addition to AdipoR1 and AdipoR2, T-cadherin also appears to function as an adiponectin receptor (101). Both AdipoR1 and AdipoR2 are seven-transmembrane domain proteins belonging to a new class of molecules with homology to proteins functioning as progesterin receptors (progesterin/adiponectin/adipoQ receptor, or PAQR, family) (100). In humans, AdipoR1 is ubiquitously expressed, whereas AdipoR2 is predominantly expressed in skeletal muscle and liver (100). Glucose-tolerant Mexican-American subjects with a family history of type 2 diabetes were found to have significantly lower skeletal muscle levels of AdipoR1 and AdipoR2 mRNA than individuals with no family history of diabetes (102). In the same study, the expression of both receptors was positively correlated with the glucose disposal rate as measured by the euglycemic-hyperinsulinemic glucose clamp technique and, in the case of AdipoR2 alone, with plasma adiponectin levels (102). Interestingly, the expression of both receptors in human macrophages is stimulated by TZDs (103). Taken together, these data, though preliminary, suggest that downregulation of AdipoR1 and/or

TABLE 3
Meta-analysis of the association between SNPs g.-11391G→A, g.-11377C→G, g.+45T→G, and g.+276G→T and type 2 diabetes

Marker	Study (ref.)	Type 2 diabetic subjects (n)	Control subjects (n)	OR (95% CI)	OR (95% CI)	P	P (heterogeneity)	P (publication bias)
<i>g.-11391G→A</i>								
				G/A vs. G/G	A/A vs. G/G			
	Fumeron (36)*	177	4,510	0.95 (0.63–1.42)	0.58 (0.08–4.28)			
	Hara (34)†	384	480	0.82 (0.44–1.52)	1.24 (0.02–62.48)			
	Gibson (37)*	775	944	1.22 (0.95–1.57)	1.45 (0.52–4.01)			
	Stenvinkel (79)*	30	145	1.59 (0.44–5.72)	0.53 (0.03–10.16)			
	Vasseur (54)*	620	690	1.21 (0.91–1.62)	2.08 (0.69–6.26)			
	Vasseur (55)*	231	270	1.22 (0.95–1.57)	1.42 (0.38–5.36)			
G/A vs. G/G		2,217	7,039	1.10 (0.94–1.28)		0.23	0.48	0.31
A/A vs. G/G					1.41 (0.78–2.58)	0.26	0.89	0.08
<i>g.-11377C→G</i>								
				C/G vs. C/C	G/G vs. C/C			
	Fumeron (36)*	175	4,502	1.10 (0.80–1.51)	0.68 (0.34–1.35)			
	Gibson (37)*	812	1,044	1.02 (0.84–1.24)	1.05 (0.73–1.51)			
	Gu (39)*	103	486	0.78 (0.49–1.24)	0.88 (0.43–1.80)			
	Hara (34)†	384	480	0.81 (0.61–1.08)	0.74 (0.43–1.27)			
	Hu (40)*	642	995	1.00 (0.82–1.24)	1.08 (0.71–1.65)			
	Polpulaire (47)†	164	183	0.45 (0.29–0.71)	0.90 (0.37–2.22)			
	Stenvinkel (79)*	30	145	1.64 (0.73–3.68)	0.57 (0.07–4.81)			
	Vasseur (54)*	620	691	1.32 (1.05–1.66)	1.30 (0.84–2.02)			
	Vasseur (55)*	231	270	1.48 (1.02–2.14)	1.04 (0.44–2.46)			
	Vojarova-De Courten (56)‡	510	429	0.59 (0.44–0.81)	1.14 (0.78–1.68)			
C/G vs. G/G		3,671	9,225	0.94 (0.75–1.18)		0.60	<0.0001	0.80
C/C vs. G/G					1.03 (0.87–1.22)	0.79	0.84	0.04
<i>g.+45T→G</i>								
				T/G vs. T/T	G/G vs. T/T			
	Fumeron (36)*	177	4,525	1.02 (0.71–1.44)	0.16 (0.01–2.66)			
	Gibson (37)*	740	915	0.94 (0.75–1.17)	0.89 (0.44–1.80)			
	Gonzales-Sanchez (38)*	61	530	1.43 (0.82–2.48)	1.10 (0.24–4.93)			
	Hara (34)†	384	480	1.41 (1.06–1.88)	1.70 (1.09–2.65)			
	Hu (40)*	642	995	0.90 (0.69–1.15)	0.88 (0.35–2.26)			
	Lee (43)†	493	427	0.89 (0.68–1.17)	0.69 (0.43–1.10)			
	Menzaghi (45)*	310	304	0.74 (0.50–1.09)	0.71 (0.26–1.93)			
	Polpulaire (47)†	164	183	1.03 (0.66–1.61)	1.54 (0.74–3.21)			
	Stenvinkel (79)*	26	145	1.60 (0.63–4.06)	0.45 (0.02–8.50)			
	Ukkola (53)*	258	283	0.96 (0.54–1.72)	0.22 (0.01–4.54)			
	Vasseur (54)*	582	680	1.05 (0.81–1.37)	0.78 (0.41–1.49)			
	Vasseur (55)*	231	270	0.68 (0.44–1.03)	0.61 (0.24–1.59)			
	Vojarova-De Courten (56)‡	603	425	1.12 (0.80–1.56)	0.48 (0.08–2.87)			
T/G vs. T/T		4,671	10,162	0.99 (0.90–1.09)		0.86	0.18	0.92
G/G vs. T/T					0.91 (0.68–1.23)	0.54	0.25	<0.001
<i>g.+276G→T</i>								
				G/T vs. G/G	T/T vs. G/G			
	Fumeron (36)*	176	4,497	0.90 (0.65–1.24)	1.23 (0.70–2.14)			
	Gibson (37)*	701	893	0.90 (0.73–1.11)	0.80 (0.55–1.17)			
	Gonzales-Sanchez (38)*	61	530	1.50 (0.86–2.62)	1.39 (0.50–3.85)			
	Hara (34)†	384	480	0.74 (0.56–0.98)	0.46 (0.26–0.83)			
	Hu (40)*	642	995	1.08 (0.88–1.33)	1.20 (0.82–1.75)			
	Lee (43)†	493	427	1.39 (1.06–1.82)	1.09 (0.66–1.79)			
	Menzaghi (45)*	310	304	1.07 (0.77–1.50)	1.05 (0.59–1.86)			
	Polpulaire (47)†	164	177	0.62 (0.39–0.99)	1.11 (0.55–2.22)			
	Stenvinkel (79)*	30	145	0.63 (0.26–1.53)	2.02 (0.61–6.62)			
	Ukkola (53)*	255	283	0.90 (0.62–1.29)	1.07 (0.63–1.82)			
	Vasseur (54)*	577	681	1.11 (0.87–1.41)	0.83 (0.56–1.21)			
	Vasseur (55)*	209	248	1.30 (0.88–1.93)	1.36 (0.72–2.57)			
	Vojarova-De Courten (56)‡	570	433	0.73 (0.55–0.95)	0.73 (0.41–1.30)			
	Yoshioka (57)†	346	98	0.76 (0.48–1.22)	1.07(0.46–2.49)			
G/T vs. G/G		4,918	10,191	0.98 (0.86–1.10)		0.42	0.008	0.64
T/T vs. G/G					0.95 (0.81–1.12)	0.99	0.35	0.32

Race/ethnicity: *Caucasian, †Asian, ‡Pima Indian.

TABLE 4
Meta-analysis of the association between SNPs g.+45T→G and g.+276G→T and BMI

Marker	Study (ref.)	Subjects (n)	WMD (95% CI)	WMD (95% CI)	P	P (heterogeneity)	P (publication bias)
<i>g.+45T→G</i>			T/G vs. T/T	G/G vs. T/T			
	Fumeron (36)*	4,448	0.20 (-0.07 to 0.47)	-0.20 (-1.15 to 0.75)			
	Jang (41)†	902	-0.10 (-0.57 to 0.37)	-0.30 (-1.09 to 0.49)			
	Kang (80)†	166	0.80 (-0.14 to 1.74)	0.10 (-1.11 to 1.31)			
	Lee (43)†	427	-0.30 (-0.92 to 0.32)	-0.40 (-1.16 to 0.36)			
	Menzaghi (45)*	399	-0.70 (-1.60 to 0.20)	-0.40 (-2.37 to 1.57)			
	Nakatani (46)†	194	0.80 (-0.12 to 1.72)	1.54 (-0.04 to 3.12)			
	Stumvoll (51)*	371	1.40 (-0.09 to 2.89)	2.30 (-0.31 to 4.91)			
	Tanko (81)*	1,380	0.00 (-0.51 to 0.51)	-0.80 (-3.12 to 1.52)			
	Yang (82)†	245	-1.41 (-3.26 to 0.44)	-3.35 (-6.00 to -0.70)			
T/G vs. T/T		8,287	0.09 (-0.09 to 0.28)		0.32	0.046	0.53
G/G vs. T/T				-0.18 (-0.58 to 0.23)	0.39	0.07	0.95
<i>g.+276G→T</i>			G/T vs. G/G	T/T vs. G/G			
	Jang (41)†	902	-0.10 (-0.57 to 0.37)	0.20 (-0.64 to 1.04)			
	Kang (80)†	166	0.00 (-0.80 to 0.80)	0.00 (-1.29 to -1.29)			
	Lee (43)†	427	0.50 (-0.12 to 1.12)	0.40 (-0.73 to 1.53)			
	Mousavinasab (65)*	252	0.20 (-0.89 to 1.29)	1.10 (-0.77 to 2.97)			
	Menzaghi (45)*	403	-1.30 (-2.21 to -0.39)	-1.20 (-2.49 to 0.09)			
	Nakatani (46)†	194	-0.98 (-1.92 to -0.04)	-1.79 (-2.95 to -0.63)			
	Tanko (81)*	1,377	0.20 (-0.22 to 0.62)	-0.10 (-0.82 to 0.62)			
	Ukkola (52)*	95	2.10 (0.69-3.51)	0.20 (-2.09 to 2.49)			
G/T vs. G/G		3,816	-0.01 (-0.59 to 0.57)		0.98	0.001	0.60
T/T vs. G/G				-0.15 (-0.80 to 0.50)	0.65	0.06	0.69

Race/ethnicity: *Caucasian, †Asian.

AdipoR2 might contribute to the development of insulin resistance in peripheral tissues.

The genes coding for AdipoR1 and AdipoR2 are placed on chromosomes 1q32.1 and 12p13.33, respectively (100). Both have been studied as candidate genes for insulin resistance and related disorders, although less extensively and with even more conflicting results than for the *ADIPOQ* gene. Wang et al. (104) screened the *ADIPOR1* gene for sequence variants in Caucasian and African-American subjects but failed to find associations between any of the identified polymorphisms and either insulin resistance or type 2 diabetes. Similarly, negative results were obtained by Hara et al. (105), who failed to observe any association between *ADIPOR1* or *ADIPOR2* gene variants and type 2 diabetes in a French Caucasian population. More recently, however, three SNPs at the *ADIPOR2* locus have been reported to confer a modest increase in the risk of type 2 diabetes in a French population (106). Multiple polymorphisms in both *ADIPOR1* and *ADIPOR2* have been also associated with type 2 diabetes and related traits in Old Order Amish (107), and two SNPs in the *ADIPOR1* gene (i.e., g.-8503G→A in the promoter and g.-1927T→C in intron 1) have been associated with insulin resistance features and amount of fat in the liver in Caucasians from Germany (108). The association between *ADIPOR1* gene variants and insulin resistance features, but not with the risk of type 2 diabetes, has been confirmed also in Finnish (109) and Mexican Americans (110). *ADIPOR2* gene variants (i.e., g.+795G→A, g.+870C→A, and g.+963C→T) have also been recently associated as a haplotype with plasma adiponectin levels (111). Finally, recent data from our laboratories indicate that polymorphisms in the 3'UTR and 3' flanking region of *ADIPOR1* may play a major role in the modulation of atherogenesis, conferring a threefold increase in the risk of CAD among individuals with type 2

diabetes (112). This effect seems to be related to a lower *ADIPOR1* expression in carriers of the at-risk genotypes, possibly determining a deficit of the antiatherogenic effects of adiponectin on vascular cells (112).

Despite this encouraging evidence, the number of studies published to date is still too small to draw firm conclusions on the role of variability in *ADIPOR1* and/or *ADIPOR2* in predicting insulin resistance and related disorders. In particular, the strong effect that we have demonstrated on cardiovascular risk must be confirmed in prospective studies before *ADIPOR1* polymorphisms can be considered as clinically useful predictors of increased susceptibility to CAD.

CONCLUSIONS AND FUTURE DIRECTIONS

Many reports have been published to date concerning the metabolic effects of variation in the gene coding for adiponectin (30-32,34-36,38-47,50-55,62-66,69,71,78,79,81,82,87-90,110). Meta-analyses of these studies support the hypothesis that variability in this gene contribute to the modulation of circulating adiponectin levels and the risk of insulin resistance and CAD. Two independent genetic effects, corresponding to the two distinct LD blocks observed at this locus, appear to be present (30). The effect corresponding to the 5' block, epitomized by SNP g.-11391G→A, can be demonstrated for plasma adiponectin levels, although its statistical significance is not especially striking, considering that these data were derived from >2,000 individuals. The effect of the 3' block, captured by SNP g.+276, is especially strong for insulin resistance and CAD, whereas it is only marginally significant for adiponectin levels. No consistent effect on BMI or risk of type 2 diabetes can be demonstrated in either block. Genetic variants in the genes coding for adiponectin receptors may also influence the risk of

insulin resistance and cardiovascular disease (106–112), but data on these genes are still too sparse to draw meaningful conclusions.

Despite the progress in this field, several aspects remain unresolved. One of these concerns the significant evidence of heterogeneity that emerges from several of the meta-analyses. At this stage, it is unclear whether these differences in association across studies reflect differences in genetic background or modifying environmental factors between populations are the result of discrepancies in ascertainment criteria or phenotyping methods or simply represent statistical fluctuations. Large collaborative studies including thousands of individuals from a few different ethnic groups, recruited and phenotyped according to standard protocols and genotyped for the same set of SNPs, are clearly needed to conclusively address this matter.

One aspect that deserves further investigation is the inconsistency between the results concerning the association with adiponectin levels and those concerning the association with metabolic traits. For instance, SNP g. –11391G→A displays the strongest association with adiponectin levels, yet it does not seem to be associated with type 2 diabetes or CAD. Conversely, SNP g. +276 shows a rather weak association with adiponectin levels, yet it appears to have a strong effect on the risk of insulin resistance and CAD. The reasons for such discrepancies are not clear at this time, but an important point to consider is that serum levels may not necessarily reflect the overall amount of adiponectin in the body or its concentration in the interstitial space where the targets for the insulin-sensitizing or antiatherogenic effect of this adipokine are located. Thus, studies considering the impact of SNPs on adiponectin production (determined for instance by measuring adiponectin expression in adipocytes) rather than its serum concentrations may be more revealing.

Another unresolved issue concerns the identities of the functional variants that are responsible for the association with adiponectin levels or metabolic phenotypes. While the genetic effect in the 5' block seems to be secondary to differences in promoter activation due to SNP g. –11391G→A (69), the situation is more complex for the 3' block where the marker displaying association with metabolic traits (i.e., SNP g. +276) does not have obvious functions. It is important to consider, however, that only a small proportion of *ADIPOQ* variants have been analyzed to date, and it is therefore possible that other as yet uninvestigated polymorphisms are responsible for the associations with metabolic traits. Thus, there is the need for a thorough characterization of the repertoire of polymorphisms in either block followed by *in vitro* studies systematically investigating the biological function of each sequence difference in strong LD with the two markers associated with metabolic phenotypes.

As these additional studies proceed, new insights are expected by the completion of the whole genome association studies that are currently in progress in a variety of populations. These efforts are expected to lead to the identification of additional genes involved in the regulation of serum adiponectin levels and, thereby, explain the proportion of genetic variance of this trait that is not accounted for by SNPs at the adiponectin and adiponectin receptor loci. With this novel knowledge, studies investigating the interaction between the *ADIPOQ* and other

genes in the etiology of the insulin resistance syndrome and related disorders will become a reality.

Another area of research that is expected to grow concerns the role of specific variants at the adiponectin and adiponectin receptor loci in modulating interindividual differences in the response to TZD treatment. While the data available in the literature are promising (80), more evidence must be gathered from properly designed pharmacogenetic studies evaluating the outcome of TZD treatment in carriers of different genotypes. If positive and replicable data are obtained, it may be possible to use the genetic information concerning these loci to build algorithms allowing physicians to tailor antidiabetes therapy to the patients' genetic features, advancing diabetes care into the realm of personalized medicine.

ACKNOWLEDGMENTS

Part of the research described in this article was supported by Accordo Programma Quadro in Materia di Ricerca Scientifica nella Regione Puglia-PST 2006 and Italian Ministry of Health grants Ricerca Corrente 2003, 2004, 2005, 2006, and 2007 (to C.M.) and Ricerca Finalizzata 2002 (to V.T.); by Ministry of University and Scientific Research FIRB RBNE01C5S2_005 (to V.T.) and PRIN 2003 (to V.T.); by National Institutes of Health grants HL73168, HL71981, DK55523, and DK60837 (to A.D.); and by the Genetics Core of the Diabetes and Endocrinology Research Center at the Joslin Diabetes Center (DK36836).

The authors gratefully acknowledge F. Pellegrini for statistical support in meta-analyses.

REFERENCES

1. Spiegelman BM, Flier JS: Obesity and the regulation of energy balance. *Cell* 104:531–543, 2001
2. Saltiel AR: You are what you secrete. *Nat Med* 7:887–888, 2001
3. Scherer PE, Williams S, Fogliano M, Baldini G, Lodish HF: A novel serum protein similar to C1q, produced exclusively in adipocytes. *J Biol Chem* 270:26746–26749, 1995
4. Nakano Y, Tobe T, Choi-Miura NH, Mazda T, Tomita M: Isolation and characterization of GBP28, a novel gelatin-binding protein purified from human plasma. *J Biochem (Tokyo)* 120:803–812, 1996
5. Maeda K, Okubo K, Shimomura I, Funahashi T, Matsuzawa Y, Matsubara K: cDNA cloning and expression of a novel adipose specific collagen-like factor, apM1 (AdiPose Most abundant Gene transcript 1). *Biochem Biophys Res Commun* 221:286–289, 1996
6. Hu E, Liang P, Spiegelman BM: AdipoQ is a novel adipose-specific gene dysregulated in obesity. *J Biol Chem* 271:10697–10703, 1996
7. Fruebis J, Tsao TS, Javorschi S, Ebbels-Reed D, Erickson MR, Yen FT, Bihain BE, Lodish HF: Proteolytic cleavage product of 30-kDa adipocyte complement-related protein increases fatty acid oxidation in muscle and causes weight loss in mice. *Proc Natl Acad Sci U S A* 98:2005–2010, 2001
8. Arita Y, Kihara S, Ouchi N, Takahashi M, Maeda K, Miyagawa J, Hotta K, Shimomura I, Nakamura T, Miyaoka K, Kuriyama H, Nishida M, Yamashita S, Okubo K, Matsubara K, Muraguchi M, Ohmoto Y, Funahashi T, Matsuzawa Y: Paradoxical decrease of an adipose-specific protein, adiponectin, in obesity. *Biochem Biophys Res Commun* 257:79–83, 1999
9. Kadowaki T, Yamauchi T: Adiponectin and adiponectin receptors. *Endocr Rev* 26:439–451, 2005
10. Yamauchi T, Kamon J, Waki H, Terauchi Y, Kubota N, Hara K, Mori Y, Ide T, Murakami K, Tsuboyama-Kasaoka N, Ezaki O, Akanuma Y, Gavrilova O, Vinson C, Reitman ML, Kagechika H, Shudo K, Yoda M, Nakano Y, Tobe K, Nagai R, Kimura S, Tomita M, Froguel P, Kadowaki T: The fat-derived hormone adiponectin reverses insulin resistance associated with both lipotrophy and obesity. *Nat Med* 7:941–946, 2001
11. Maeda N, Shimomura I, Kishida K, Nishizawa H, Matsuda M, Nagaretani H, Furuyama N, Kondo H, Takahashi M, Arita Y, Komuro R, Ouchi N, Kihara S, Tochino Y, Okutomi K, Horie M, Takeda S, Aoyama T, Funahashi T, Matsuzawa Y: Diet-induced insulin resistance in mice lacking adiponectin/ACRP30. *Nat Med* 8:731–737, 2002
12. Berg AH, Combs TP, Du X, Brownlee M, Scherer PE: The adipocyte-

- secreted protein Acrp30 enhances hepatic insulin action. *Nat Med* 7:947–953, 2001
13. Combs TP, Pajvani UB, Berg AH, Lin Y, Jelicks LA, Laplante M, Nawrocki AR, Rajala MW, Parlow AF, Cheeseboro L, Ding YY, Russell RG, Lindemann D, Hartley A, Baker GR, Obici S, Deshaies Y, Ludgate M, Rossetti L, Scherer PE: A transgenic mouse with a deletion in the collagenous domain of adiponectin displays elevated circulating adiponectin and improved insulin sensitivity. *Endocrinology* 145:367–383, 2004
 14. Kubota N, Terauchi Y, Yamauchi T, Kubota T, Moroi M, Matsui J, Eto K, Yamashita T, Kamon J, Satoh H, Yano W, Froguel P, Nagai R, Kimura S, Kadowaki T, Noda T: Disruption of adiponectin causes insulin resistance and neointimal formation. *J Biol Chem* 277:25863–25866, 2002
 15. Matsuda M, Shimomura I, Sata M, Arita Y, Nishida M, Maeda N, Kumada M, Okamoto Y, Nagaretani H, Nishizawa H, Kishida K, Komuro R, Ouchi N, Kihara S, Nagai R, Funahashi T, Matsuzawa Y: Role of adiponectin in preventing vascular stenosis: the missing link of adipo-vascular axis. *J Biol Chem* 277:37487–37491, 2002
 16. Hotta K, Funahashi T, Arita Y, Takahashi M, Matsuda M, Okamoto Y, Iwahashi H, Kuriyama H, Ouchi N, Maeda K, Nishida M, Kihara S, Sakai N, Nakajima T, Hasegawa K, Muraguchi M, Ohmoto Y, Nakamura T, Yamashita S, Hanafusa T, Matsuzawa Y: Plasma concentrations of a novel, adipose-specific protein, adiponectin, in type 2 diabetic patients. *Arterioscler Thromb Vasc Biol* 20:1595–1599, 2000
 17. Weyer C, Funahashi T, Tanaka S, Hotta K, Matsuzawa Y, Pratley RE, Tataranni PA: Hypoadiponectinemia in obesity and type 2 diabetes: close association with insulin resistance and hyperinsulinemia. *J Clin Endocrinol Metab* 86:1930–1935, 2001
 18. Yang WS, Lee WJ, Funahashi T, Tanaka S, Matsuzawa Y, Chao CL, Chen CL, Tai TY, Chuang LM: Weight reduction increases plasma levels of an adipose-derived anti-inflammatory protein, adiponectin. *J Clin Endocrinol Metab* 86:3815–3819, 2001
 19. Kumada M, Kihara S, Sumitsuji S, Kawamoto T, Matsumoto S, Ouchi N, Arita Y, Okamoto Y, Shimomura I, Hiraoka H, Nakamura T, Funahashi T, Matsuzawa Y: Association of hypoadiponectinemia with coronary artery disease in men. *Arterioscler Thromb Vasc Biol* 23:85–89, 2003
 20. Lindsay RS, Funahashi T, Hanson RL, Matsuzawa Y, Tanaka S, Tataranni PA, Knowler WC, Krakoff J: Adiponectin and development of type 2 diabetes in the Pima Indian population. *Lancet* 360:57–58, 2002
 21. Spranger J, Kroke A, Mohlig M, Bergmann MM, Ristow M, Boeing H, Pfeiffer AF: Adiponectin and protection against type 2 diabetes mellitus. *Lancet* 361:226–228, 2003
 22. Snehalatha C, Mukesh B, Simon M, Viswanathan V, Haffner SM, Ramachandran A: Plasma adiponectin is an independent predictor of type 2 diabetes in Asian Indians. *Diabetes Care* 26:3226–3229, 2003
 23. Duncan BB, Schmidt MI, Pankow JS, Bang H, Couper D, Ballantyne CM, Hoogeveen RC, Heiss G: Adiponectin and the development of type 2 diabetes: the atherosclerosis risk in communities study. *Diabetes* 53:2473–2478, 2004
 24. Frystyk J, Berne C, Berglund L, Jensen K, Flyvbjerg A, Zethelius B: Serum adiponectin is a predictor of coronary heart disease: a population-based 10-year follow-up study in elderly men. *J Clin Endocrinol Metab* 92:571–576, 2007
 25. Sattar N, Wannamethee G, Sarwar N, Tchernova J, Cherry L, Wallace AM, Danesh J, Whincup PH: Adiponectin and coronary heart disease: a prospective study and meta-analysis. *Circulation* 114:623–629, 2006
 26. Kissebah AH, Sonnenberg GE, Myklebust J, Goldstein M, Broman K, James RG, Marks JA, Krakower GR, Jacob HJ, Weber J, Martin L, Blangero J, Comuzzie AG: Quantitative trait loci on chromosomes 3 and 17 influence phenotypes of the metabolic syndrome. *Proc Natl Acad Sci U S A* 97:14478–14483, 2000
 27. Vionnet N, Hani El H, Dupont S, Gallina S, Francke S, Dotte S, De Matos F, Durand E, Lepretre F, Lecoecur C, Gallina P, Zekiri L, Dina C, Froguel P: Genome-wide search for type 2 diabetes-susceptibility genes in French whites: evidence for a novel susceptibility locus for early-onset diabetes on chromosome 3q27-qter and independent replication of a type 2-diabetes locus on chromosome 1q21–q24. *Am J Hum Genet* 67:1470–1480, 2000
 28. Francke S, Manraj M, Lacquemant C, Lecoecur C, Lepretre F, Passa P, Hebe A, Corset L, Yan SL, Lahmidi S, Jankee S, Gunnness TK, Ramjuttun US, Balgobin V, Dina C, Froguel P: A genome-wide scan for coronary heart disease suggests in Indo-Mauritians a susceptibility locus on chromosome 16p13 and replicates linkage with the metabolic syndrome on 3q27. *Hum Mol Genet* 10:2751–2765, 2001
 29. Mori Y, Otabe S, Dina C, Yasuda K, Populaire C, Lecoecur C, Vatin V, Durand E, Hara K, Okada T, Tobe K, Boutin P, Kadowaki T, Froguel P: Genome-wide search for type 2 diabetes in Japanese affected sib-pairs confirms susceptibility genes on 3q, 15q, and 20q and identifies two new candidate loci on 7p and 11p. *Diabetes* 51:1247–1255, 2002
 30. Heid IM, Wagner SA, Gohlke H, Iglseider B, Mueller JC, Cip P, Ladurner G, Reiter R, Stadlmayr A, Mackevics V, Illig T, Kronenberg F, Paulweber B: Genetic architecture of the APM1 gene and its influence on adiponectin plasma levels and parameters of the metabolic syndrome in 1,727 healthy Caucasians. *Diabetes* 55:375–384, 2006
 31. Tso AW, Sham PC, Wat NM, Xu A, Cheung BM, Rong R, Fong CH, Xu JY, Cheng KK, Janus ED, Lam KS: Polymorphisms of the gene encoding adiponectin and glycaemic outcome of Chinese subjects with impaired glucose tolerance: a 5-year follow-up study. *Diabetologia* 49:1806–1815, 2006
 32. Sutton BS, Weinert S, Langefeld CD, Williams AH, Campbell JK, Saad MF, Haffner SM, Norris JM, Bowden DW: Genetic analysis of adiponectin and obesity in Hispanic families: the IRAS Family Study. *Hum Genet* 117:107–118, 2005
 33. Takahashi M, Arita Y, Yamagata K, Matsukawa Y, Okutomi K, Horie M, Shimomura I, Hotta K, Kuriyama H, Kihara S, Nakamura T, Yamashita S, Funahashi T, Matsuzawa Y: Genomic structure and mutations in adipose-specific gene, adiponectin. *Int J Obes Relat Metab Disord* 24:861–868, 2000
 34. Hara K, Boutin P, Mori Y, Tobe K, Dina C, Yasuda K, Yamauchi T, Otabe S, Okada T, Eto K, Kadowaki H, Hagura R, Akanuma Y, Yazaki Y, Nagai R, Taniyama M, Matsubara K, Yoda M, Nakano Y, Tomita M, Kimura S, Ito C, Froguel P, Kadowaki T: Genetic variation in the gene encoding adiponectin is associated with an increased risk of type 2 diabetes in the Japanese population. *Diabetes* 51:536–540, 2002
 35. Filippi E, Sentinelli F, Trischitta V, Romeo S, Arca M, Leonetti F, Di Mario U, Baroni MG: Association of the human adiponectin gene and insulin resistance. *Eur J Hum Genet* 12:199–205, 2004
 36. Fumeron F, Aubert R, Siddiq A, Betoulle D, Pean F, Hadjadj S, Tichet J, Wilpart E, Chesnier MC, Balkau B, Froguel P, Marre M: Adiponectin gene polymorphisms and adiponectin levels are independently associated with the development of hyperglycemia during a 3-year period: the epidemiologic data on the insulin resistance syndrome prospective study. *Diabetes* 53:1150–1157, 2004
 37. Gibson F, Froguel P: Genetics of the APM1 locus and its contribution to type 2 diabetes susceptibility in French Caucasians. *Diabetes* 53:2977–2983, 2004
 38. Gonzalez-Sanchez JL, Zabena CA, Martinez-Larrad MT, Fernandez-Perez C, Perez-Barba M, Laakso M, Serrano-Rios M: An SNP in the adiponectin gene is associated with decreased serum adiponectin levels and risk for impaired glucose tolerance. *Obes Res* 13:807–812, 2005
 39. Gu HF, Abulaiti A, Ostenson CG, Humphreys K, Wahlestedt C, Brookes AJ, Efendic S: Single nucleotide polymorphisms in the proximal promoter region of the adiponectin (APM1) gene are associated with type 2 diabetes in Swedish caucasians. *Diabetes* 53 (Suppl. 1):S31–S35, 2004
 40. Hu FB, Doria A, Li T, Meigs JB, Liu S, Memisoglu A, Hunter D, Manson JE: Genetic variation at the adiponectin locus and risk of type 2 diabetes in women. *Diabetes* 53:209–213, 2004
 41. Jang Y, Lee JH, Chae JS, Kim OY, Koh SJ, Kim JY, Cho H, Lee JE, Ordovas JM: Association of the 276GT polymorphism of the adiponectin gene with cardiovascular disease risk factors in nondiabetic Koreans. *Am J Clin Nutr* 82:760–767, 2005
 42. Jang Y, Lee JH, Kim OY, Koh SJ, Chae JS, Woo JH, Cho H, Lee JE, Ordovas JM: The SNP276G>T polymorphism in the adiponectin (ACDC) gene is more strongly associated with insulin resistance and cardiovascular disease risk than SNP45T>G in nonobese/nondiabetic Korean men independent of abdominal adiposity and circulating plasma adiponectin. *Metabolism* 55:59–66, 2006
 43. Lee YY, Lee NS, Cho YM, Moon MK, Jung HS, Park YJ, Park HJ, Youn BS, Lee HK, Park KS, Shin HD: Genetic association study of adiponectin polymorphisms with risk of type 2 diabetes mellitus in Korean population. *Diabet Med* 22:569–575, 2005
 44. Mackevics V, Heid IM, Wagner SA, Cip P, Doppelmayr H, Lejnieks A, Gohlke H, Ladurner G, Illig T, Iglseider B, Kronenberg F, Paulweber B: The adiponectin gene is associated with adiponectin levels but not with characteristics of the insulin resistance syndrome in healthy Caucasians. *Eur J Hum Genet* 14:349–356, 2006
 45. Menzaghi C, Ercolino T, Di Paola R, Berg AH, Warram JH, Scherer PE, Trischitta V, Doria A: A haplotype at the adiponectin locus is associated with obesity and other features of the insulin resistance syndrome. *Diabetes* 51:2306–2312, 2002
 46. Nakatani K, Noma K, Nishioka J, Kasai Y, Morioka K, Katsuki A, Hori Y, Yano Y, Sumida Y, Wada H, Nobori T: Adiponectin gene variation associates with the increasing risk of type 2 diabetes in non-diabetic Japanese subjects. *Int J Mol Med* 15:173–177, 2005

47. Populaire C, Mori Y, Dina C, Vasseur F, Vaxillaire M, Kadowaki T, Froguel P: Does the -11377 promoter variant of APM1 gene contribute to the genetic risk for Type 2 diabetes mellitus in Japanese families? *Diabetologia* 46:443–445, 2003
48. Rudofsky G Jr, Schlimme M, Schlotterer A, von Eynatten M, Reismann P, Tafel J, Grafe I, Morcos M, Nawroth P, Bierhaus A, Hamann A: No association of the 94T/G polymorphism in the adiponectin gene with diabetic complications. *Diabetes Obes Metab* 7:455–459, 2005
49. Salmenniemi U, Zacharova J, Ruotsalainen E, Vauhkonen I, Pihlajamaki J, Kainulainen S, Punnonen K, Laakso M: Association of adiponectin level and variants in the adiponectin gene with glucose metabolism, energy expenditure, and cytokines in offspring of type 2 diabetic patients. *J Clin Endocrinol Metab* 90:4216–4223, 2005
50. Schwarz PE, Towers GW, Fischer S, Govindarajulu S, Schulze J, Bornstein SR, Hanefeld M, Vasseur F: Hypoadiponectinemia is associated with progression toward type 2 diabetes and genetic variation in the ADIPOQ gene promoter. *Diabetes Care* 29:1645–1650, 2006
51. Stumvoll M, Tschrirter O, Fritsche A, Staiger H, Renn W, Weisser M, Machicao F, Haring H: Association of the T-G polymorphism in adiponectin (exon 2) with obesity and insulin sensitivity: interaction with family history of type 2 diabetes. *Diabetes* 51:37–41, 2002
52. Ukkola O, Ravussin E, Jacobson P, Sjöström L, Bouchard C: Mutations in the adiponectin gene in lean and obese subjects from the Swedish obese subjects cohort. *Metabolism* 52:881–884, 2003
53. Ukkola O, Santaniemi M, Rankinen T, Leon AS, Skinner JS, Wilmore JH, Rao DC, Bergman R, Kesaniemi YA, Bouchard C: Adiponectin polymorphisms, adiposity and insulin metabolism: HERITAGE family study and Oulu diabetic study. *Ann Med* 37:141–150, 2005
54. Vasseur F, Helbecque N, Dina C, Lobbens S, Delannoy V, Gaget S, Boutin P, Vaxillaire M, Lepretre F, Dupont S, Hara K, Clement K, Bihain B, Kadowaki T, Froguel P: Single-nucleotide polymorphism haplotypes in the both proximal promoter and exon 3 of the APM1 gene modulate adipocyte-secreted adiponectin hormone levels and contribute to the genetic risk for type 2 diabetes in French Caucasians. *Hum Mol Genet* 11:2607–2614, 2002
55. Vasseur F, Helbecque N, Lobbens S, Vasseur-Delannoy V, Dina C, Clement K, Boutin P, Kadowaki T, Scherer PE, Froguel P: Hypoadiponectinaemia and high risk of type 2 diabetes are associated with adiponectin-encoding (ACDC) gene promoter variants in morbid obesity: evidence for a role of ACDC in diabetes. *Diabetologia* 48:892–899, 2005
56. Vozarova de Courten B, Hanson RL, Funahashi T, Lindsay RS, Matsuzawa Y, Tanaka S, Thameem F, Gruber JD, Froguel P, Wolford JK: Common polymorphisms in the adiponectin gene ACDC are not associated with diabetes in Pima Indians. *Diabetes* 54:284–289, 2005
57. Yoshioka K, Yoshida T, Umekawa T, Kogure A, Takakura Y, Toda H, Yoshikawa T: Adiponectin gene polymorphism (G276T) is not associated with incipient diabetic nephropathy in Japanese type 2 diabetic patients. *Metabolism* 53:1223–1226, 2004
58. Chuang LM, Chiu YF, Sheu WH, Hung YJ, Ho LT, Grove J, Rodriguez B, Quertermous T, Chen YD, Hsiung CA, Tai TY: Biethnic comparisons of autosomal genomic scan for loci linked to plasma adiponectin in populations of Chinese and Japanese origin. *J Clin Endocrinol Metab* 89:5772–5778, 2004
59. Comuzzie AG, Funahashi T, Sonnenberg G, Martin LJ, Jacob HJ, Black AE, Maas D, Takahashi M, Kihara S, Tanaka S, Matsuzawa Y, Blangero J, Cohen D, Kissebah A: The genetic basis of plasma variation in adiponectin, a global endophenotype for obesity and the metabolic syndrome. *J Clin Endocrinol Metab* 86:4321–4325, 2001
60. Guo X, Saad MF, Langefeld CD, Williams AH, Cui J, Taylor KD, Norris JM, Jinagouda S, Darwin CH, Mitchell BD, Bergman RN, Sutton B, Chen YD, Wagenknecht LE, Bowden DW, Rotter JI: Genome-wide linkage of plasma adiponectin reveals a major locus on chromosome 3q distinct from the adiponectin structural gene: the IRAS Family Study. *Diabetes* 55:1723–1730, 2006
61. Lindsay RS, Funahashi T, Krakoff J, Matsuzawa Y, Tanaka S, Kobes S, Bennett PH, Tataranni PA, Knowler WC, Hanson RL: Genome-wide linkage analysis of serum adiponectin in the Pima Indian population. *Diabetes* 52:2419–2425, 2003
62. Menzaghi C, Ercolino T, Salvemini L, Coco A, Kim SH, Fini G, Doria A, Trischitta V: Multigenic control of serum adiponectin levels: evidence for a role of the APM1 gene and a locus on 14q13. *Physiol Genomics* 19:170–174, 2004
63. Pollin TI, Tanner K, O'Connell JR, Ott SH, Damcott CM, Shuldiner AR, McLenithan JC, Mitchell BD: Linkage of plasma adiponectin levels to 3q27 explained by association with variation in the APM1 gene. *Diabetes* 54:268–274, 2005
64. Iwashima Y, Katsuya T, Ishikawa K, Ouchi N, Ohishi M, Sugimoto K, Fu Y, Motone M, Yamamoto K, Matsuo A, Ohashi K, Kihara S, Funahashi T, Rakugi H, Matsuzawa Y, Ogihara T: Hypoadiponectinemia is an independent risk factor for hypertension. *Hypertension* 43:1318–1323, 2004
65. Mousavinasab F, Tahtinen T, Jokelainen J, Koskela P, Vanhala M, Oikarinen J, Keinanen-Kiukkaanniemi S, Laakso M: Common polymorphisms (single-nucleotide polymorphisms SNP+45 and SNP+276) of the adiponectin gene regulate serum adiponectin concentrations and blood pressure in young Finnish men. *Mol Genet Metab* 87:147–151, 2006
66. Poitou C, Lacorte JM, Coupaye M, Bertrais S, Bedel JF, Lafon N, Bouillot JL, Galan P, Borson-Chazot F, Basdevant A, Coussieu C, Clement K: Relationship between single nucleotide polymorphisms in leptin, IL6 and adiponectin genes and their circulating product in morbidly obese subjects before and after gastric banding surgery. *Obes Surg* 15:11–23, 2005
67. van Houwelingen HC, Arends LR, Stijnen T: Advanced methods in meta-analysis: multivariate approach and meta-regression. *Stat Med* 21:589–624, 2002
68. Macaskill P, Walter SD, Irwig L: A comparison of methods to detect publication bias in meta-analysis. *Stat Med* 20:641–654, 2001
69. Bouatia-Naji N, Meyre D, Lobbens S, Seron K, Fumeron F, Balkau B, Heude B, Jouret B, Scherer PE, Dina C, Weill J, Froguel P: ACDC/Adiponectin polymorphisms are associated with severe childhood and adult obesity. *Diabetes* 55:545–550, 2006
70. Conne B, Stutz A, Vassalli JD: The 3' untranslated region of messenger RNA: a molecular 'hotspot' for pathology? *Nat Med* 6:637–641, 2000
71. Ohashi K, Ouchi N, Kihara S, Funahashi T, Nakamura T, Sumitsuji S, Kawamoto T, Matsumoto S, Nagaretani H, Kumada M, Okamoto Y, Nishizawa H, Kishida K, Maeda N, Hiraoka H, Iwashima Y, Ishikawa K, Ohishi M, Katsuya T, Rakugi H, Ogihara T, Matsuzawa Y: Adiponectin I164T mutation is associated with the metabolic syndrome and coronary artery disease. *J Am Coll Cardiol* 43:1195–1200, 2004
72. Waki H, Yamauchi T, Kamon J, Ito Y, Uchida S, Kita S, Hara K, Hada Y, Vasseur F, Froguel P, Kimura S, Nagai R, Kadowaki T: Impaired multimerization of human adiponectin mutants associated with diabetes: molecular structure and multimer formation of adiponectin. *J Biol Chem* 278:40352–40363, 2003
73. Menzaghi C, Ercolino T, Salvemini L, Coco A, Fini G, Di Paola R, Doria A, Trischitta V: Lack of evidence for interaction between APM1 and PPAR-gamma2 genes in modulating insulin sensitivity in nondiabetic Caucasians from Italy. *J Intern Med* 257:315–317, 2005
74. Mousavinasab F, Tahtinen T, Jokelainen J, Koskela P, Vanhala M, Oikarinen J, Keinanen-Kiukkaanniemi S, Laakso M: Effect of the Pro12Ala polymorphism of the PPAR-gamma2 gene on serum adiponectin changes. *Endocrine* 27:307–309, 2005
75. Yamamoto Y, Hirose H, Miyashita K, Nishikai K, Saito I, Taniyama M, Tomita M, Saruta T: PPAR(gamma)2 gene Pro12Ala polymorphism may influence serum level of an adipocyte-derived protein, adiponectin, in the Japanese population. *Metabolism* 51:1407–1409, 2002
76. Gonzalez-Sanchez JL, Martinez-Calatrava MJ, Martinez-Larrad MT, Zabenca C, Fernandez-Perez C, Laakso M, Serrano-Rios M: Interaction of the -308G/A promoter polymorphism of the tumor necrosis factor-alpha gene with single-nucleotide polymorphism 45 of the adiponectin gene: effect on serum adiponectin concentrations in a Spanish population. *Clin Chem* 52:97–103, 2006
77. Thamer C, Machicao F, Fritsche A, Stumvoll M, Haring H: No influence of the PPAR-gamma2 Pro12Ala genotype on serum adiponectin concentrations in healthy Europeans. *Metabolism* 52:798; author reply 798–799, 2003
78. Zacharova J, Chiasson JL, Laakso M: The common polymorphisms (single nucleotide polymorphism [SNP] +45 and SNP +276) of the adiponectin gene predict the conversion from impaired glucose tolerance to type 2 diabetes: the STOP-NIDDM Trial. *Diabetes* 54:893–899, 2005
79. Stenvinkel P, Marchlewski A, Pecoits-Filho R, Heimbürger O, Zhang Z, Hoff C, Holmes C, Axelsson J, Arvidsson S, Schalling M, Barany P, Lindholm B, Nordfors L: Adiponectin in renal disease: relationship to phenotype and genetic variation in the gene encoding adiponectin. *Kidney Int* 65:274–281, 2004
80. Kang ES, Park SY, Kim HJ, Ahn CW, Nam M, Cha BS, Lim SK, Kim KR, Lee HC: The influence of adiponectin gene polymorphism on the rosiglitazone response in patients with type 2 diabetes. *Diabetes Care* 28:1139–1144, 2005
81. Tanko LB, Siddiq A, Lecoeur C, Larsen PJ, Christiansen C, Walley A, Froguel P: ACDC/adiponectin and PPAR-gamma gene polymorphisms: implications for features of obesity. *Obes Res* 13:2113–2121, 2005
82. Yang WS, Hsiung CA, Ho LT, Chen YT, He CT, Curb JD, Grove J, Quertermous T, Chen YD, Kuo SS, Chuang LM: Genetic epistasis of

- adiponectin and PPARgamma2 genotypes in modulation of insulin sensitivity: a family-based association study. *Diabetologia* 46:977–983, 2003
83. Okamoto Y, Kihara S, Ouchi N, Nishida M, Arita Y, Kumada M, Ohashi K, Sakai N, Shimomura I, Kobayashi H, Terasaka N, Inaba T, Funahashi T, Matsuzawa Y: Adiponectin reduces atherosclerosis in apolipoprotein E-deficient mice. *Circulation* 106:2767–2770, 2002
 84. Yamauchi T, Kamon J, Waki H, Imai Y, Shimozawa N, Hioki K, Uchida S, Ito Y, Takakuwa K, Matsui J, Takata M, Eto K, Terauchi Y, Komeda K, Tsunoda M, Murakami K, Ohnishi Y, Naitoh T, Yamamura K, Ueyama Y, Froguel P, Kimura S, Nagai R, Kadowaki T: Globular adiponectin protected ob/ob mice from diabetes and ApoE-deficient mice from atherosclerosis. *J Biol Chem* 278:2461–2468, 2003
 85. Sakuta H, Suzuki T, Yasuda H, Ito T: Adiponectin levels and cardiovascular risk factors in Japanese men with type 2 diabetes. *Endocr J* 52:241–244, 2005
 86. Imatoh T, Miyazaki M, Kadowaki K, Babazono A, Sato M, Une H: Interaction of low serum adiponectin levels and smoking on coronary stenosis in Japanese men. *Int J Cardiol* 110:251–255, 2006
 87. Lacquemant C, Froguel P, Lobbens S, Izzo P, Dina C, Ruiz J: The adiponectin gene SNP+45 is associated with coronary artery disease in type 2 (non-insulin-dependent) diabetes mellitus. *Diabet Med* 21:776–781, 2004
 88. Bacci S, Menzaghi C, Ercolino T, Ma X, Rauseo A, Salvemini L, Vigna C, Fanelli R, Di Mario U, Doria A, Trischitta V: The +276 G/T single nucleotide polymorphism of the adiponectin gene is associated with coronary artery disease in type 2 diabetic patients. *Diabetes Care* 27:2015–2020, 2004
 89. Qi L, Doria A, Manson JE, Meigs JB, Hunter D, Mantzoros CS, Hu FB: Adiponectin genetic variability, plasma adiponectin, and cardiovascular risk in patients with type 2 diabetes. *Diabetes* 55:1512–1516, 2006
 90. Qi L, Li T, Rimm E, Zhang C, Rifai N, Hunter D, Doria A, Hu FB: The +276 polymorphism of the APM1 gene, plasma adiponectin concentration, and cardiovascular risk in diabetic men. *Diabetes* 54:1607–1610, 2005
 91. Yki-Jarvinen H: Thiazolidinediones. *N Engl J Med* 351:1106–1118, 2004
 92. Kubota N, Terauchi Y, Kubota T, Kumagai H, Itoh S, Satoh H, Yano W, Ogata H, Tokuyama K, Takamoto I, Mineyama T, Ishikawa M, Moroi M, Sugi K, Yamauchi T, Ueki K, Tobe K, Noda T, Nagai R, Kadowaki T: Pioglitazone ameliorates insulin resistance and diabetes by both adiponectin-dependent and -independent pathways. *J Biol Chem* 281:8748–8755, 2006
 93. Phillips SA, Ciaraldi TP, Kong AP, Bandukwala R, Aroda V, Carter L, Baxi S, Mudaliar SR, Henry RR: Modulation of circulating and adipose tissue adiponectin levels by anti-diabetic therapy. *Diabetes* 52:667–674, 2003
 94. Yu JG, Javorschi S, Hevener AL, Kruszynska YT, Norman RA, Sinha M, Olefsky JM: The effect of thiazolidinediones on plasma adiponectin levels in normal, obese, and type 2 diabetic subjects. *Diabetes* 51:2968–2974, 2002
 95. Antonucci T, Whitcomb R, McLain R, Lockwood D, Norris RM: Impaired glucose tolerance is normalized by treatment with the thiazolidinedione troglitazone. *Diabetes Care* 20:188–193, 1997
 96. Aronoff S, Rosenblatt S, Braithwaite S, Egan JW, Mathisen AL, Schneider RL: Pioglitazone hydrochloride monotherapy improves glycemic control in the treatment of patients with type 2 diabetes: a 6-month randomized placebo-controlled dose-response study: the Pioglitazone 001 Study Group. *Diabetes Care* 23:1605–1611, 2000
 97. Buchanan TA, Xiang AH, Peters RK, Kjos SL, Marroquin A, Goico J, Ochoa C, Tan S, Berkowitz K, Hodis HN, Azen SP: Preservation of pancreatic beta-cell function and prevention of type 2 diabetes by pharmacological treatment of insulin resistance in high-risk Hispanic women. *Diabetes* 51:2796–2803, 2002
 98. Nolan JJ, Ludvik B, Beerdsen P, Joyce M, Olefsky J: Improvement in glucose tolerance and insulin resistance in obese subjects treated with troglitazone. *N Engl J Med* 331:1188–1193, 1994
 99. Wolford JK, Yeatts KA, Dhanjal SK, Black MH, Xiang AH, Buchanan TA, Watanabe RM: Sequence variation in PPARG may underlie differential response to troglitazone. *Diabetes* 54:3319–3325, 2005
 100. Yamauchi T, Kamon J, Ito Y, Tsuchida A, Yokomizo T, Kita S, Sugiyama T, Miyagishi M, Hara K, Tsunoda M, Murakami K, Ohteki T, Uchida S, Takekawa S, Waki H, Tsuno NH, Shibata Y, Terauchi Y, Froguel P, Tobe K, Koyasu S, Taira K, Kitamura T, Shimizu T, Nagai R, Kadowaki T: Cloning of adiponectin receptors that mediate antidiabetic metabolic effects. *Nature* 423:762–769, 2003
 101. Hug C, Wang J, Ahmad NS, Bogan JS, Tsao TS, Lodish HF: T-cadherin is a receptor for hexameric and high-molecular-weight forms of Acrp30/adiponectin. *Proc Natl Acad Sci U S A* 101:10308–10313, 2004
 102. Civitarese AE, Jenkinson CP, Richardson D, Bajaj M, Cusi K, Kashyap S, Berria R, Belfort R, DeFronzo RA, Mandarino LJ, Ravussin E: Adiponectin receptors gene expression and insulin sensitivity in non-diabetic Mexican Americans with or without a family history of type 2 diabetes. *Diabetologia* 47:816–820, 2004
 103. Chinetti G, Zawadzki C, Fruchart JC, Staels B: Expression of adiponectin receptors in human macrophages and regulation by agonists of the nuclear receptors PPARalpha, PPARgamma, and LXR. *Biochem Biophys Res Commun* 314:151–158, 2004
 104. Wang H, Zhang H, Jia Y, Zhang Z, Craig R, Wang X, Elbein SC: Adiponectin receptor 1 gene (ADIPOR1) as a candidate for type 2 diabetes and insulin resistance. *Diabetes* 53:2132–2136, 2004
 105. Hara K, Horikoshi M, Kitazato H, Yamauchi T, Ito C, Noda M, Ohashi J, Froguel P, Tokunaga K, Nagai R, Kadowaki T: Absence of an association between the polymorphisms in the genes encoding adiponectin receptors and type 2 diabetes. *Diabetologia* 48:1307–1314, 2005
 106. Vaxillaire M, Dechaume A, Vasseur-Delannoy V, Lahmidi S, Vatin V, Lepretre F, Boutin P, Herberg S, Charpentier G, Dina C, Froguel P: Genetic analysis of ADIPOR1 and ADIPOR2 candidate polymorphisms for type 2 diabetes in the Caucasian population. *Diabetes* 55:856–861, 2006
 107. Damcott CM, Ott SH, Pollin TI, Reinhart LJ, Wang J, O'Connell JR, Mitchell BD, Shuldiner AR: Genetic variation in adiponectin receptor 1 and adiponectin receptor 2 is associated with type 2 diabetes in the Old Order Amish. *Diabetes* 54:2245–2250, 2005
 108. Stefan N, Machicao F, Staiger H, Machann J, Schick F, Tschritter O, Spieth C, Weigert C, Fritsche A, Stumvoll M, Haring HU: Polymorphisms in the gene encoding adiponectin receptor 1 are associated with insulin resistance and high liver fat. *Diabetologia* 48:2282–2291, 2005
 109. Siitonen N, Pulkkinen L, Mager U, Lindstrom J, Eriksson JG, Valle TT, Hamalainen H, Ilanne-Parikka P, Keinanen-Kiukkaanniemi S, Tuomilehto J, Laakso M, Uusitupa M: Association of sequence variations in the gene encoding adiponectin receptor 1 (ADIPOR1) with body size and insulin levels: the Finnish Diabetes Prevention Study. *Diabetologia* 49:1795–1805, 2006
 110. Richardson DK, Schneider J, Fourcaudot MJ, Rodriguez LM, Arya R, Dyer TD, Almasy L, Blangero J, Stern MP, Defronzo RA, Duggirala R, Jenkinson CP: Association between variants in the genes for adiponectin and its receptors with insulin resistance syndrome (IRS)-related phenotypes in Mexican Americans. *Diabetologia* 49:2317–2328, 2006
 111. Broedl UC, Lehrke M, Fleischer-Briellmaier E, Tietz AB, Nagel JM, Goke B, Lohse P, Parhofer KG: Genetic variants of adiponectin receptor 2 are associated with increased adiponectin levels and decreased triglyceride/VLDL levels in patients with metabolic syndrome. *Cardiovasc Diabetol* 5:11, 2006
 112. Soccio T, Zhang YY, Bacci S, Mlynarski W, Placha G, Raggio G, Di Paola R, Marucci A, Johnstone MT, Gervino EV, Abumrad NA, Klein S, Trischitta V, Doria A: Common haplotypes at the adiponectin receptor 1 (ADIPOR1) locus are associated with increased risk of coronary artery disease in type 2 diabetes. *Diabetes* 55:2763–2770, 2006