

Increasing Dietary Leucine Intake Reduces Diet-Induced Obesity and Improves Glucose and Cholesterol Metabolism in Mice via Multiple Mechanisms

Yiying Zhang,^{1,2} Kaiying Guo,¹ Robert E. LeBlanc,¹ Daniella Loh,¹ Gary J. Schwartz,³ and Yi-Hao Yu⁴

Leucine, as an essential amino acid and activator of mTOR (mammalian target of rapamycin), promotes protein synthesis and suppresses protein catabolism. However, the effect of leucine on overall glucose and energy metabolism remains unclear, and whether leucine has beneficial effects as a long-term dietary supplement has not been examined. In the present study, we doubled dietary leucine intake via leucine-containing drinking water in mice with free excess to either a rodent chow or a high-fat diet (HFD). While it produced no major metabolic effects in chow-fed mice, increasing leucine intake resulted in up to 32% reduction of weight gain ($P < 0.05$) and a 25% decrease in adiposity ($P < 0.01$) in HFD-fed mice. The reduction of adiposity resulted from increased resting energy expenditure associated with increased expression of uncoupling protein 3 in brown and white adipose tissues and in skeletal muscle, while food intake was not decreased. Increasing leucine intake also prevented HFD-induced hyperglycemia, which was associated with improved insulin sensitivity, decreased plasma concentrations of glucagon and glucogenic amino acids, and downregulation of hepatic glucose-6-phosphatase. Additionally, plasma levels of total and LDL cholesterol were decreased by 27% ($P < 0.001$) and 53% ($P < 0.001$), respectively, in leucine supplemented HFD-fed mice compared with the control mice fed the same diet. The reduction in cholesterol levels was largely independent of leucine-induced changes in adiposity. In conclusion, increases in dietary leucine intake substantially decrease diet-induced obesity, hyperglycemia, and hypercholesterolemia in mice with ad libitum consumption of HFD likely via multiple mechanisms. *Diabetes* 56:1647–1654, 2007

From the ¹Department of Pediatrics, Division of Molecular Genetics, Columbia University, New York, New York; the ²Naomi Berrie Diabetes Center, Columbia University, New York, New York; the ³Department of Medicine and Neuroscience, Diabetes Research and Training Center, Albert Einstein College of Medicine, Bronx, New York; and the ⁴Department of Medicine, Division of Preventive Medicine, Columbia University, New York, New York.

Address correspondence and reprint requests to Yiying Zhang, PhD, Division of Molecular Genetics, Department of Pediatrics, Columbia University, Russ Berrie Pavilion, Rm. 620, 1150 St. Nicholas Ave., New York, NY 10032. E-mail: yz84@columbia.edu. Or Yi-Hao Yu, MD, PhD, Division of Preventive Medicine, Department of Medicine, Columbia University, 630 W. 168th St., PH 10-305J, New York, NY 10032. E-mail: yy102@columbia.edu.

Received for publication 27 January 2007 and accepted in revised form 19 February 2007.

Published ahead of print at <http://diabetes.diabetesjournals.org> on 14 March 2007. DOI: 10.2337/db07-0123.

BAT, brown adipose tissue; G6P, glucose-6-phosphatase; HFD, high-fat diet; HOMA-IR, homeostasis model assessment of insulin resistance; mTOR, mammalian target of rapamycin; UCP, uncoupling protein; WAT, white adipose tissue.

© 2007 by the American Diabetes Association.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Despite the increased awareness of health risks associated with obesity and its attendant metabolic disorders, the prevalence of obesity has doubled in the past 20 years (1). Many dieting strategies have been developed and practiced in the past two decades, with variable short-term success rates (2,3). Long-term success in maintaining weight loss has proven to be difficult, in part because energy metabolism becomes more efficient at the reduced weight as a result of the normal physiological responses to weight loss and energy deficits (4,5).

Manipulation of dietary composition of macronutrients merits careful investigation because macronutrients not only provide calories but some components may also function as signaling molecules to affect feeding behavior, energy balance, and fuel efficiency. Dietary supplementation is particularly attractive in that it may improve diet qualities and reduce cardiovascular risks associated with obesity and the metabolic syndrome without affecting the gross macronutrient compositions and the palatability of regular food, which will likely result in better long-term compliance.

Leucine is an essential branched chain amino acid that cannot be produced and can only be obtained from diet in human. Leucine serves not only as a building block for protein synthesis but is also a potent activator of the mammalian target of rapamycin (mTOR), a serine/threonine kinase involved in many cellular processes, including protein synthesis, cell growth, and metabolism (6–10). Indeed, leucine supplementation increases the net protein anabolism in various tissues (11–16). However, by activating mTOR and S6 kinase, leucine also feedback inhibits insulin signaling and decrease glucose utilization in skeletal muscle (17–20).

Several recent studies have shown that protein-rich diets produce a better glycemic control and a greater loss of body fat and less of lean tissue during weight loss than do carbohydrate-rich diets with the same fat and caloric content (21–24). Branched chain amino acids, especially leucine, have been speculated to play a key role in this process (25). Supporting this hypothesis, Donato et al. (26) reported that leucine supplementation during caloric restriction also results in more fat loss and improves protein synthesis in liver and muscle. Recently, Seeley and colleagues (10) showed that infusion of leucine into the third ventricle of the rat brain decreases the animal's food intake and body weight via activation of the mTOR pathway in the arcuate nucleus of the hypothalamus. Thus, as

leucine may function to promote fat loss while also, independently, increase insulin resistance in peripheral tissues (see above), it remains unclear whether increased leucine intake in animals affects the development of dietary obesity as well as whether it has an overall detrimental or beneficial health effect.

We sought to determine the net effects of dietary supplementation of leucine on whole-body macronutrient and energy metabolism in mice with free access to a high-fat diet (HFD) and to assess how cardiovascular risks associated with diet-induced obesity may be modified by increased dietary leucine intake. Here, we show that doubling dietary leucine intake significantly reduces HFD-induced weight gain and improves hyperglycemia and hypercholesterolemia, which is associated with a previously unrecognized effect of leucine, i.e., increased resting energy expenditure. Furthermore, modifications of the various cardiometabolic risk factors associated with increased leucine intake appear to involve multiple mechanisms that are not solely dependent on reduced adiposity.

RESEARCH DESIGN AND METHODS

Animal husbandry, diets, and leucine supplementation. Male C57BL/6J mice were purchased from The Jackson Laboratories. Animal protocols were in compliance with the accepted standards of animal care and were approved by the Columbia University institutional animal care and use committee. Mice were maintained at 22°C on a 12:12 light-dark cycle (0700–1900 h) and had ad libitum access to a regular rodent chow (Purina Rodent Diet 20 5053) or, when indicated, a HFD containing 60% fat calories (D12492; Research Diets, New Brunswick, NJ). Both diets contained 20% protein calories. Leucine was supplemented via drinking water made of 1.5% (wt/vol) L-leucine (Sigma, St. Louis, MO).

Blood glucose, glucose tolerance tests, and insulin tolerance tests. Blood glucose levels were measured in tail-vein blood using a Glucometer Elite (Bayer, Elkhart, IN). For glucose tolerance tests, blood glucose was measured at 0, 30, 60, and 90 min after a bolus intraperitoneal glucose administration (1 mg/g body wt) to overnight-fasted mice. For insulin tolerance tests, regular human insulin was administered intraperitoneally (0.75 mU/g) to mice after 4-h food and leucine deprivation, and blood glucose was measured at 0, 30, 60, and 90 min after insulin injection. The homeostasis model assessment of insulin resistance (HOMA-IR) index was calculated based on the conventional formula: $HOMA-IR = \text{basal glucose (mmol/l)} \times \text{basal insulin (mU/l)} / 22.5$.

Indirect calorimetry. Five-month-old male C57BL/6J mice that had been on a HFD with or without leucine supplementation for 10 weeks were implanted with E-mitters (Mini-Mitter) for temperature measurements under anesthesia and were allowed a 4-day recovery before being housed in metabolic chambers. Oxygen consumption, locomotor activity, core temperature, and respiration exchange ratio were obtained continuously during 12:12 light-dark cycles using a comprehensive laboratory animal monitoring system (Columbus Instruments, Columbus, OH) open-circuit indirect calorimetry system. The same diet and leucine regimens for the mice were continued during indirect calorimetry. Data were collected over at least 5 days following 2–4 days of adaptation to the metabolic cages.

Determination of mRNA expression. Quantitative real-time RT-PCR was used to determine mRNA expression as previously described (27,28). Briefly, total RNA was isolated using miniRNaseasy columns (Qiagen, Valencia, CA) and reverse transcribed into single-stranded cDNA using random hexamers and M-MLV Reverse Transcriptase (Invitrogen, Carlsbad, CA). Quantitative amplification by PCR was carried out using Bio-Rad iQ SYBR Green Supermix (Bio-Rad Laboratories, Hercules, CA). Cyclophilin A was used as a control. Cycle numbers at a set fluorescence threshold within the exponential amplification range (C_t) were used to calculate expression levels of the genes of interest (relative to cyclophilin A). Changes in gene expression were expressed as fold increase or decrease over a specified reference sample. The following primers were used in this study: uncoupling protein (UCP)1, 5'-ggtgagttcga-caactccg-3' (forward) and 5'-ggtgatggtccctaa-gacacc-3' (reverse); UCP2, 5'-gtgtgctggagatacagagc-3' (forward) and 5'-gaggttgcttccag-gagag-3' (reverse); UCP3, 5'-caaggagcggaccactcc-3' (forward) and 5'-ctctctccctcagttccatg-3' (reverse); and cyclophilin A, 5'-atggcactggcggcaggtcc-3' (forward) and 5'-ttgccattctctggaccacaaa-3' (reverse).

Protein extraction and immunoblotting. Gastrocnemius muscle (30 mg) or adipose tissues (100 mg) were powdered in liquid nitrogen and homogenized in a homogenization buffer (50 mmol/l Tris-HCl, pH 8.0, 1% NP-40, 1 mmol/l

TABLE 1

Effects of leucine supplementation on plasma hormone levels in HFD-fed mice

	HFD-W	HFD-Leu
Insulin (ng/ml)	5.28 ± 0.78	3.26 ± 0.56*
Glucagon (pg/ml)	52.3 ± 9.9	23.9 ± 1.3*
Leptin (ng/ml)	36.3 ± 3.3	18.2 ± 5.4†
Adiponectin (mg/ml)	11.1 ± 1.4	10.3 ± 0.5
Corticosterone (ng/ml)	149 ± 39	112 ± 34
Epinephrine (ng/ml)	0.89 ± 0.19	1.11 ± 0.20
Norepinephrine (ng/ml)	0.89 ± 0.22	0.86 ± 0.16

Data are means ± SE. * $P < 0.05$ and † $P < 0.01$, HFD-W vs. HFD-Leu; $n = 5$.

EDTA, 150 mmol/l NaCl₂, 50 mmol/l NaF, 0.5 mmol/l NaVO₄, 5 mmol/l NaP₂O₇, and 1 mmol/l DTT and protease inhibitors). Tissue homogenates were sonicated for 2 min in an ice-water bath, followed by centrifugation at 14,000 rpm for 15 min in an Eppendorf Microcentrifuge to remove insoluble cellular debris. Proteins (50–100 μg) were resolved by SDS-PAGE and then transferred to nitrocellulose membranes. Levels of specific proteins were determined by immunoblotting using enhanced chemiluminescence detection horseradish peroxidase reagents (Amersham Biosciences, Piscataway, NJ) and antibodies to mouse UCP1 and UCP3 (Sigma-Aldrich, St. Louis, MO) and β-actin (Santa Cruz Biotechnology, Santa Cruz, CA). Immunoblotting signals were quantified using the Quantity One program (Bio-Rad).

Determination of body composition and plasma parameters. Body composition was determined using a Minispec TD-NMR Spectrometer (Bruker Optics). Unless otherwise indicated, plasma lipids, amino acids, and hormones, including insulin, leptin, glucagon, adiponectin, corticosterone, and catecholamines, were measured at the basal state (after a 5-h food and leucine deprivation) in HFD leucine and HFD control mice. Assays were done by the Mouse Phenotyping Center, Vanderbilt University. Plasma hormone concentrations are shown in Table 1.

Statistical analysis. The effects of leucine supplementation on weight, body composition, food intake, and blood glucose levels in HFD-fed and chow-fed mice were assessed separately using one-way ANOVA (Statistica version 6; StatSoft, Tulsa, OK); Newman-Keuls tests were used for post hoc comparisons. Multiple regression analysis was used to assess direct effects of leucine versus leucine-mediated reduction in adiposity on glucose and cholesterol metabolism. A two-tailed $P < 0.05$ was considered statistically significant. For indirect calorimetry measurements, repeated-measures ANOVA (Prism; GraphPad Software, San Diego, CA) were used. Post hoc analyses were conducted using Tukey post hoc or Bonferroni comparisons. Data are expressed as means with SEs.

RESULTS

Leucine supplementation decreases diet-induced obesity by increasing resting energy expenditure associated with increased UCP3 expression in thermogenic tissues. Eight week-old male C57BL/6J mice on a regular rodent chow diet (chow) or on a HFD were supplemented with leucine via drinking water (chow-Leu and HFD-Leu, respectively). HFD and leucine supplementation were started simultaneously. Control mice on the respective diets were provided with regular tap water (chow-W or HFD-W). The amounts of regular or leucine-supplemented water consumed by the mice were the same. Leucine-supplemented mice ingested an additional ~55 mg leucine daily via water consumption, which nearly doubled the total daily leucine intake from food (~56 mg from chow; ~63 mg from HFD) (Fig. 1A). Leucine supplementation increased plasma leucine concentrations by ~30% during feeding but decreased overall plasma amino acid levels, particularly Asp, Glu, Gln, and Gly, in postabsorptive states and Asn, Pro, and Lys during feeding (Fig. 1C and D), suggesting a decreased protein catabolism in leucine-supplemented mice.

Mice on the HFD gained significantly more weight, mainly as fat, than did those on the chow diet as was




FIG. 1. Leucine supplementation decreases protein catabolism and reduces HFD-induced weight gain and adiposity. *A* and *B*: Daily leucine intake and caloric intake, respectively, calculated from the amount of ingested food and drinking water measured in individually caged mice over 14 days. Chow-Leu and HFD-Leu designate leucine-supplemented mice on regular chow and HFD, respectively; Chow-W and HFD-W designate the control mice supplied with regular water, on chow and HFD, respectively. *C* and *D*: Plasma concentrations of amino acids during feeding (fed state; measured at the fourth hour of the dark cycle) and at the postabsorptive state (measured at the seventh hour of the light cycle after a 5-h food and leucine deprivation), respectively. Values were determined from mice at the 14th week of indicated diet/supplement treatments. *E*: Body weights and growth curves during 14 weeks of indicated diet and supplementation treatments. The indicated diet/supplement regimens were started in 8-week-old weight-matched male mice, designated as week 0. *F*: Body composition at the end of 14 weeks of diet/supplement treatments. Lean body mass (Lean), fat mass (Fat), and total body mass (Total) were measured by nuclear magnetic resonance spectrometry, and the percentage of fat mass (%Fat) was calculated as fat mass/(lean + fat mass). Data are expressed as mean \pm SE, $n = 5$ in each group. * $P < 0.05$ and ** $P = 0.01$, HFD-Leu vs. HFD-W; # $P < 0.05$, Chow-Leu vs. Chow-W. Results shown in this figure are from a single representative experiment (from a total of three experiments, $n = 5$ for each treatment group in each experiment).




FIG. 2. Leucine supplementation increases resting energy expenditure. Four pairs of HFD-Leu and HFD-W mice were studied by indirect calorimetry after 10 weeks of the indicated diet/supplement regimens. **A:** Twenty-four hour oxygen consumption. **B:** Locomotor activity. ctr, control; leu, leucine. **C:** Core body temperature. **D:** Respiratory exchange ratio (RER). Data shown are average values measured over a 5- to 8-day period after a 4-day adaptation to the metabolic chamber.

expected (Fig. 1E and F). However, this diet-induced weight gain and expansion of fat mass were significantly attenuated in leucine-supplemented HFD-fed mice (Fig. 1E and F). Eight weeks into the diet/supplement regimens, HFD-W mice weighed 41.4 ± 1.2 g, whereas HFD-Leu mice weighed 37.6 ± 1.0 g. This represents a 32% ($P < 0.05$) reduction of HFD-induced weight gain (12.1 g in HFD-W vs. 8.3 g in HFD-Leu, $P < 0.05$) (Fig. 1E) and a 25% decrease in adiposity (41.3% fat in HFD-W vs. 31.0% fat in HFD-Leu, $P < 0.01$). Differences in body weight, fat mass, and adiposity persisted throughout the 14-week study (Fig. 1E and F). Leucine supplementation had no effect on body weight or composition in mice on the chow diet (Fig. 1E and F). Total caloric intake was increased by 8.4% in chow-Leu mice, compared with chow-W mice, but was not significantly different between HFD-Leu and HFD-W mice (Fig. 1B).

Since weight loss without decreased food intake (the HFD group) and increased food intake without weight gain (the chow group) both suggested a leucine-mediated increase in energy expenditure, we directly measured energy metabolism in four pairs of HFD-W and HFD-Leu

mice by indirect calorimetry. Total energy expenditure (24-h VO_2) normalized to body weight was markedly increased ($\sim 15\%$, $P < 0.01$) in HFD-Leu mice relative to HFD-W mice (Fig. 2A), despite a decrease in locomotor activity in HFD-Leu mice (Fig. 2B). Core body temperature was not significantly different between HFD-Leu and HFD-W mice (Fig. 2C). The respiratory exchange ratio (V_{CO_2}/VO_2) was slightly lower in HFD-Leu mice than in HFD-W mice (an average 0.808 vs. 0.814, respectively, $P < 0.01$) (Fig. 2D), suggesting increased fat oxidation in HFD-W mice.

To gain insight into molecular mechanisms for increased resting energy expenditure in HFD-Leu mice, we examined mRNA and/or protein levels of UCP1, -2, and -3, three major uncoupling proteins, in skeletal muscle and in brown adipose tissue (BAT) and white adipose tissue (WAT). Messenger RNA levels for UCP3, a major uncoupling protein in skeletal muscle and BAT, were increased by 1.51-fold ($P < 0.05$) in gastrocnemius muscle, 2.08-fold ($P < 0.001$) in perigonadal adipose tissue (WAT), and 1.67-fold ($P < 0.05$) in BAT of HFD-Leu mice, relative to the respective tissues of HFD-W mice (Fig. 3A). At the




FIG. 3. Leucine supplementation increases UCP3 expression in skeletal muscle and adipose tissues. **A:** Messenger RNA levels of UCP2 and -3 in gastrocnemius muscle, perigonadal adipose tissue (WAT), and intrascapular BAT and UCP1 in BAT of HFD-Leu (L) and HFD-W (W) mice described in Fig. 1 ($n = 5$). Data are shown as means \pm SE. * $P < 0.05$ and *** $P = 0.001$, HFD-Leu vs. HFD-W. **B:** Representative immunoblotting results for UCP3 protein expression in the same tissues as indicated. UCP1 protein levels were also shown in BAT.

protein level, UCP3 was increased by 22.7% ($P < 0.05$), 28.9% ($P < 0.01$), and 20.1% ($P < 0.05$) in gastrocnemius muscle, perigonadal adipose tissue, and BAT, respectively, in HFD-Leu mice compared with control tissues (Fig. 3B). However, the levels of the ubiquitously expressed UCP2 and BAT UCP1 were not significantly different (Fig. 3A and B). These results suggest that leucine may specifically regulate UCP3 gene expression in skeletal muscle, BAT, and WAT and that increased UCP3 expression may account, at least in part, for the increased resting energy expenditure in HFD-Leu mice.

Leucine supplementation improves glucose metabolism, reduces diet-induced insulin resistance, and lowers plasma glucagon levels and hepatic glucose-6-phosphatase expression. Four weeks after HFD treatment, fasting blood glucose levels became elevated in HFD-W mice (112.5 ± 7.0 mg/dl) compared with chow-W mice (74.2 ± 3.6 mg/dl, $P < 0.001$) (Fig. 4A), but leucine supplementation largely prevented this HFD-induced hyperglycemia (89.2 ± 7.6 vs. 112.5 ± 7.0 mg/dl, $P < 0.05$). The glucose-lowering effect of leucine persisted throughout the study period, and HFD-Leu mice remained euglycemic at week 10 (Fig. 4A). Basal plasma insulin concentrations were significantly decreased in HFD-Leu mice compared with HFD-W mice (Table 1 and Fig. 4E). Glucose tolerance tests and insulin tolerance tests performed at weeks 10 and 11, respectively, confirmed that HFD-Leu mice were significantly more glucose tolerant and insulin sensitive than HFD-W mice (Fig. 4B and C). The HOMA-IR index was $>50\%$ lower in HFD-Leu mice than in HFD-W mice (34.2 ± 7.6 vs. 67.7 ± 11.7 , $P < 0.05$) (Fig. 4D). A forward stepwise regression analysis shows that leucine treatment ($F = 5.6$, $P < 0.05$) is a better predicting factor than adiposity ($F = 3.5$, $P = 0.09$) for HOMA-IR, suggesting that the glucose-lowering effect of

leucine may not be entirely dependent on the decrease in adiposity.

Basal plasma glucagon levels were decreased by $\sim 50\%$ ($P < 0.05$) in HFD-Leu mice (23.9 ± 1.3 pg/ml) compared with HFD-W mice (52.3 ± 9.9 pg/ml) (Fig. 4E and Table 1). Messenger RNA levels for glucose-6-phosphatase (G6P), a key enzyme in regulating hepatic glucose production, were also significantly lower in the livers of HFD-Leu mice, although mRNA levels for PEPCK and PGC-1 α (peroxisome proliferator-activated receptor- γ coactivator), another two genes involved in hepatic gluconeogenesis (29,30), were not significantly different (Fig. 4F). G6P mRNA levels were significantly correlated with plasma glucagon levels ($r = 0.85$, $P < 0.01$) but not with insulin levels in HFD-fed mice (data not shown). Unlike HFD-fed mice, chow-fed mice remained euglycemic over the 14-week study period, and leucine supplementation had no additional effect on these values (Fig. 4A).

Leucine supplementation reduces diet-induced hypercholesterolemia independent of adiposity. Leucine supplementation for 14 weeks decreased total cholesterol levels by 27% ($P < 0.001$) and LDL-cholesterol by 53% ($P < 0.001$) in HFD-fed mice (Fig. 5A). The cholesterol-reducing effect of leucine remained significant when adiposity was also included as an independent variable in a multiple regression analysis (Fig. 5B). This finding contrasts with the effect of leucine on plasma leptin levels. Leucine supplementation lowered plasma leptin levels (18.2 ± 5.4 ng/ml in HFD-Leu mice vs. 36.3 ± 3.3 ng/ml in HFD-W mice, $P < 0.01$) (Table 1), but this effect was essentially lost when the decrease of adiposity was taken into account (Fig. 5B). No significant differences were detected in HDL cholesterol, total triglyceride, or free fatty acid levels (Fig. 5A). Taken together, these data suggest that leucine has a selective





FIG. 4. Leucine supplementation improves glucose metabolism by increasing insulin sensitivity and decreasing glucagon secretion in HFD-fed mice. Study design is as in Fig. 1E. **A:** Fasting (overnight food and leucine deprivation, regular water allowed) blood glucose concentrations were measured at weeks 4 and 10, and basal (after a 5-h food and leucine deprivation) blood glucose concentrations were measured at week 10. Diets are as indicated. ctrl, control mice supplied with regular water; Leu, leucine supplementation. The data are representative of three similar experiments. **B and C:** Intraperitoneal glucose tolerance tests (ITT) (1 mg glucose per g body wt) and insulin tolerance tests (0.75 mU insulin per g body wt), respectively, performed at weeks 10 and 11 of the diet/supplement treatments. **D:** HOMA-IR index calculated from basal plasma glucose and insulin levels at the end of a 14-week study period. **E:** Basal plasma glucose, insulin, and glucagon levels at the end of 14 weeks of diet/supplement study. **F:** Relative hepatic mRNA levels of G6P, peroxisome proliferator-activated receptor- γ coactivator (PGC)-1 α , and PEPCK in the same mice as in E. All data are shown as means \pm SE, $n = 5$ in each group. * $P < 0.05$ and ** $P < 0.01$, HFD-Leu vs. HFD-W.



B

Dependent variables	Independent variables			
	Leu treatment		Adiposity	
	β	p	β	p
Total C	-0.69	<0.05	0.23	0.46
LDL-C	-0.63	<0.05	0.34	0.21
leptin	0.12	0.665	1.01	<0.01

FIG. 5. Leucine supplementation decreases plasma levels of total and LDL cholesterol in HFD-fed mice. Study design is as in Fig. 1E. **A:** Fasting plasma levels of total cholesterol (TC), HDL cholesterol (HDL-C), LDL cholesterol (LDL-C), triglyceride (TG), and free fatty acids (FFA), measured at week 14 of the diet/supplement treatments as described. Data are shown as means \pm SE, $n = 5$ in each group. *** $P < 0.001$. **B:** Multiple regression analyses using β , standardized correlation coefficient, and P values to describe the outcomes, as shown. Leucine treatment and adiposity are independent variables; total cholesterol, LDL cholesterol, and leptin are dependent variables.

effect on LDL cholesterol reduction, independent of its effect on adiposity.

DISCUSSION

We have shown that doubling dietary leucine intake substantially reduced diet-induced weight gain and improved glucose and cholesterol metabolism in mice. The attenuation of HFD-induced weight gain by leucine supplementation resulted from increased resting energy expenditure associated with increased UCP3 protein expression in skeletal muscle, BAT, and WAT. Food intake was not reduced by the increased leucine intake. Both we and others have observed that mice on chronic HFD become highly fuel efficient; they gain more weight per unit of calorie consumed compared with chow-fed mice (31–34). In HFD-Leu mice, increased UCP3-mediated mitochondrial uncoupling of oxidative phosphorylation in thermogenic tissues likely accounts for the increased resting energy expenditure with reduced fuel efficiency. Our result is consistent with the reports that increased UCP3 expression reduces HFD-induced weight gain in transgenic mice (35) and is associated with the obesity-resistant phenotype in rats (34). Additionally, increases in muscle UCP3 expression level are generally seen in obese rats and humans (36,37), and weight loss results in decreases in muscle UCP3 expression in humans (37,38). However, HFD-Leu mice, while having lower adiposity relative to HFD-W mice, express higher levels of UCP3 protein in both skeletal muscle and adipose tissues. Thus, the effect of increased leucine intake on tissue UCP3 expression appears to be independent of the attenuated adiposity in HFD-Leu mice. How leucine increases UCP3 expression is unclear, it may depend on leucine-mediated activation of the mTOR-S6K pathway, since insulin also both activates mTOR-S6K signaling pathway and increases UCP3 expression in skeletal muscle (39).

It is of interest to note that while increased UCP3 expression in skeletal muscle and in adipose tissues is consistent with the observed increased in resting energy expenditure, core body temperatures were not significantly different between HFD-Leu and HFD-W mice. Normal core body temperatures were associated with normal diurnal fluctuations in both HFD-Leu and HFD-W mice, suggesting that the thermoregulation mechanism was intact and that excess heat generated by increased energy expenditure was effectively dissipated in HFD-Leu mice. A similar result has also been observed in mice with genetic ablation of MCH (melanin-concentrating hormone), a hypothalamic neuropeptide involved in regulating food intake and energy expenditure; a 20% increase in the rate of oxygen consumption in MCH^{-/-} mice has no effect on core body temperature (40).

Cota et al. (10) showed that centrally administered leucine activates the mTOR pathway in the hypothalamic arcuate nucleus and suppresses food intake in rats. Orally administered leucine in our study, however, had no significant suppressive effect on food intake. Possible explanations for the apparent discrepancy include the doses and routes by which leucine was introduced into animals (intracerebroventricular infusion versus oral administration). Orally administered leucine may be substantially metabolized in the liver before it reaches other tissue targets. Plasma leucine concentrations were increased by 30% during feeding in HFD-Leu mice and were not significantly different in the postabsorptive state. Thus, it is possible that leucine concentrations in the hypothalamus of HFD-Leu mice did not reach a level comparable with that achieved by intracerebroventricular leucine infusion reported by Cota et al. However, this does not preclude the possible involvement of the central nervous system in mediating leucine's effects on energy expenditure, whether in animals receiving centrally administered leucine or in animals with increased oral intake of leucine, such as HFD-Leu mice. Cota et al. did not report the status of energy expenditure, but it is possible that changes in centrally mediated energy expenditure may require a lower central nervous system level of leucine than what is necessary for suppressing food intake. Another potential explanation for the apparent discrepancy between our study and that of Cota et al. is the species difference. Mice (ours) and rats (Cota et al.) may have species-specific difference in ways in which they respond to leucine.

Locomotor activity was decreased in HFD-Leu mice compared with HFD-W mice. The cause for this change is unclear. However, this change was not due to toxicity from leucine supplementation. Leucine is among the most tolerated amino acids as no adverse effects of increased leucine intake (usually three times the daily requirement) were reported in various animal and human studies (16,41). In our study, leucine-supplemented mice consumed normal amounts of food and water and had a normal growth curve compared with controls. In fact, on the rodent chow diet group, leucine-supplemented mice consumed more food than control mice. Thus, the decreased locomotor activity does not appear to be related to food-seeking activity either.

Increasing dietary leucine intake significantly improves glucose metabolism in HFD-fed mice. It appears that better glycemic control is primarily due to the improved insulin sensitivity that is associated with decreased adiposity in HFD-Leu mice. However, the involvement of other mechanisms cannot be ruled out. In fact, multiple

regression analyses suggest that leucine's effect on the HOMA-IR index was only partially accounted for by the reduced adiposity. Two important players in the regulation of hepatic glucose production, plasma glucagon concentrations and hepatic G6P expression levels, were both lower in HFD-Leu mice. Additionally, plasma concentrations of glycogenic amino acids, which provide substrates for hepatic gluconeogenesis in the postabsorptive state (42), were also lower in HFD-Leu mice. Taken together, these results suggest that hepatic glucose production may be decreased in HFD-Leu mice, which could contribute to the improved glycemic control.

Leucine has a biphasic effect on glucagon secretion in the perfused rat pancreas: stimulatory in low glucose states (3.3 mmol/l) but inhibitory in high glucose states (8.3 mmol/l) (43) and may suppress glucagon secretion indirectly by lowering concentrations of glucagon secretagogues such as glutamine, glycine, and alanine in HFD-Leu mice (44,45). Leucine may also affect glucose metabolism by stimulating insulin secretion (43–45). Although fasting insulin levels were lower in HFD-Leu mice, most likely secondary to the improved insulin sensitivity, it cannot be ruled out that leucine may transiently stimulate insulin secretion in HFD-Leu mice. A direct action of leucine in skeletal muscle and liver may involve mTOR-mediated feedback inhibition of insulin signaling to affect glucose disposal and glucose production, respectively (17–20). However, our data show that increasing dietary leucine intake resulted in a net improvement of whole-body insulin sensitivity and glycemic control, along with reduced adiposity in mice on a chronic HFD.

Leucine markedly reduced plasma total and LDL cholesterol levels, independent of weight reduction. Cholesterol-reducing effects of L-arginine and keto and amino acid supplementation in diabetic rats and in patients on low protein diets have also been reported by others (46,47). Further elucidation of leucine signaling, via mTOR and/or other potential mediators, will be required to uncover the relevant molecular mechanisms.

Finally, despite the remarkable effects of leucine observed in HFD-fed mice, increased leucine intake had no notable effects on chow-fed mice, except that leucine-supplemented mice had slightly increased food intake. Since the increased food intake in these mice was not associated with increases in body weight, adiposity, or blood glucose levels, we speculate that this behavioral change was only secondary to a leucine-induced increase in basal metabolic rates, as a homeostatic mechanism to prevent downward excursion from the "normal" body weight.

In summary, we have shown that doubling dietary leucine intake on a chronic basis produces a net health benefit that includes reduction of diet-induced weight gain, hyperglycemia, and hypercholesterolemia in mice on a HFD. The reduction in adiposity results from a leucine-induced increase in resting energy expenditure and is associated with increased UCP3 protein expression in skeletal muscle and adipose tissues. The overall beneficial effects on lipid and glucose metabolism produced by leucine supplementation appear to involve multiple mechanisms that need to be further defined.

ACKNOWLEDGMENTS

This study was supported by grants from National Institutes of Health, DK063034 (to Y.Z.), DK071866 (to Y.Y.),

and DK066618 (to G.J.S.) and grants P30DK026687 and P30DK063068.

We thank Dr. Rudy Leibel for many discussions during the study and Drs. Rudy Leibel, Henry Ginsberg, and Ira Goldberg for reviewing the manuscript.

REFERENCES

- Hedley AA, Ogden CL, Johnson CL, Carroll MD, Curtin LR, Flegal KM: Prevalence of overweight and obesity among US children, adolescents, and adults, 1999–2002. *JAMA* 291:2847–2850, 2004
- Katz DL: Competing dietary claims for weight loss: finding the forest through truculent trees. *Annu Rev Public Health* 26:61–88, 2005
- Freedman MR, King J, Kennedy E: Popular diets: a scientific review. *Obes Res* 9:1S–40S, 2001
- Leibel RL, Rosenbaum M, Hirsch J: Changes in energy expenditure resulting from altered body weight. *N Engl J Med* 332:621–628, 1995
- Rosenbaum M, Goldsmith R, Bloomfield D, Magnano A, Weimer L, Heymsfield S, Gallagher D, Mayer L, Murphy E, Leibel RL: Low-dose leptin reverses skeletal muscle, autonomic, and neuroendocrine adaptations to maintenance of reduced weight. *J Clin Invest* 115:3579–3586, 2005
- Anthony JC, Anthony TG, Kimball SR, Jefferson LS: Signaling pathways involved in translational control of protein synthesis in skeletal muscle by leucine. *J Nutr* 131:856S–860S, 2001
- Gingras AC, Raught B, Sonenberg N: Regulation of translation initiation by FRAP/mTOR. *Genes Dev* 15:807–826, 2001
- Kimball SR, Jefferson LS: Signaling pathways and molecular mechanisms through which branched-chain amino acids mediate translational control of protein synthesis. *J Nutr* 136:227S–231S, 2006
- Inoki K, Corradetti MN, Guan KL: Dysregulation of the TSC-mTOR pathway in human disease. *Nat Genet* 37:19–24, 2005
- Cota D, Proulx K, Smith KA, Kozma SC, Thomas G, Woods SC, Seeley RJ: Hypothalamic mTOR signaling regulates food intake. *Science* 312:927–930, 2006
- Anthony JC, Anthony TG, Layman DK: Leucine supplementation enhances skeletal muscle recovery in rats following exercise. *J Nutr* 129:1102–1106, 1999
- Dardevet D, Sornet C, Bayle G, Prugnaud J, Pouyet C, Grizard J: Postprandial stimulation of muscle protein synthesis in old rats can be restored by a leucine-supplemented meal. *J Nutr* 132:95–100, 2002
- Gomes-Marcondes MC, Ventrucci G, Toledo MT, Cury L, Cooper JC: A leucine-supplemented diet improved protein content of skeletal muscle in young tumor-bearing rats. *Braz J Med Biol Res* 36:1589–1594, 2003
- Levenhagen DK, Carr C, Carlson MG, Maron DJ, Borel MJ, Flakoll PJ: Postexercise protein intake enhances whole-body and leg protein accretion in humans. *Med Sci Sports Exerc* 34:828–837, 2002
- Shimomura Y, Yamamoto Y, Bajotto G, Sato J, Murakami T, Shimomura N, Kobayashi H, Mawatari K: Nutraceutical effects of branched-chain amino acids on skeletal muscle. *J Nutr* 136:529S–532S, 2006
- Lynch CJ, Hutson SM, Patson BJ, Vaval A, Vary TC: Tissue-specific effects of chronic dietary leucine and norleucine supplementation on protein synthesis in rats. *Am J Physiol Endocrinol Metab* 283:E824–E835, 2002
- Patti ME, Brambilla E, Luzi L, Landaker EJ, Kahn CR: Bidirectional modulation of insulin action by amino acids. *J Clin Invest* 101:1519–1529, 1998
- Tremblay F, Marette A: Amino acid and insulin signaling via the mTOR/p70 S6 kinase pathway: a negative feedback mechanism leading to insulin resistance in skeletal muscle cells. *J Biol Chem* 276:38052–38060, 2001
- Krebs M, Krssak M, Bernroider E, Anderwald C, Brehm A, Meyerspeer M, Nowotny P, Roth E, Waldhausl W, Roden M: Mechanism of amino acid-induced skeletal muscle insulin resistance in humans. *Diabetes* 51:599–605, 2002
- Um SH, Frigerio F, Watanabe M, Picard F, Joaquin M, Sticker M, Fumagalli S, Allegri PR, Kozma SC, Auwerx J, Thomas G: Absence of S6K1 protects against age- and diet-induced obesity while enhancing insulin sensitivity. *Nature* 431:200–205, 2004
- Layman DK, Boileau RA, Erickson DJ, Painter JE, Shiue H, Sather C, Christou DD: A reduced ratio of dietary carbohydrate to protein improves body composition and blood lipid profiles during weight loss in adult women. *J Nutr* 133:411–417, 2003
- Layman DK, Shiue H, Sather C, Erickson DJ, Baum J: Increased dietary protein modifies glucose and insulin homeostasis in adult women during weight loss. *J Nutr* 133:405–410, 2003
- Skov AR, Toubro S, Ronn B, Holm L, Astrup A: Randomized trial on protein vs carbohydrate in ad libitum fat reduced diet for the treatment of obesity. *Int J Obes Relat Metab Disord* 23:528–536, 1999
- Parker B, Noakes M, Luscombe N, Clifton P: Effect of a high-protein, high-monounsaturated fat weight loss diet on glycemic control and lipid levels in type 2 diabetes. *Diabetes Care* 25:425–430, 2002
- Layman DK, Walker DA: Potential importance of leucine in treatment of obesity and the metabolic syndrome. *J Nutr* 136:319S–323S, 2006
- Donato J Jr, Pedrosa RG, Cruzat VF, Pires IS, Tirapegui J: Effects of leucine supplementation on the body composition and protein status of rats submitted to food restriction. *Nutrition* 22:520–527, 2006
- Guo KY, Halo P, Leibel RL, Zhang Y: Effects of obesity on the relationship of leptin mRNA expression and adipocyte size in anatomically distinct fat depots in mice. *Am J Physiol Regul Integr Comp Physiol* 287:R112–R119, 2004
- Yu YH, Zhu H: Chronological changes in metabolism and functions of cultured adipocytes: a hypothesis for cell aging in mature adipocytes. *Am J Physiol Endocrinol Metab* 286:E402–E410, 2004
- Trinh KY, O'Doherty RM, Anderson P, Lange AJ, Newgard CB: Perturbation of fuel homeostasis caused by overexpression of the glucose-6-phosphatase catalytic subunit in liver of normal rats. *J Biol Chem* 273:31615–31620, 1998
- Yoon JC, Xu G, Deeney JT, Yang SN, Rhee J, Puigserver P, Levens AR, Yang R, Zhang CY, Lowell BB, Berggren PO, Newgard CB, Bonner-Weir S, Weir G, Spiegelman BM: Suppression of beta cell energy metabolism and insulin release by PGC-1alpha. *Dev Cell* 5:73–83, 2003
- Chen Q, Chan LL, Li ET: Bitter melon (*Momordica charantia*) reduces adiposity, lowers serum insulin and normalizes glucose tolerance in rats fed a high fat diet. *J Nutr* 133:1088–1093, 2003
- Hu CC, Qing K, Chen Y: Diet-induced changes in stearoyl-CoA desaturase 1 expression in obesity-prone and -resistant mice. *Obes Res* 12:1264–1270, 2004
- Corbett SW, Stern JS, Keeseey RE: Energy expenditure in rats with diet-induced obesity. *Am J Clin Nutr* 44:173–180, 1986
- Weigle DS, Levin BE: Defective dietary induction of uncoupling protein 3 in skeletal muscle of obesity-prone rats. *Obes Res* 8:385–391, 2000
- Son C, Hosoda K, Ishihara K, Bevilacqua L, Masuzaki H, Fushiki T, Harper ME, Nakao K: Reduction of diet-induced obesity in transgenic mice overexpressing uncoupling protein 3 in skeletal muscle. *Diabetologia* 47:47–54, 2004
- Rodriguez AM, Roca P, Bonet ML, Pico C, Oliver P, Palou A: Positive correlation of skeletal muscle UCP3 mRNA levels with overweight in male, but not in female, rats. *Am J Physiol Regul Integr Comp Physiol* 285:R880–R888, 2003
- Bao S, Kennedy A, Wojciechowski B, Wallace P, Ganaway E, Garvey WT: Expression of mRNAs encoding uncoupling proteins in human skeletal muscle: effects of obesity and diabetes. *Diabetes* 47:1935–1940, 1998
- Vidal-Puig A, Rosenbaum M, Considine RC, Leibel RL, Dohm GL, Lowell BB: Effects of obesity and stable weight reduction on UCP2 and UCP3 gene expression in humans. *Obes Res* 7:133–140, 1999
- Pedersen SB, Lund S, Buhl ES, Richelsen B: Insulin and contraction directly stimulate UCP2 and UCP3 mRNA expression in rat skeletal muscle in vitro. *Biochem Biophys Res Commun* 283:19–25, 2001
- Shimada M, Tritos NA, Lowell BB, Flier JS, Maratos-Flier E: Mice lacking melanin-concentrating hormone are hypophagic and lean. *Nature* 396:670–674, 1998
- Baker DH: Tolerance for branched-chain amino acids in experimental animals and humans. *J Nutr* 135:1585S–1590S, 2005
- Chevalier S, Burgess SC, Malloy CR, Gougeon R, Marliss EB, Morais JA: The greater contribution of gluconeogenesis to glucose production in obesity is related to increased whole-body protein catabolism. *Diabetes* 55:675–681, 2006
- Leclercq-Meyer V, Marchand J, Woussen-Colle MC, Giroix MH, Malaisse WJ: Multiple effects of leucine on glucagon, insulin, and somatostatin secretion from the perfused rat pancreas. *Endocrinology* 116:1168–1174, 1985
- Kuhara T, Ikeda S, Ohneda A, Sasaki Y: Effects of intravenous infusion of 17 amino acids on the secretion of GH, glucagon, and insulin in sheep. *Am J Physiol* 260:E21–E26, 1991
- Bassett JM, Madill D, Burks AH, Pinches RA: Glucagon and insulin release in the lamb before and after birth: effects of amino acids in vitro and in vivo. *J Dev Physiol* 4:379–389, 1982
- Miguez I, Marino G, Rodriguez B, Taboada C: Effects of dietary L-arginine supplementation on serum lipids and intestinal enzyme activities in diabetic rats. *J Physiol Biochem* 60:31–37, 2004
- Teplan V, Schuck O, Votruba M, Poledne R, Kazdova L, Skibova J, Maly J: Metabolic effects of keto acid-amino acid supplementation in patients with chronic renal insufficiency receiving a low-protein diet and recombinant human erythropoietin—a randomized controlled trial. *Wien Klin Wochenschr* 113:661–669, 2001