

Association of *CDKAL1*, *IGF2BP2*, *CDKN2A/B*, *HHEX*, *SLC30A8*, and *KCNJ11* With Susceptibility to Type 2 Diabetes in a Japanese Population

Shintaro Omori,^{1,2} Yasushi Tanaka,² Atsushi Takahashi,³ Hiroshi Hirose,⁴ Atsunori Kashiwagi,⁵ Kohei Kaku,⁶ Ryuzo Kawamori,⁷ Yusuke Nakamura,⁸ and Shiro Maeda¹

OBJECTIVE—Recently, several genes have been shown to be associated with an increased risk of type 2 diabetes by genome-wide association studies in white populations. To further investigate the involvement of these polymorphisms in conferring susceptibility to type 2 diabetes, we examined the association of 14 single nucleotide polymorphisms (SNPs) within 11 candidate loci with type 2 diabetes in a Japanese population.

RESEARCH DESIGN AND METHODS—We analyzed 14 SNPs (rs4402960 in *IGF2BP2*, rs10811661 in *CDKN2A/B*, rs1111875 and rs7923837 in *HHEX*, rs13266634 in *SLC30A8*, rs1113132 and rs11037909 in *EXT2*, rs9939609 and rs8050136 in *FTO*, rs7756992 in *CDKAL1*, rs1801282 in *PPARG* Pro12Ara, rs5219 in *KCNJ11* Glu23Lys, rs7480010 in *LOC387761*, and rs9300039 in Ch11) in 1,630 Japanese subjects with type 2 diabetes and in 1,064 control subjects by using an invader assay or a TaqMan assay.

RESULTS—Among the 11 loci examined, 6 were significantly associated with type 2 diabetes in our population by a logistic regression analysis, similar to previously reported results (rs4402960, $P = 0.00009$; rs10811661, $P = 0.0024$; rs5219, $P = 0.0034$; rs1111875, $P = 0.0064$; rs13266634, $P = 0.0073$; rs7756992, $P = 0.0363$). In this population, the remaining five loci were not significantly associated with type 2 diabetes. In addition, we identified significant association of the SNPs in *FTO* gene with BMI in the control subjects.

CONCLUSIONS—We have identified 6 of the 11 loci that were identified by genome-wide association studies in white populations, and these loci are considered strong candidates for type 2 diabetes susceptibility across different ethnicities. *Diabetes* 57: 791–795, 2008

From the ¹Laboratory for Diabetic Nephropathy, SNP Research Center, RIKEN, Yokohama, Kanagawa, Japan; the ²Department of Internal Medicine, Division of Metabolism and Endocrinology, St. Marianna University School of Medicine, Kawasaki, Kanagawa, Japan; the ³Laboratory for Statistical Analysis, SNP Research Center, RIKEN, Tokyo, Japan; ⁴Health Center, Keio University School of Medicine, Tokyo, Japan; the ⁵Department of Medicine, Shiga University of Medical Science, Otsu, Shiga, Japan; the ⁶Division of Endocrinology and Metabolism, Department of Internal Medicine, Kawasaki Medical School, Kurashiki, Okayama, Japan; the ⁷Department of Medicine, Metabolism and Endocrinology, School of Medicine, Juntendo University, Tokyo, Japan; and the ⁸Laboratory of Molecular Medicine, Human Genome Center, Institute of Medical Science, University of Tokyo, Tokyo, Japan.

Address correspondence and reprint requests to Shiro Maeda, Laboratory for Diabetic Nephropathy, SNP Research Center, RIKEN, 1-7-22 Suehiro-cho, Tsurumi-ku, Yokohama, Kanagawa 230-0045, Japan. E-mail: smaeda@src.riken.jp.

Received for publication 14 July 2007 and accepted in revised form 12 December 2007.

Published ahead of print at <http://diabetes.diabetesjournals.org> on 27 December 2007. DOI: 10.2337/db07-0979.

Additional information for this article can be found in an online appendix at <http://dx.doi.org/10.2337/db07-0979>.

SNP, single nucleotide polymorphism.

© 2008 by the American Diabetes Association.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Type 2 diabetes affects >200 million individuals worldwide, and its prevalence continues to increase in many countries, including Japan. Although the precise mechanisms underlying the development and progression of type 2 diabetes have not been elucidated, a combination of multiple genetic and/or environmental factors is considered to contribute to the pathogenesis of the disease (1).

Recently, genome-wide association studies conducted by several independent European and American groups have identified multiple susceptible variants in white populations including *TCF7L2* variants, which had been originally identified by a genome-wide linkage study (2) and confirmed in several replication studies across different ethnicities (3–7). Sladek et al. (8) additionally identified the solute carrier family 30 member 8 (*SLC30A8*), homeobox hematopoietically expressed (*HHEX*), *LOC387761*, and exostosin 2 (*EXT2*) genes, and the WTCCC/UKT2D, FUSION, and DGI study groups identified the insulin-like growth factor 2 mRNA binding protein 2 (*IGF2BP2*), cyclin-dependent kinase 5 regulatory subunit associated protein 1-like 1 (*CDKAL1*), cyclin-dependent kinase inhibitor-2A/B (*CDKN2A/B*), fat mass- and obesity-associated (*FTO*) genes, and the rs9300039 locus as additional loci that were strongly associated with the susceptibility to this disease (9–11). The latter study also confirmed the association between the susceptibility to type 2 diabetes and the peroxisome proliferators-activated receptor- γ (*PPARG*) Pro12Ala or potassium inwardly rectifying channel subfamily J member 11 (*KCNJ11*) Glu23Lys polymorphism that had already been reported as strong candidates. The associations among single nucleotide polymorphisms (SNPs) within *CDKAL1* and *SLC30A8* were also identified by a genome-wide association study conducted for the Icelandic population (12).

These additional loci are also considered to be strong candidates for conferring susceptibility to type 2 diabetes in white populations. However, the contributions of these new loci should be evaluated in other ethnic populations, because it is well known that there are significant differences in the frequencies of some genetic variations among different ethnic groups (6,7,13).

The aim of the present study is to determine whether the variations identified by the genome-wide association studies in white populations are associated with the susceptibility to type 2 diabetes in a Japanese population.

TABLE 1
Clinical characteristics of the subjects

	Type 2 diabetic subjects	Control subjects	<i>P</i> *
<i>n</i>	1,630	1,064	
Sex (M/F)	978/652	638/426	0.9845
Age (years)	61.5 ± 11.6	45.5 ± 9.5	<0.0001
BMI (kg/m ²)	23.7 ± 3.9	22.9 ± 3.0	<0.0001
FPG (mmol/l)	9.1 ± 3.5	5.1 ± 0.5	<0.0001
A1C (%)	7.4 ± 1.6	4.7 ± 0.4	<0.0001
Duration of diabetes (years)	11.5 ± 13.9	—	—

Data are means ± SE unless otherwise indicated. *Pearson's χ^2 test.

RESEARCH DESIGN AND METHODS

Subject and DNA preparation. DNA samples were obtained from the peripheral blood samples of 1,630 type 2 diabetic patients recruited from the outpatient clinic of the Shiga University of Medical Science, Kawasaki Medical School (978 men and 652 women; age 61.5 ± 11.6 years; duration of diabetes 11.5 ± 13.9 years; A1C 7.4 ± 1.6%; fasting plasma glucose 9.1 ± 3.5 mmol/l; BMI 23.7 ± 3.9 kg/m² [all values are expressed as means ± SD], Table 1). Diabetes was diagnosed according to the WHO criteria. Type 2 diabetes is clinically defined as a disease with gradual adult onset. Subjects who tested positive for anti-GAD antibodies and those diagnosed as mitochondrial disease (mitochondrial myopathy, encephalopathy, lactic acidosis, and stroke-like episodes [MELAS]) or maturity-onset diabetes of young (MODY) were not included in the case group. We also examined 1,064 control subjects who were enrolled from an annual health check conducted either at the Juntendo University or Keio University (Tokyo, Japan; 638 men and 426 women; age 45.5 ± 9.5 years; A1C 4.7 ± 0.4%; fasting plasma glucose 5.1 ± 0.5 mmol/l; BMI 22.9 ± 3.0 kg/m²; Table 1).

Written informed consent was obtained from all the participants, and DNA was extracted using the standard phenol-chloroform procedure. The protocol was approved by the ethics committee of the Institute of Physical and Chemical Research (RIKEN).

Genotyping. Each SNP genotyping was performed by the TaqMan assay (Applied Biosystems, Foster City, CA) or by the multiplex-PCR invader assay as described previously (13). The success rates of these assays were >95%, and there was almost a 100% agreement between the results of genotyping and direct sequencing.

Statistical analysis. Statistical methods for determining associations and to calculate linkage disequilibrium (LD) coefficients (r^2) were described previously (14). We performed the Hardy-Weinberg equilibrium (HWE) test according to the method described by Nielsen et al. (15). Although the genotype distributions of all the SNPs were within the Hardy-Weinberg equilibrium ($P \geq 0.01$), some of them had borderline results for HWE test (rs5219 in control, rs7480010 in case, rs8050136 in case; see supplementary Table 1 at <http://dx.doi.org/10.2337/db07-0979>). Therefore, we performed Wright's *F* statistics (16) to evaluate the difference in the population structure between our case and control groups using randomly selected 96 SNPs. The result indicated that the population structures of our case and control groups were almost the

same in view of a very small F_{ST} value between the two groups ($F_{ST} = 0.001556$).

The differences between the case and control groups in terms of genotype distribution were analyzed using a logistic regression analysis. To test the additive model of each SNP after adjusting for sex, age, and log-transformed BMI, the analysis was performed using StatView software. In addition, χ^2 test to evaluate the additive, dominant, and recessive models of each SNP were also performed by the method of Sladek et al. (8).

The difference in the BMI according to the genotypes was analyzed using a multiple linear regression with log-transformed BMI as the dependent variable and genotype as the independent variable with sex as a covariate for log-transformed BMI (17).

RESULTS

As shown in Table 2, six SNPs within six distinct loci (rs4402960, rs10811661, rs5219, rs1111875, rs13266634, and rs7756992) were found to be significantly associated with type 2 diabetes in our Japanese population. No significant association was observed between the remaining five loci and type 2 diabetes in this population ($P \geq 0.05$; Table 2). We also evaluated the association of the 14 SNP loci with type 2 diabetes using χ^2 test by the method of Sladek et al. (8) (supplementary Table 2) and identified almost consistent results with those obtained by a logistic regression analysis even after selecting control subjects whose age was >50 years old ($n = 382$, supplementary Table 3). Since *FTO* variants have been reported to be associated with BMI, we also examined the association of these polymorphisms with BMI in our control subjects, and we found that the polymorphisms within *FTO* and *HHEX* were modestly associated with BMI (Table 3).

TABLE 2
Association of candidate SNP loci with type 2 diabetes

SNP	Gene	<i>P</i> *	Odds ratio (95% CI)
rs4402960	<i>IGF2BP2</i>	0.00009	1.368 (1.169-1.600)
rs10811661	<i>CDKN2A/B</i>	0.0024	1.255 (1.084-1.454)
rs5219	<i>KCNJ11</i>	0.0034	1.254 (1.078-1.459)
rs1111875†	<i>HHEX</i>	0.0064	1.243 (1.063-1.453)
rs7923837†	<i>HHEX</i>	0.3773	1.083 (0.907-1.293)
rs13266634	<i>SLC30A8</i>	0.0073	1.225 (1.056-1.420)
rs7756992	<i>CDKAL1</i>	0.0363	1.164 (1.010-1.342)
rs9939609	<i>FTO</i>	0.2376	1.114 (0.931-1.332)
rs8050136	<i>FTO</i>	0.3520	1.089 (0.910-1.302)
rs1801282	<i>PPARG</i>	0.4137	0.843 (0.559-1.270)
rs7480010	<i>LOC387761</i>	0.4393	1.073 (0.898-1.281)
rs1113132	<i>EXT2</i>	0.4728	1.056 (0.910-1.225)
rs11037909	<i>EXT2</i>	0.5365	1.048 (0.903-1.216)
rs9300039	41871942‡	0.6966	1.034 (0.874-1.222)

**P* value is calculated on logistic regression with additive model (sex, age, BMI adjusted, and BMI were log transformed for the analysis); † $r^2 = 0.50$ (this study), 0.346 (HapMap-JPT), and 0.698 (HapMap-CEU), respectively. ‡Position on the chromosome is indicated.

TABLE 3
Association of the SNP loci with BMI in control subjects

SNP/gene	Genotype (number of subjects)/BMI*			P
rs9939609 <i>FTO</i>	TT (676) 22.2 (21.9–22.4)	AT (331) 23.0 (22.7–23.4)	AA (37) 23.1 (22.5–23.8)	0.0271
rs8050136 <i>FTO</i>	CC (678) 22.2 (22.0–22.4)	CA (331) 23.0 (22.7–23.3)	AA (35) 23.2 (22.5–23.9)	0.0436
rs7923837 <i>HHEX</i>	AA (653) 22.4 (22.2–22.6)	AG (333) 22.5 (22.2–22.8)	GG (46) 22.7 (21.9–23.5)	0.032
rs1111875 <i>HHEX</i>	TT (529) 22.5 (22.2–22.7)	CT (419) 22.5 (22.2–22.7)	CC (84) 22.3 (21.7–22.9)	0.05
rs5219 <i>KCNJ11</i> Glu23Lys	CC (421) 22.4 (22.1–22.7)	CT (509) 22.6 (22.3–22.8)	TT (118) 22.1 (21.6–22.6)	0.0777
rs1801282 <i>PPARG</i> Pro12Ala	CC (2) 24.4	CG (53) 23.1 (22.4–23.9)	GG (995) 22.4 (22.2–22.6)	0.1039
rs11037909 <i>EXT2</i>	CC (141) 22.4 (21.9–22.9)	CT (471) 22.7 (22.4–22.9)	TT (425) 22.3 (22.0–22.6)	0.1243
rs1113132 <i>EXT2</i>	CC (143) 22.4 (21.9–22.9)	CG (467) 22.6 (22.4–22.9)	GG (427) 22.3 (22.0–22.6)	0.1315
rs9300039 41871942†	AA (68) 22.7 (21.9–23.5)	AC (370) 22.5 (22.2–22.8)	CC (592) 22.4 (22.2–22.6)	0.4153
rs13266634 <i>SLC30A8</i>	TT (173) 22.6 (22.1–23.1)	CT (491) 22.5 (22.2–22.7)	CC (376) 22.4 (22.2–22.7)	0.6014
rs10811661 <i>CDKN2A/B</i>	CC (200) 22.3 (21.9–22.7)	CT (518) 22.5 (22.2–22.7)	TT (326) 22.5 (22.2–22.8)	0.6185
rs7480010 <i>LOC387761</i>	AA (684) 22.4 (22.1–22.6)	AG (317) 22.8 (22.5–23.1)	GG (42) 22.3 (21.5–23.2)	0.7183
rs7756992 <i>CDKAL1</i>	AA (289) 22.4 (22.0–22.7)	AG (508) 22.6 (22.4–22.9)	GG (236) 22.3 (22.0–22.7)	0.7985
rs4402960 <i>IGF2BP2</i>	GG (520) 22.5 (22.3–22.8)	GT (433) 22.4 (22.1–22.7)	TT (88) 22.5 (21.8–23.1)	0.9313

*Data are presented as geometric means, and values for 95% CI are in parentheses; †position on the chromosome is indicated.

DISCUSSION

In the present study, we identified significant associations of SNPs within the *CDKAL1*, *IGF2BP2*, *CDKN2A/B*, *HHEX*, *SLC30A8* and *KCNJ11* genes with the susceptibility to type 2 diabetes in a Japanese population. We also found that the SNPs in the *FTO* gene and the *HHEX* gene were significantly associated with BMI in our control group.

Recent advances in human genetic research have facilitated the identification of genes conferring susceptibility to common diseases such as type 2 diabetes from across the entire human genome by using a large number of subjects, and genome-wide association studies have been conducted worldwide (8–12). Although the importance of *TCF7L2* as a susceptibility gene for type 2 diabetes has been well established, its polymorphism could account for ~20% of all cases for type 2 diabetes in white populations (2,4,5). The population-attributable risk of the *TCF7L2* polymorphism in the Japanese was ~2% because the risk allelic frequency in a previously studied Japanese population was very low (6,7). Further, many important genes for the disease remain to be identified, especially in East Asian populations.

Several groups have independently performed genome-wide association studies for type 2 diabetes in white populations (8–12). All these studies have demonstrated that the *TCF7L2* polymorphism is most strongly associated with the susceptibility to type 2 diabetes, and from a large set of replication studies, they have identified additional candidate loci also associated with the disease. The results of these genome-wide association studies are also considered highly consistent with regard to white popula-

tions. However, there are considerable differences in phenotype between Japanese (lean and less hyperinsulinemic Asian type 2 diabetes) and white (European descent) type 2 diabetes, and these differences might affect the genetic contribution of each gene to conferring susceptibility to type 2 diabetes. Therefore, the new candidates should also be evaluated in different ethnic groups because there are clear ethnic differences in terms of genetic contribution to diseases (13,18).

In the present study, we also identified the six SNPs within the *CDKAL1*, *IGF2BP2*, *CDKN2A/B*, *HHEX*, *SLC30A8*, and *KCNJ11* genes that were significantly associated with type 2 diabetes in our Japanese population. Since the risk alleles for these variations in the tested population were entirely consistent with those in white populations (8–12), the contribution of these polymorphisms to type 2 diabetes susceptibility is highly convincing across the different ethnicities, although there are considerable differences in allele frequencies between the Japanese and white populations (Table 4 and supplementary Table 4). Regarding the *HHEX* locus, only rs1111875 had significant association with type 2 diabetes in the present study, whereas both rs1111875 and rs7923837 were associated with the disease in white populations. The risk allele frequencies of SNPs in the *HHEX* gene were significantly different between our population and white populations (rs1111875 28.4 vs. 56.1%, rs7923837 20.6 vs. 60.1% in Japanese and white populations, respectively; supplementary Table 4), and there were some differences in the LD coefficients (r^2) between rs1111875 and rs7923837 among those populations (0.5 in the Japanese and 0.698 in the white population). Therefore, these dif-

TABLE 4

The comparison of risk allele frequency and population-attributable risk (PAR) between Japanese and white populations

SNP	Gene	Risk allele frequency (%)		<i>P</i> *	PAR (%)	
		This study	White populations		This study	White populations†
rs4402960	<i>IGF2BP2</i>	29.3	31.3	0.1439	11.1	4.8
rs10811661	<i>CDKN2A/B</i>	56.1	84.1	2.2×10^{-183}	13.4	18.1
rs5219	<i>KCNJ11</i>	35.5	46.4	1.9×10^{-16}	9.1	5.7
rs1111875	<i>HHEX</i>	28.4	57.7	3.2×10^{-146}	7.3	19.0
rs7923837	<i>HHEX</i>	20.6	62.3	3.6×10^{-218}	—	20.0
rs13266634	<i>SLC30A8</i>	60.0	63.4	0.0097	12.5	14.6
rs7756992	<i>CDKAL1</i>	47.4	23.2	3.0×10^{-114}	7.7	6.1
rs9939609	<i>FTO</i>	19.4	38.5	1.3×10^{-19}	—	—
rs8050136	<i>FTO</i>	19.3	39.0	6.0×10^{-67}	—	—
rs1801282	<i>PPARG</i>	97.3	82.4	4.2×10^{-63}	—	14.2
rs7480010	<i>LOC387761</i>	19.1	30.1	9.5×10^{-21}	—	8.0
rs1113132	<i>EXT2</i>	63.6	73.3	1.8×10^{-16}	—	19.0
rs11037909	<i>EXT2</i>	63.6	72.9	2.6×10^{-14}	—	25.0
rs9300039	41871942‡	75.3	89.2	2.5×10^{-46}	—	30.6

**P* values for χ^2 test for genotype distribution (2×3 contingency table); †combined data from WTCCC, UKRS, FUSION, French, and Icelandic studies; ‡position on the chromosome is indicated.

ferences might explain the discrepancies in the results for the association of SNPs in the *HHEX* gene with type 2 diabetes, although the possibility of insufficient study power in the present study could not be excluded.

We also found that the SNPs within *FTO* (rs9939609 and rs8050136) and *HHEX* (rs1111875 and rs7923837) were modestly associated with BMI in our control subjects (Table 3). The association of the SNPs in *FTO* with type 2 diabetes was not significant in our population (Table 2 and supplementary Tables 2 and 3). Therefore, the variations in the *FTO* gene might directly affect body weight rather than type 2 diabetes itself; this finding is also consistent with that in white populations (17,19).

Among the other four loci, the association of the Pro12Ala polymorphism in the *PPARG* gene with type 2 diabetes has been well established (20,21), and this association was also observed in Japanese populations (22,23), although we could not determine whether the association of this polymorphism with type 2 diabetes was significant. Because the frequencies of the Ala allele or its carrier (X/Ala) found in the present study (three and 6%, respectively) are consistent with those in previous studies on Japanese populations, the frequencies of the Ala allele can be considered very low in the Japanese as compared to those in white populations. Because an estimated power to detect the association of the SNP with type 2 diabetes in the present study is ~20%, the results do not appear strong enough to determine the association between the *PPARG* Pro12Ala polymorphism and type 2 diabetes.

With regard to the remaining three loci, the results for *EXT2*, *LOC387761*, and rs9300039 were not always in agreement with those of the genome-wide association studies in white populations. In addition, the allele frequencies of those SNPs were also significantly different between the Japanese and the white populations (Table 4 and supplementary Table 4). Because the estimated powers of the present study were >90%, >90%, and >70% for rs1113132 (*EXT2*), rs7480010 (*LOC387761*), and rs9300039, respectively, the contribution of these three loci in the Japanese populations is considered minor, if present at all; however, more replication studies are required for the precise evaluation of these loci.

In conclusion, we identified significant associations be-

tween SNPs within the *CDKAL1*, *IGF2BP2*, *CDKN2A/B*, *HHEX*, *SLC30A8*, and *KCNJ11* genes and type 2 diabetes in a Japanese population. These loci are considered strong candidates for conferring susceptibility to type 2 diabetes across different ethnicities. However, further studies are required to elucidate the association of other loci with the susceptibility to type 2 diabetes and the biological significance of these genes and gene polymorphisms.

ACKNOWLEDGMENTS

We thank Shuichi Tsukada, Masa-aki Kobayashi, and the technical staff of the Laboratory for Diabetic Nephropathy at the SNP Research Center for providing technical assistance. This work was partly supported by the Japanese Millennium Project.

REFERENCES

- O'Rahilly S, Barroso I, Wareham NJ: Genetic factors in type 2 diabetes: the end of the beginning? *Science* 307:370–373, 2005
- Grant SF, Thorleifsson G, Reynisdottir I, et al: Variant of transcription factor 7-like 2 (*TCF7L2*) gene confers risk of type 2 diabetes. *Nat Genet* 38:320–323, 2006
- Florez JC, Jablonski KA, Bayley N, Pollin TI, de Bakker PI, Shuldiner AR, Knowler WC, Nathan DM, Altshuler D, Diabetes Prevention Program Research Group: *TCF7L2* polymorphisms and progression to diabetes in the diabetes prevention program. *N Engl J Med* 355:241–250, 2006
- Groves CJ, Zeggini E, Minton J, Frayling TM, Weedon MN, Rayner NW, Hitman GA, Walker M, Wiltshire S, Hattersley AT, McCarthy MI: Association analysis of 6,736 U.K. subjects provides replication and confirms *TCF7L2* as a type 2 diabetes susceptibility gene with a substantial effect on individual risk. *Diabetes* 55:2640–2644, 2006
- Zhang C, Qi L, Hunter DJ, Meigs JB, Manson JE, van Dam RM, Hu FB: Variant of transcription factor 7-like 2 (*TCF7L2*) gene and the risk of type 2 diabetes in large cohorts of U.S. women and men. *Diabetes* 55:2645–2648, 2006
- Hayashi T, Iwamoto Y, Kaku K, Hirose H, Maeda S: Replication study for the association of *TCF7L2* with susceptibility to the type 2 diabetes in a Japanese population. *Diabetologia* 50:980–984, 2007
- Horikoshi M, Hara K, Ito C, Nagai R, Froguel P, Kadowaki T: A genetic variation of the transcription factor 7-like 2 gene is associated with risk of type 2 diabetes in the Japanese population. *Diabetologia* 50:747–751, 2007
- Sladek R, Rocheleau G, Rung J, et al: A genome-wide association study identifies novel risk loci for type 2 diabetes. *Nature* 445:881–885, 2007
- Zeggini E, Weedon MN, Lindgren CM, et al: Replication of genome-wide association signal in U.K. samples reveals risk loci for type 2 diabetes. *Science* 316:1336–1341, 2007

10. Scott LJ, Mohlke KL, Bonnycastle LL, et al.: A genome-wide association study of type 2 diabetes in Finns detects multiple susceptibility variants. *Science* 316:1341–1345, 2007
11. Diabetes Genetics Initiative of Broad Institute of Harvard and MIT, Lund University, and Novartis Institutes of BioMedical Research, Saxena R, Voight BF, Lyssenko V, et al. Gage D, Nizzari M, Gabriel SB, Chirn GW, Ma Q, Parikh H, Richardson D, Ricke D, Purcell S: Genome-wide association analysis identifies loci for type 2 diabetes and triglyceride levels. *Science* 316:1331–1336, 2007
12. Steinthorsdottir V, Thorleifsson G, Reynisdottir I, et al.: A variant in *CDKAL1* influences insulin response and risk of type 2 diabetes. *Nat Genet* 39:770–775, 2007
13. Maeda S, Tsukada S, Kanazawa A, et al.: Genetic variations in the gene encoding *TFAP2B* are associated with type 2 diabetes mellitus. *J Hum Genet* 50:283–292, 2005
14. Devlin B, Risch N: A comparison of linkage disequilibrium measures for fine-scale mapping. *Genomics* 20:311–322, 1995
15. Nielsen DM, Ehm MG, Weir BS: Detecting marker-disease association by testing for Hardy-Weinberg disequilibrium at a marker locus. *Am J Hum Genet* 63:1531–1540, 1998
16. Wright S: *Evolution and the Genetics of Populations. Volume 2: The Theory of Gene Frequencies*. Chicago, IL, University of Chicago Press, 1969, p. 295–295
17. Frayling TM, Timpson NJ, Weedon MN, et al.: A common variant in the *FTO* gene is associated with body mass index and predisposes to childhood and adult obesity. *Science* 316:889–894, 2007
18. The International HapMap Consortium: A haplotype map of the human genome. *Nature* 437:1299–1320, 2005
19. Dina C, Meyre D, Gallina S, et al.: Variation in *FTO* contributes to childhood obesity and severe adult obesity. *Nat Genet* 39:724–726, 2007
20. Altshuler D, Hirschhorn JN, Klannemark M, Lindgren CM, Vohl MC, Nemesh J, Lane CR, Schaffner SF, Bolk S, Brewer C, Tuomi T, Gaudet D, Hudson TJ, Daly M, Groop L, Lander ES: The common PPARgamma Pro12Ala polymorphism is associated with decreased risk of type 2 diabetes. *Nat Genet* 26:76–80, 2000
21. Tonjes A, Loeffler M, Scholz M, Stumvoll M: Association of Pro12Ala Polymorphism in peroxisome proliferator-activated receptor gamma with pre-diabetic phenotypes. *Diabetes Care* 29:2489–2497, 2006
22. Hara K, Okada T, Tobe K, Yasuda K, Mori Y, Kadowaki H, Hagura R, Akanuma Y, Kimura S, Ito C, Kadowaki T: The Pro12Ala polymorphism in PPAR gamma 2 may confer resistance to type 2 diabetes. *Biochem Biophys Res Commun* 271:212–216, 2000
23. Kawasaki I, Tahara H, Emoto M, Shoji T, Shioji A, Okuno Y, Inaba M, Nishizawa Y: Impact of Pro12Ala variant in the peroxisome proliferator-activated receptor (PPAR) gamma2 on obesity and insulin resistance in Japanese type 2 diabetic and healthy subjects. *Osaka City Med J* 48:23–28, 2002