

Loss of Both ABCA1 and ABCG1 Results in Increased Disturbances in Islet Sterol Homeostasis, Inflammation, and Impaired β -Cell Function

Janine K. Kruit,¹ Nadeeja Wijesekara,¹ Clara Westwell-Roper,² Tim Vanmierlo,³ Willeke de Haan,¹ Alpna Bhattacharjee,¹ Renmei Tang,¹ Cheryl L. Wellington,² Dieter Lütjohann,³ James D. Johnson,⁴ Liam R. Brunham,¹ C. Bruce Verchere,⁵ and Michael R. Hayden¹

Cellular cholesterol homeostasis is important for normal β -cell function. Disruption of cholesterol transport by decreased function of the ATP-binding cassette (ABC) transporter ABCA1 results in impaired insulin secretion. Mice lacking β -cell ABCA1 have increased islet expression of ABCG1, another cholesterol transporter implicated in β -cell function. To determine whether ABCA1 and ABCG1 have complementary roles in β -cells, mice lacking ABCG1 and β -cell ABCA1 were generated and glucose tolerance, islet sterol levels, and β -cell function were assessed. Lack of both ABCG1 and β -cell ABCA1 resulted in increased fasting glucose levels and a greater impairment in glucose tolerance compared with either ABCG1 deletion or loss of ABCA1 in β -cells alone. In addition, glucose-stimulated insulin secretion was decreased and sterol accumulation increased in islets lacking both transporters compared with those isolated from knockout mice with each gene alone. Combined deficiency of ABCA1 and ABCG1 also resulted in significant islet inflammation as indicated by increased expression of interleukin-1 β and macrophage infiltration. Thus, lack of both ABCA1 and ABCG1 induces greater defects in β -cell function than deficiency of either transporter individually. These data suggest that ABCA1 and ABCG1 each make complementary and important contributions to β -cell function by maintaining islet cholesterol homeostasis in vivo. *Diabetes* 61:659–664, 2012

Type 2 diabetes is characterized by both progressive β -cell dysfunction and loss of β -cell mass. Recently, abnormalities of cholesterol metabolism have emerged as a potential contributor to β -cell dysfunction (1,2). In humans, low levels of HDL, which is the predominant acceptor of cellular cholesterol, is a risk factor for the development of type 2 diabetes (3) and is associated with β -cell dysfunction (4). Although multiple mechanisms may explain the protective properties of HDL on β -cell dysfunction, enhancement of cellular

cholesterol efflux seems to play a crucial role. Cholesterol accumulation in islets compromises β -cell function and reduces insulin secretion in mice (5–8). In vitro data indicate that HDL-mediated increase in insulin secretion is dependent on the ATP-binding cassette (ABC) transporters ABCA1 and ABCG1 (9). We previously demonstrated that ABCA1 plays a critical role in β -cell cholesterol homeostasis and β -cell function in mice (5). Deletion of ABCA1 specifically in β -cells leads to markedly impaired glucose tolerance, defective insulin secretion, and cholesterol accumulation in islets (5). Islets lacking β -cell ABCA1 have increased expression of the related cholesterol transporter ABCG1, perhaps as a compensatory mechanism to maintain islet cholesterol homeostasis.

ABCG1 promotes cholesterol efflux to HDL and acts sequentially with ABCA1 to remove cellular cholesterol (10). Mice with deletion of ABCG1 also have impaired glucose-induced insulin secretion (11). ABCA1 and ABCG1 have complementary roles in macrophage function (12,13). In addition to massive lipid accumulation, loss of both ABCA1 and ABCG1 in macrophages leads to increased expression of proinflammatory cytokines and enhanced susceptibility to apoptosis (13). However, the relative importance of ABCA1 and ABCG1 in islets is thus far unknown.

In this study, we examined whether ABCA1 and ABCG1 have a complementary role in mediating cholesterol efflux in β -cells. To determine whether loss of both cholesterol transporters induces an exacerbated phenotype compared with loss of either transporter alone, we assessed cholesterol accumulation, glucose tolerance, insulin secretion, and islet inflammation in mice with β -cells deficient in both ABCA1 and ABCG1.

RESEARCH DESIGN AND METHODS

Animals. ABCA1^{fl/fl};Rip-Cre mice (5) and ABCG1^{-/-} mice (Deltagen, San Mateo, CA), both on pure C57Bl/6 backgrounds, were crossed to generate F1 heterozygotes. Heterozygote F1 animals were crossbred to obtain the following mice: ABCA1^{fl/fl};ABCG1^{+/+} (designated in text as “control”), ABCA1^{fl/fl};Rip-Cre;ABCG1^{+/+} (ABCA1^{-P/-P}), ABCA1^{fl/fl};ABCG1^{-/-} (ABCG1^{-/-}), and ABCA1^{fl/fl};Rip-Cre;ABCG1^{-/-} (ABCA1^{-P/-P};ABCG1^{-/-}) mice. We have previously reported that under our experimental conditions, mice with floxed ABCA1 alleles (ABCA1^{fl/fl}) or mice transgenic for the RIP-Cre transgene do not have altered glucose homeostasis compared with ABCA1^{+/+} mice (5). All mice were 3–4 months of age. All studies were approved by the University of British Columbia Animal Care Committee.

Physiological and metabolic studies. Intraperitoneal glucose tolerance tests (GTTs) were performed on 4 h-fasted mice injected with 2 g/kg glucose as previously described (5). Insulin secretion during static incubation and perfusion were performed on hand-picked islets isolated after intraductal collagenase injection (5,7). Plasma cholesterol was determined by enzymatic assay (Thermo Electron Corporation). Islet cholesterol, desmosterol, lathosterol, cholesterol, campesterol, and sitosterol were determined by gas-liquid chromatography-mass spectrometry (14).

From the ¹Department of Medical Genetics, Centre for Molecular Medicine and Therapeutics, University of British Columbia, Vancouver, British Columbia, Canada; the ²Department of Pathology and Laboratory Medicine, Child and Family Research Institute, University of British Columbia, Vancouver, British Columbia, Canada; the ³Laboratory for Special Lipid Diagnostics, Institute of Clinical Chemistry and Clinical Pharmacology, University Clinics of Bonn, Bonn, Germany; the ⁴Department of Cellular and Physiological Sciences, University of British Columbia, Vancouver, British Columbia, Canada; and the ⁵Departments of Surgery and Pathology and Laboratory Medicine, Child and Family Research Institute, University of British Columbia, Vancouver, British Columbia, Canada.

Corresponding author: Michael R. Hayden, mrh@cmmt.ubc.ca.

Received 22 September 2011 and accepted 11 December 2011.

DOI: 10.2337/db11-1341

© 2012 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.




FIG. 1. Loss of β-cell ABCA1 and ABCG1 exacerbates glucose intolerance and β-cell dysfunction. **A:** Body weight of 3-month-old mice ($n = 8-12$ per group). **B:** Fasted plasma glucose levels ($n = 10-14$ per group). **C:** Plasma glucose levels during intraperitoneal GTT ($n = 5-9$). Values represent mean \pm SEM. **D:** Area under curve of GTT. **E:** Plasma glucose levels during intraperitoneal insulin tolerance test ($n = 5-9$). Values represent mean \pm SEM. **F:** Insulin secretion from isolated islets during static incubation. Islets were cultured overnight and then stimulated for 1 h in the conditions indicated. Values represent pooled data from three separate experiments, each consisting of pooled islets from two mice per genotype. **G:** Insulin release during islet perfusion experiments ($n = 3$ per group). Values represent mean \pm SEM. **H:** Area under the curve of insulin release during islet perfusion experiments. ^a $P < 0.05$ versus control; ^b $P < 0.05$ versus ABCA1^{-P/-P}; ^c $P < 0.05$ versus ABCG1^{-/-}.

Quantitative PCR. Real-time PCR for C/EBP homologous protein (CHOP), interleukin-1 β (IL-1 β), and F4/80 was performed as previously described (5). Primer sequences are available upon request.

β -Cell mass and immunofluorescence. β -Cell mass was determined from five evenly spaced, paraffin-embedded pancreatic sections as previously described (5). Macrophage staining was performed with antibodies to F4/80 (Cedarlane) and insulin (DAKO) and quantified using ImagePro software (MediaCybernetics). All islets in three sections from different areas of the pancreas were analyzed.

Statistical analysis. Differences between groups were calculated by Kruskal-Wallis test with Conover post-test (for four groups) with a *P* value of 0.05 considered significant.

RESULTS

Loss of β -cell ABCA1 and ABCG1 exacerbates glucose intolerance and β -cell dysfunction. In order to determine the effects of combined deficiency of ABCA1 and ABCG1 on β -cell function, we crossed mice with β -cell-specific ABCA1 deficiency (ABCA1^{-P/-P}) with globally deficient ABCG1 (ABCG1^{-/-}) mice to generate ABCA1^{-P/-P}; ABCG1^{-/-} double knockout animals. Double knockout mice showed no changes in body weight (Fig. 1A), but had increased fasting glucose levels (Fig. 1B) compared with single knockout mice, which had similar fasting glucose levels as control animals. Lack of ABCA1 specifically in β -cells resulted in glucose intolerance (Fig. 1C and D), as previously reported (5). Although deletion of ABCG1 alone had no significant effect on glucose tolerance, combined deletion of ABCG1 and ABCA1 greatly exacerbated glucose intolerance compared with either ABCG1^{-/-} or ABCA1^{-P/-P} mice alone. Insulin sensitivity was similar in all groups (Fig. 1E).

As reported previously (5), islets lacking β -cell ABCA1 showed decreased glucose-induced insulin secretion during static incubation (Fig. 1F), as well as perfusion experiments (Fig. 1G and H). Both first and second-phase, glucose-stimulated insulin secretion were reduced. Glucose-stimulated insulin secretion was significantly but less markedly decreased in ABCG1^{-/-} mice (Fig. 1F). Notably, islets lacking both ABCA1 and ABCG1 had an even greater reduction in glucose-stimulated insulin secretion compared with islets from mice lacking either transporter (Fig. 1F, G, and H).

Loss of β -cell ABCA1 and ABCG1 exacerbates islet sterol accumulation. As loss of both ABCA1 and ABCG1 leads to massive sterol accumulation in several tissues (15), we measured sterol levels in isolated islets. In agreement with our previous findings (5), loss of β -cell ABCA1 resulted in significant islet cholesterol accumulation (Table 1). Although ABCG1 deletion alone had no effect on cholesterol accumulation, combined loss of both transporters resulted in increased islet cholesterol levels compared with control, ABCA1^{-P/-P}, and ABCG1^{-/-} islets (Table 1). Other sterols,

such as desmosterol, cholestanol, campesterol, and sitosterol, were also increased in islets lacking both transporters (Table 1). Importantly, plasma cholesterol levels were similar in all groups (control mice, 2.26 \pm 0.08 mmol/L; ABCA1^{-P/-P} mice, 2.37 \pm 0.07 mmol/L; ABCG1^{-/-} mice, 2.36 \pm 0.12 mmol/L; ABCA1^{-P/-P}; ABCG1^{-/-} mice, 2.19 \pm 0.23 mmol/L).

Increased CHOP expression, but no differences in β -cell mass, in islets lacking ABCA1 and ABCG1. Cholesterol accumulation in macrophages leads to activation of the unfolded protein response and CHOP-induced apoptosis (16). CHOP is an unfolded protein response-induced transcription factor that links endoplasmic reticulum stress to β -cell dysfunction and apoptosis in animal models of type 2 diabetes (17). We found that β -cell-specific deletion of ABCA1 caused a significant increase in CHOP expression in isolated islets, whereas ABCG1 deficiency had no effect (Fig. 2A). Lack of both transporters led to a further increase in CHOP expression (Fig. 2A), but this was not associated with any change in β -cell mass at the time point studied (Fig. 2B).

Loss of both β -cell ABCA1 and ABCG1 leads to macrophage infiltration and increased IL-1 β expression. Islet inflammation is emerging as an important contributor to type 2 diabetes (18). Recent studies have shown that both ABCA1 and ABCG1 modulate inflammation (19). Thus, we examined mRNA levels of IL-1 β , a proinflammatory cytokine that plays a central role in modulating islet chemokine release and impairs islet function (18). IL-1 β mRNA was significantly increased in islets lacking both ABCA1 and ABCG1, but not in islets lacking either ABCA1 or ABCG1 alone (Fig. 3A). To assess islet macrophage infiltration, we analyzed expression of the macrophage marker F4/80 in islets. Both ABCA1^{-P/-P} and ABCA1^{-P/-P}; ABCG1^{-/-} islets were found to have increased mRNA levels of the macrophage marker F4/80 (Fig. 3B). In support of these data, increased numbers of F4/80-positive cells were observed, by immunostaining, to be present in islets of ABCA1^{-P/-P} and ABCA1^{-P/-P}; ABCG1^{-/-} mice (Fig. 3C).

DISCUSSION

β -Cell cholesterol homeostasis is an emerging factor that has been shown to influence β -cell function and insulin secretion (1,2). Islet cholesterol accumulation due to impaired cholesterol efflux or hypercholesterolemia leads to decreased insulin secretion (5–8). The current study shows that two major cellular cholesterol efflux transporters, ABCA1 and ABCG1, play complementary roles in mediating cholesterol efflux from β -cells and protecting against islet sterol accumulation and resultant impairment of insulin secretion and islet inflammation.

TABLE 1

Islet sterol levels are increased in ABCA1^{-P/-P}; ABCG1^{-/-} mice

	Control	ABCA1 ^{-P/-P}	ABCG1 ^{-/-}	ABCA1 ^{-P/-P} ; ABCG1 ^{-/-}
Cholesterol (μ g/mg protein)	33.2 \pm 3.5	52.8 \pm 2.9 ^{a,c}	38.3 \pm 5.3	118.8 \pm 4.8 ^{a,b,c}
Lathosterol (ng/mg protein)	18 \pm 4	8 \pm 3	15 \pm 4	10 \pm 2
Desmosterol (ng/mg protein)	249 \pm 50	268 \pm 48	151 \pm 44	466 \pm 36 ^{a,b,c}
Cholestanol (ng/mg protein)	463 \pm 47	636 \pm 36 ^{a,c}	481 \pm 55	1,760 \pm 200 ^{a,b,c}
Campesterol (ng/mg protein)	400 \pm 43	543 \pm 76	478 \pm 78	1,876 \pm 325 ^{a,b,c}
Sitosterol (ng/mg protein)	428 \pm 71	520 \pm 30	536 \pm 72	990 \pm 253 ^{a,b,c}

Data are mean \pm SEM. ^a*P* < 0.05 versus control. ^b*P* < 0.05 versus ABCA1^{-P/-P}. ^c*P* < 0.05 versus ABCG1^{-/-} (*n* = 4–6 in each group).




FIG. 2. Increased CHOP expression, but normal β-cell mass in ABCA1^{-P/-P} and ABCA1^{-P/-P};ABCG1^{-/-} islets. **A:** Relative CHOP mRNA levels in isolated islets (*n* = 5–7 per group). **B:** β-Cell mass (*n* = 4–8 per group). ^a*P* < 0.05 versus control; ^b*P* < 0.05 versus ABCA1^{-P/-P}; ^c*P* < 0.05 versus ABCG1^{-/-}.

The dramatic sterol accumulation in islets lacking both ABCA1 and ABCG1 confirmed the important role of these transporters in lipid metabolism in β-cells. Importantly, the degree of sterol accumulation, as well as glucose intolerance in mice deficient for both transporters, was more than additive, indicating that these two transporters have a synergistic effect on islet sterol homeostasis. Islets lacking ABCA1 and ABCG1 not only accumulate cholesterol, the cholesterol metabolite cholestanol, and the cholesterol precursor desmosterol, but also plant sterols such as campesterol and sitosterol, underlining the importance of these transporters in sterol efflux. Previously, it was suggested that ABCG1 does not mediate cholesterol efflux in β-cells (11), as ABCG1 deficiency did not impact cholesterol levels or cholesterol efflux in β-cells. Cholesterol levels in insulin vesicles was decreased in ABCG1 knock-down β-cells, which led to the suggestion that ABCG1 in β-cells primarily regulates subcellular cholesterol distribution (11). Although we similarly observed that ABCG1 deficiency alone does not result in islet cholesterol accumulation, a role for ABCG1 in β-cell cholesterol export was unmasked in the absence of β-cell ABCA1.

In the current study, we used β-cell-specific ABCA1-deficient mice crossed with the global ABCG1^{-/-} mice to




FIG. 3. Loss of β-cell ABCA1 and ABCG1 results in islet inflammation. **A:** Relative IL-1β mRNA levels in isolated islets (*n* = 5–7 per group). **B:** Relative mRNA levels of the macrophage marker F4/80 in isolated islets (*n* = 5–7 per group). **C:** Number of F4/80-positive cells per insulin-positive area in pancreas isolated from control, ABCA1^{-P/-P}, ABCG1^{-/-}, and ABCA1^{-P/-P};ABCG1^{-/-} mice (*n* = 4–6 per group). ^a*P* < 0.05 versus control; ^b*P* < 0.05 versus ABCA1^{-P/-P}; ^c*P* < 0.05 versus ABCG1^{-/-}.

determine whether ABCA1 and ABCG1 have complementary roles in β -cells. We chose not to use mice with global ABCA1 deficiency that have very low plasma cholesterol levels (20), as we previously showed that β -cells lacking ABCA1 are susceptible to dysfunction in vivo only in the presence of sufficient levels of plasma cholesterol (5). An obvious limitation of the current study is that the loss of ABCG1 is not β -cell specific and we can therefore not exclude effects of ABCG1 deficiency in other cell types mediating the phenotype reported here.

Absence of both ABCA1 and ABCG1 in β -cells resulted in no overt alteration in β -cell mass in 4-month-old mice. In contrast, glucose-stimulated insulin secretion was further impaired in islets lacking both transporters, indicating that impaired β -cell function and not β -cell loss resulted in the hyperglycemia in the double knockout mice. It remains to be determined whether β -cell mass will be diminished in older mice, since we did notice a marked increase in expression of the endoplasmic reticulum stress-related apoptosis factor CHOP in islets lacking both ABCA1 and ABCG1. In keeping with our previous findings (5), β -cell dysfunction was characterized by loss of both glucose- and KCl-stimulated insulin secretion, consistent with a defect in granule exocytosis associated with cholesterol accumulation (21).

Loss of both ABCA1 and ABCG1 resulted in increased IL-1 β expression and increased macrophage infiltration in islets. Islet inflammation has been reported in type 2 diabetes (22). Elevated glucose and free fatty acids increase expression of proinflammatory cytokines in islets (22). Our results suggest that decreased activity of ABCA1 and ABCG1 in β -cells could also lead to islet inflammation. Loss of both transporters in macrophages results in an exaggerated cytokine response to toll-like receptor (TLR) stimuli (19). Although the precise mechanism is unknown, recent data suggest that macrophages lacking ABCA1 exhibit increased trafficking of TLRs to lipid rafts, which leads to enhanced signaling (23). The IL-1 receptor, which appears to be critical for inflammatory responses in β -cells (18), also depends on lipid rafts for signaling (24). Therefore, lack of both ABCA1 and ABCG1 in β -cells could potentially lead to enhanced TLR or IL-1 receptor signaling, resulting in islet inflammation. Furthermore, cholesterol accumulation could lead to the formation of cholesterol crystals, which have been shown to induce inflammation by stimulating the NLRP3 inflammasome in macrophages (25). Our data provide an important link between islet cholesterol accumulation and islet inflammation, which both potentially contribute to increased islet dysfunction in the absence of ABCA1 and ABCG1.

In summary, our data show that ABCA1 and ABCG1 have complementary roles in protecting against islet cholesterol accumulation, inflammation and impaired insulin secretion. These data add further support to the concept that regulation of β -cell cholesterol homeostasis is essential for normal islet function, and suggest that upregulating the activity of ABCA1 and ABCG1 could be a promising approach to decrease islet cholesterol accumulation, decrease islet inflammation, and improve β -cell function in type 2 diabetes.

ACKNOWLEDGMENTS

J.K.K. was supported by postdoctoral fellowship awards from the Canadian Institutes of Health Research (CIHR) and the Michael Smith Foundation for Health Research

(MSFHR). N.W. was supported by a postdoctoral fellowship from the Heart and Stroke Foundation of Canada. C.W.-R. is a recipient of a Vanier Canada Graduate Scholarship and an MSFHR Junior Graduate Studentship. W.d.H. was supported by postdoctoral fellowship awards from the Canadian Diabetes Association and MSFHR. L.R.B. received research support from the University of British Columbia Department of Medicine. C.B.V. is an MSFHR Senior Scholar. M.R.H. holds a Canada Research Chair in Human Genetics and is a University of British Columbia Killam Professor. This work was supported by CIHR grants to M.R.H. and C.B.V.

No potential conflicts of interest related to this article were reported.

J.K.K. designed and performed the research and wrote the manuscript. N.W., C.W.-R., and W.d.H. performed the research, contributed to discussion, and reviewed and edited the manuscript. T.V., A.B., and R.T. performed the research. C.L.W., D.L., J.D.J., and L.R.B. contributed to discussion and reviewed and edited the manuscript. C.B.V. designed the research, contributed to discussion, and reviewed and edited the manuscript. M.R.H. designed the research and wrote the manuscript. M.R.H. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

The authors would like to thank Piers Ruddle, Mark Wang, and Xiaoke Hu (University of British Columbia) for technical assistance; John S. Parks (Wake Forest University School of Medicine, Winston-Salem, NC) for providing the ABCA1^{fl/fl} mice; Jan A. Ehses (University of British Columbia) for insightful comments, and Vincent W. Bloks (University Medical Center Groningen, Groningen, the Netherlands) for statistical advice.

REFERENCES

- Brunham LR, Kruit JK, Verchere CB, Hayden MR. Cholesterol in islet dysfunction and type 2 diabetes. *J Clin Invest* 2008;118:403–408
- Kruit JK, Brunham LR, Verchere CB, Hayden MR. HDL and LDL cholesterol significantly influence beta-cell function in type 2 diabetes mellitus. *Curr Opin Lipidol* 2010;21:178–185
- von Eckardstein A, Schulte H, Assmann G. Risk for diabetes mellitus in middle-aged Caucasian male participants of the PROCAM study: implications for the definition of impaired fasting glucose by the American Diabetes Association. *Prospective Cardiovascular Münster. J Clin Endocrinol Metab* 2000;85:3101–3108
- Hermans MP, Ahn SA, Rousseau MF. log(TG)/HDL-C is related to both residual cardiometabolic risk and β -cell function loss in type 2 diabetes males. *Cardiovasc Diabetol* 2010;9:88
- Brunham LR, Kruit JK, Pape TD, et al. Beta-cell ABCA1 influences insulin secretion, glucose homeostasis and response to thiazolidinedione treatment. *Nat Med* 2007;13:340–347
- Hao M, Head WS, Gunawardana SC, Hasty AH, Piston DW. Direct effect of cholesterol on insulin secretion: a novel mechanism for pancreatic beta-cell dysfunction. *Diabetes* 2007;56:2328–2338
- Kruit JK, Kremer PH, Dai L, et al. Cholesterol efflux via ATP-binding cassette transporter A1 (ABCA1) and cholesterol uptake via the LDL receptor influences cholesterol-induced impairment of beta cell function in mice. *Diabetologia* 2010;53:1110–1119
- Peyot ML, Pepin E, Lamontagne J, et al. Beta-cell failure in diet-induced obese mice stratified according to body weight gain: secretory dysfunction and altered islet lipid metabolism without steatosis or reduced beta-cell mass. *Diabetes* 2010;59:2178–2187
- Fryirs MA, Barter PJ, Appavoo M, et al. Effects of high-density lipoproteins on pancreatic beta-cell insulin secretion. *Arterioscler Thromb Vasc Biol* 2010;30:1642–1648
- Gelissen IC, Harris M, Rye KA, et al. ABCA1 and ABCG1 synergize to mediate cholesterol export to apoA-I. *Arterioscler Thromb Vasc Biol* 2006;26:534–540
- Sturek JM, Castle JD, Trace AP, et al. An intracellular role for ABCG1-mediated cholesterol transport in the regulated secretory pathway of mouse pancreatic beta cells. *J Clin Invest* 2010;120:2575–2589

12. Out R, Hoekstra M, Habets K, et al. Combined deletion of macrophage ABCA1 and ABCG1 leads to massive lipid accumulation in tissue macrophages and distinct atherosclerosis at relatively low plasma cholesterol levels. *Arterioscler Thromb Vasc Biol* 2008;28:258–264
13. Yvan-Charvet L, Ranalletta M, Wang N, et al. Combined deficiency of ABCA1 and ABCG1 promotes foam cell accumulation and accelerates atherosclerosis in mice. *J Clin Invest* 2007;117:3900–3908
14. Lütjohann D, Stroick M, Bertsch T, et al. High doses of simvastatin, pravastatin, and cholesterol reduce brain cholesterol synthesis in guinea pigs. *Steroids* 2004;69:431–438
15. Out R, Jessup W, Le Goff W, et al. Coexistence of foam cells and hypocholesterolemia in mice lacking the ABC transporters A1 and G1. *Circ Res* 2008;102:113–120
16. Feng B, Yao PM, Li Y, et al. The endoplasmic reticulum is the site of cholesterol-induced cytotoxicity in macrophages. *Nat Cell Biol* 2003;5:781–792
17. Song B, Scheuner D, Ron D, Pennathur S, Kaufman RJ. Chop deletion reduces oxidative stress, improves beta cell function, and promotes cell survival in multiple mouse models of diabetes. *J Clin Invest* 2008;118:3378–3389
18. Donath MY, Böni-Schnetzler M, Ellingsgaard H, Ehses JA. Islet inflammation impairs the pancreatic beta-cell in type 2 diabetes. *Physiology (Bethesda)* 2009;24:325–331
19. Yvan-Charvet L, Wang N, Tall A. Role of HDL, ABCA1, and ABCG1 transporters in cholesterol efflux and immune responses. *Arterioscler Thromb Vasc Biol* 2010;30:139–143
20. McNeish J, Aiello RJ, Guyot D, et al. High density lipoprotein deficiency and foam cell accumulation in mice with targeted disruption of ATP-binding cassette transporter-1. *Proc Natl Acad Sci USA* 2000;97:4245–4250
21. Kruit JK, Wijesekara N, Fox JE, et al. Islet cholesterol accumulation due to loss of ABCA1 leads to impaired exocytosis of insulin granules. *Diabetes* 2011;60:3186–3196
22. Ehses JA, Perren A, Eppler E, et al. Increased number of islet-associated macrophages in type 2 diabetes. *Diabetes* 2007;56:2356–2370
23. Zhu X, Owen JS, Wilson MD, et al. Macrophage ABCA1 reduces MyD88-dependent Toll-like receptor trafficking to lipid rafts by reduction of lipid raft cholesterol. *J Lipid Res* 2010;51:3196–3206
24. Oakley FD, Smith RL, Engelhardt JF. Lipid rafts and caveolin-1 coordinate interleukin-1beta (IL-1beta)-dependent activation of NFkappaB by controlling endocytosis of Nox2 and IL-1beta receptor 1 from the plasma membrane. *J Biol Chem* 2009;284:33255–33264
25. Duewell P, Kono H, Rayner KJ, et al. NLRP3 inflammasomes are required for atherogenesis and activated by cholesterol crystals. *Nature* 2010;464:1357–1361